

3 1761 06705692 9



Presented to
The Library
of the
University of Toronto
by

R. A. Paterson, Esq.

R. A. Peterson
Hous. Coll.
Toronto

1884

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

Cambridge
School and College
Text Books.

Cambridge :

**PRINTED BY C. J. CLAY & SON,
AT THE UNIVERSITY PRESS.**

Physics
Mech
B

CAMBRIDGE SCHOOL AND COLLEGE
TEXT BOOKS.

ELEMENTARY
HYDROSTATICS

BY
W. H. BESANT, M.A., F.R.S.

LECTURER OF ST JOHN'S COLLEGE, CAMBRIDGE.

266962
4.4.32

TENTH EDITION REVISED.

CAMBRIDGE:
DEIGHTON, BELL, AND CO.
LONDON: GEORGE BELL AND SONS.

1882



P R E F A C E.

I HAVE endeavoured in the following treatise to place before the student a complete series of those propositions in Hydrostatics, the solution of which can be effected without the aid of the Differential Calculus, and to illustrate the theory by the description of many Hydrostatic Instruments, and by the insertion of a large number of examples and problems.

In doing this I have had in view the courses of preparation necessary for the first three days of the Examination for the Mathematical Tripos, for some of the Examinations of the University of London, and for various other Examinations in which more or less knowledge of Hydrostatics is required.

As far as possible the whole of the propositions are strictly deduced from the definitions and axioms of the subject, but it is occasionally necessary to assume empirical results, and these assumptions are distinctly pointed out. I have thought it advisable to give a slight account of some cases of fluid motion, and also to give an explanation of some of the more important phenomena of sound; in each of these cases I have assumed, as the basis of reasoning, certain facts which can be deduced from theory by an analytical investigation, but which it may be useful to the student to accept as experimental results.

The Geometrical facts which are enunciated at the end of the Introduction are such as can be demonstrated without the aid of the Differential Calculus.

By Professor Miller's kind permission, I have been allowed to make use of the Chapter on Instruments in his Hydrostatics: of this permission I have availed myself in many cases, and, in particular, I am entirely indebted to Professor Miller for the descriptions of the Piezometer and Stereometer, and for information and references having regard to those instruments.

The slight historical notices appended to some of the Chapters are intended to mark the principal steps in the progress of the science, and to assign to their respective authors the exact values of the advances made at different times.

I have given, in most cases, the answers to the examples and problems, and these will, I hope, sufficiently illustrate the subject, and form for the student a collection of useful and instructive exercises.

W. H. BESANT.

ST JOHN'S COLLEGE,
April, 1863.

PREFACE TO THE TENTH EDITION.

IN the present edition the text has been carefully revised, a chapter on Capillarity has been inserted, and other additions have been made. I venture to hope that these additions will be an aid to the Student and will increase the utility of the treatise as a Textbook.

W. H. BESANT.

ST JOHN'S COLLEGE,
April 1882.

TABLE OF CONTENTS.

		PAGE
	INTRODUCTION	1
CHAPTER I.		
ARTICLES		
1—18.	Definitions and Fundamental Properties of Fluids, Hydrostatic Paradox, Hydraulic Presses	4
CHAPTER II.		
19—31.	Density and Specific Gravity	17
CHAPTER III.		
32—50.	Pressures at different points of a fluid, Pressures on plane surfaces, Whole Pressure, Centre of Pressure	25
CHAPTER IV.		
51—67.	Resultant horizontal and vertical pressures, Equilibrium of a floating body, stability, Metacentre	45
CHAPTER V.		
68—94.	Mechanical Properties of Air and Gases, Thermometers, Barometer, Siphon, Differential Thermometer	69

CHAPTER VI.

ARTICLES

PAGE

- 95—121. The Diving Bell, Pumps, Bramah's Press,
Air Pumps, Condenser, Manometers,
Barker's Mill, Piezometer, Hydraulic
Ram, Steam Engines 94

CHAPTER VII.

- 122—132. The determination of Specific Gravities,
Hydrostatic Balance, Hydrometers, Ste-
reometer 123

CHAPTER VIII.

- 133—157. Mixture of Gases, Radiation, Conduction and
Convection of Heat, Dew, Hygrometers,
Effects of Heat and Cold on liquids, Spe-
cific Heat 134

CHAPTER IX.

- 158—164. On Rotating Liquid 150

CHAPTER X.

- 165—170. On the Tension of cylindrical and spherical
vessels containing fluids 161

CHAPTER XI.

- 171—177. On Capillarity 168

CHAPTER XII.

- 178—183. The Motion of Fluids 173

CHAPTER XIII.

- 184—200. On Sound 179

- Appendix I. 197

- „ II. Solutions of Problems 201

- Miscellaneous Problems 216

- Answers to Examples and Problems 215

- Index 224

ELEMENTARY HYDROSTATICS.

INTRODUCTION.

THE object of the science of Hydrostatics is to discuss the mechanical properties of fluids, or to determine the nature of the action which fluids exert upon each other and upon bodies with which they are in contact, and to explain and classify, under general laws, the varied phenomena relating to fluids which are offered to the attention of an observer. To effect this purpose it is necessary to construct a consistent theory, founded upon observation and experiment, from which, by processes of deductive reasoning, and the aid of Geometry and Algebra, the explanations of phenomena shall flow as consequences of the definitions and fundamental properties assumed; the test of the theory will be the coincidence with observed facts of the results of such reasoning.

We shall assume in the following pages that the student is acquainted with the elements of Plane Geometry, with the simpler portions of Algebra and Trigonometry, and of Statics, and, in the later chapters, with a few of the properties of Conic Sections, and certain results of Dynamics.

In dealing with any mechanical science, we may take as the basis of our reasoning certain known laws, derived from experiment, or we may deduce these laws from a set of

axioms and definitions, the axioms being the result of inductive reasonings from observed facts. With our present subject it is generally necessary to rest upon empirical laws, but in some cases these laws can be deduced from the axiomatic definition of a fluid. For instance, in the first chapter we have stated as experimental laws the principles of the equality of pressure in all directions and the transmission of pressure, but this formal statement of fact is followed by a deduction of the laws, by strict reasoning, from the axiomatic definition.

The idea of a varying fluid pressure and of the measure of such pressure is one of the first which presents itself as a difficulty; the student will perceive that it is a difficulty of the same kind as the idea of varying velocity and its measure. A body in motion with a changing velocity has, at any instant, a rate of motion which can be appreciated and measured; and, in a similar manner, the pressure at any point of a fluid can be conceived, and, by reference to proper units, can be made the subject of calculation.

In problems relating to the equilibrium of fluids, an artificial mode of thought enables us to reduce such problems to the form of statical problems, and we are thus enabled to employ the laws of equilibrium, which have been proved for rigid bodies.

Some of the most important results of the science will be found in the construction of Hydrostatic instruments; a consideration of these instruments, many of which we shall describe, will shew how universal are the practical applications of fluids, and that, while doing the hardest work of levers and pulleys, they at the same time assist in the most delicate manipulations for determining weights and measures. The Hydraulic Press and the Stereometer illustrate these extreme applications of the properties of fluids.

The articles printed in smaller characters in the following chapters may if necessary be omitted during a first reading of the subject, and the Examination papers which follow the first eight chapters are intended as a first course of questions upon the chapters. The examples which follow the Examination papers are somewhat more difficult, and should be dealt with after the former have been studied and discussed.

The following geometrical facts are assumed and employed in some of the examples.

The volume of a pyramid or of a cone is one-third of the prism or cylinder on the same base and of the same altitude.

The volume of a sphere is $\frac{4}{3}\pi r^3$, and its surface is $4\pi r^2$, r being the radius.

The volume of a paraboloid of revolution is one-half the cylinder on the same base and of the same altitude.

The surface of a cone is $\pi r^2 \operatorname{cosec} \alpha$, r being the radius of the base and α the semivertical angle. This may also be written $\pi r \sqrt{r^2 + h^2}$, h being the altitude of the cone.

The area of an ellipse is πab , $2a$ and $2b$ being the lengths of its axes.

The area of the portion of a parabola cut off by any ordinate is two-thirds of the rectangle, the sides of which are the ordinate and corresponding abscissa.

It is also assumed that *the weight of a cubic foot of water is 1000 oz.*

CHAPTER I.

Definition of a Fluid, Compressibility of Liquids, Fluid Pressure, Transmission of Pressure, Equality of Pressure in all directions, Hydrostatic Bellows, Hydrostatic Paradox, Hydraulic Presses, and Safety Valves.

1. **I**T is a matter of ordinary observation that fluids are capable of exerting pressure.

A certain amount of effort is necessary in order to immerse the hand in water, and the effort is much more sensible when a light substance, such as a piece of wood or cork, is held under water, the resistance offered to the immersion being greater as the piece immersed is larger. This resistance can only be caused by the fluid pressure acting upon the surface of the body immersed.

If an aperture be made in the side of a vessel containing water, and be covered by a plate so as to prevent the escape of the water, a definite amount of force must be exerted in order to maintain the plate in its position, and this force is opposed to, and is a direct measure of, the pressure of the water.

That the atmosphere when at rest exerts pressure is shewn directly by means of an air-pump. Amongst many experiments a simple one is to exhaust the air within a receiver made of very thin glass; when the exhaustion has reached a certain point depending on the strength of the glass, the receiver will be shivered by the pressure of the external air. The action of wind, the motion of a wind-mill, the propulsion of a boat by means of sails, and other familiar facts offer themselves naturally as instances of the pressure of the air when in motion.

2. All such substances as water, oil, mercury, steam, air, or any kind of gas are called fluids, but in order to obtain a definition of a fluid, we have to find a property which is common to all these different kinds of substances, and which does not depend upon any of the characteristics by which they are distinguished from each other. This property is found in the extreme mobility of their particles and in the ease with which these particles can be separated from the mass of fluid and from each other, no sensible resistance being offered to the separation from a mass of fluid of a portion whether large or small.

If a very thin plate be immersed in water, the resistance to its immersion in the direction of its plane is so small as to lead to the idea that a perfectly fluid mass is incapable of exerting any tangential action, or, in other words, any action of the nature of friction, such for instance as would be exerted if the plate were pushed between two flat boards held close to each other. Observations of such experiments have led to the following definition :

A Fluid is a substance, such that a mass of it can be very easily divided in any direction, and of which portions, however small, can be very easily separated from the whole mass ;

And also to the statement of the fundamental property of a fluid, viz. :

The Pressure of a fluid on any surface with which it is in contact is perpendicular to the surface.

3. Fluids are of two kinds, liquid and gaseous, the former being practically incompressible, while the latter, by the application of ordinary force, can be easily compressed, and, if the compressing force be removed or diminished, will expand in volume.

Liquids are however really compressible, but to a slight degree.

Experiments made by Canton in 1761, Perkins in 1819, Oersted in 1823, Colladon and Sturm in 1829, and others, have proved the compressibility of liquids.

The last two obtained the following results, employing a pressure of one atmosphere, that is $14\frac{1}{2}$ lbs. on a square inch, at the temperature 0° .

Compression of unit of volume.

Mercury000005
Distilled water000049
” ” deprived of air000051
Sulphuric ether000133

Moreover *the decrease in volume, for the same liquid, is proportional to the pressure.*

If V be the original volume of a liquid, and V' its volume under a pressure p , $V - V'$ is the decrease in the volume V , and therefore $\frac{V - V'}{V}$ is the decrease in each unit of volume.

Hence the law may be thus stated :

$$\frac{V - V'}{V} \propto p = \mu p,$$


where μ is different for different fluids.

Thus for mercury, if p be measured by taking one atmospheric pressure as the unit, we have $\mu = .000005$. We shall however, in all questions relating to equilibrium, consider liquids as incompressible fluids.

Measure of fluid pressure.

4. The pressure of a fluid on a plane is measured, when uniform over the plane, by the force exerted on an unit of area.

Thus, if a vessel with a moveable base contain water, and if it be necessary to employ a force of 60 lbs. upwards to keep the base at rest, then 60 lbs. is the pressure of the water on the base; and, supposing the area of the base to be 4 square inches, and that a square inch is the unit of area, the measure of the pressure *at any point of the base* is 15 lbs.



The pressure *on* a point of the base is of course zero; the pressure *at* a point is used conventionally to express the pressure on a square unit containing the point.

If the pressure be variable over the plane, as, for instance, on the vertical side of a vessel, the pressure at any point is measured by the pressure which would be exerted on an unit of area, supposing the pressure over the whole unit to be exerted at the same rate as it is at the point.

In order to measure the pressure of a fluid at any point *within its mass*, imagine a small rigid plane placed so as to contain the point, and conceive the fluid removed from one side of the plane and the plane kept at rest by a force of P lbs. Then if a be the area of the plane, and the pressure over it be uniform, $\frac{P}{a}$ is the pressure on each unit of area, and this is usually represented by p .


If the pressure over the plane be variable, we may suppose the area a made so small that the pressure shall be sensibly uniform, and in this case P will be small as well as a , but $\frac{P}{a}$ or p will measure the rate of pressure at the point*.

Or we may say that $\frac{P}{a}$ is the measure of the mean pressure over the area a , and that, when a is small, this mean pressure is the actual pressure.

5. *Transmission of fluid pressure.*


Any pressure, applied to the surface of a fluid, is transmitted equally to all parts of the fluid.

If a closed vessel be filled with water, and if A and B be two equal openings in the top of the vessel, closed by pistons, it is found that any pressure applied at A must be counteracted by an equal pressure at B to prevent its being forced out, and if C be a piston of different size, it is found that the pressure applied at C must bear to the pressure on A the ratio of the area of C to that of A , and that this is the case whether the piston B exists or not.



* This may be expressed by saying that p is the ultimate value of $\frac{P}{a}$ when a , and therefore P , are indefinitely diminished.

Taking a more general case, if a vessel of any shape have several openings closed by pistons, kept at rest by suitable forces, it will be found that any additional force P applied to one piston will require the application, to all the other pistons, of additional forces, which have the same ratio to P as the areas of the respective pistons have to that of the piston to which P is applied.



6. To explain the reason of this equal transmission, imagine a tube of uniform bore filled with water and closed by pistons at A and B . Then it may be assumed as self-



evident, that any additional force applied at A will require an equal additional force at B to counteract it and keep the fluid at rest.

Now suppose in the figure that A and B are equal pistons, and draw a tube of uniform bore and of any form connecting the two, and imagine all the fluid except that contained in the tube to be solidified. This will not affect the equilibrium, inasmuch as the fluid pressure on the surface of the tube is at all points perpendicular to the surface whether the fluid be or be not solidified, and the additional pressures on A and B are equal as before.

Also, one piston (A) remaining fixed, the other (B) may be placed with its plane in any direction, and it follows that the pressure upon it is the same for all positions of its plane, or, in other words, the pressure of the fluid is the same in every direction. This proposition we shall enunciate in a general manner in the next article.

The experimental fact that the pressures on pistons of different areas are proportional to those areas may be deduced as follows.

Suppose in a closed vessel two apertures be made in which pistons are fitted, one being a square A , and the other a plane area B , formed by placing together two, three, or any number of squares equal to A ; then the additional pressure on each square being equal to the additional pressure on A , the whole additional pressure will be to the additional pressure on A as the area of B is to that of A^* .

7. *The pressure at any point of a fluid is the same in every direction.*

It is intended by this statement to assert that if at any point of a fluid a small plane area be placed containing the point, the pressure of the fluid upon the plane at that point will be independent of the position of the plane.

The second figure of Art. (5) will serve to illustrate the meaning of the proposition. The aperture in which one of the pistons is fitted may be so constructed as to allow of its plane being changed; and it will be found that in any position, the pressure, or additional pressure, upon the piston is the same.

8. *If a mass of fluid be at rest, any portion may be supposed to become rigid without affecting its equilibrium or the pressure of the surrounding fluid.*

For any portion of a fluid mass may be contemplated as a separate body surrounded by fluid, which presses upon its surface perpendicularly at all points, and its solidification will introduce no change in the pressures upon it, and therefore no change in the pressure at any other point of the fluid.

This proposition enables us to apply the laws of statics to cases of the equilibrium of fluids.

9. The two principles of the equal transmission of pressure and of the equality of pressure in all directions, for the truth of


* If A and B be two pistons of any shape and size, they can be divided into small areas of the same shape and size, and by making these areas small enough, it will be seen that their numbers will be ultimately in the ratio of the areas A and B .

10 Pressure the same in all directions.

which we have appealed to experience, can be deduced from the fundamental property of a fluid, stated as an axiom in Art. (2).

10. The equality of pressure at any point in all directions.

We shall prove this for the case of fluids at rest under the action of gravity, that is, for heavy fluids at rest.



Suppose a small rectangular wedge or prism of fluid, having its sides horizontal and vertical, and its plane ends vertical, to be solidified, and let ABC be its section by a vertical plane bisecting its length. This prism is kept at rest under the action of gravity and of the pressures of the fluid on its ends and sides. The ends are supposed to be perpendicular to the sides of the prism; hence, the pressures on these ends being perpendicular to all the other forces must balance each other, and the pressures on the sides AC , CB , BA , must balance the weight.

Taking d for the length of the wedge, a , b , c the sides of the triangle, w for the weight of an unit of volume of the fluid, and p , p' , p'' for the measures of the pressures on the sides AC , CB , BA , these pressures are

$$pbd, \quad p'ad, \quad \text{and} \quad p''dc,$$

and the weight is $\frac{1}{2}abdw$.

Hence resolving vertically and horizontally,

$$\frac{1}{2}abw = p'a - p''c \cos B,$$

$$pb = p''c \sin B;$$


$$\text{but } a = c \cos B, \text{ and } b = c \sin B;$$

$$\therefore p = p'', \text{ and } p' - p'' = \frac{1}{2}bw.$$

If now we suppose the sides a , b indefinitely diminished, in which case p , p' and p'' will be the pressures in different directions at the point C , we shall have $p' = p''$, and therefore the three pressures are equal*.

By turning the wedge round AC and changing the angle A and B it will be seen that the proposition is true for all directions.

* In strictness $p'p''$ are the measures of the mean pressures on the sides of the wedge, but a reference to Art. 4 on the measure of variable pressure will shew why it is unnecessary to repeat an explanation already made.

11. *The transmission of pressure.*

Let A and B be two points in a fluid at rest, and about the straight line AB as axis describe a cylinder having plane ends perpendicular to AB , and imagine this cylinder solidified.

The equilibrium of the cylinder is maintained by the fluid pressures on its ends, which are parallel to its axis, by the fluid pressures on its curved surface, which are perpendicular to its axis, and by its weight.

Now resolving along AB , the difference of the pressures at A and B must be equal to the resolved part of the weight in the direction BA , and the weight remaining the same, any change of pressure at A involves the same change at B . Moreover, if fluid be contained in a vessel of any shape, and the straight line AB do not lie entirely in the fluid, the two points may be connected by a series of straight lines such as $ACDB$, and any change of pressure at A produces an equal change at C , and therefore, taking account of the previous article, the same change is produced at D , and therefore at B .



12. *The Hydrostatic Bellows* is a machine illustrating the principle of the transmission of fluid pressure.

B is the top of a cylinder having its sides made of leather, and CA is a pipe leading into it. If this vessel and the pipe be filled with water and a pressure applied at A , a very great weight upon B may be raised by a small pressure at A , the weight lifted being greater in proportion to the size of B .



Even without water weights may be raised by simply blowing into the tube *A*.


13. *The Hydrostatic Paradox.*

Any quantity of fluid, however small, may be made to support any weight, however large.

This is another mode of enunciating the same principle. For in the previous figure we may suppose the tube *CA* extended vertically, and the pressure produced by pouring in water to a considerable height, so as to produce a pressure at *A* by means of the column of liquid above it. The tube may be very thin, so that the pressure upon the section *A* of the tube may be very small, but, as this pressure is transmitted to every portion of the surface *B*, which is equal to the section *A*, the force produced can be as large as we please. To increase the upward force on *B* we must enlarge the surface *B*, or increase the height of the column of liquid in the tube, and the only limitation to the increase of the force will be the want of sufficient strength in the pipe and cylinder to resist the increased pressure. By making the height *BC* very small, and the tube *A* of very small bore, the quantity of fluid can be made as small as we please, and hence the paradoxical statement made above.

Hydraulic Presses.

14. The transmission of fluid pressure is the principle upon which Hydraulic or Hydrostatic Presses are constructed.




Thus, if *A*, *B* be two pistons working in hollow cylinders connected by a pipe *C*, and filled with water, any force applied to the piston *B* is transmitted to *A*, and the force upon *A* is greater than the force on *B* in the ratio of the area of *A* to *B*.

This is a Hydraulic Press in its simplest form. Practically it is requisite to have a reservoir from which more water can be obtained by a pump, and we therefore defer the description of a complete Hydrostatic Press until the principle of the Pump has been explained.

The Safety-Valve.

15. In many machines, and especially in steam engines, a very great fluid pressure may be produced, and the strength of the machine may be very severely tried: in order to guard against accidents arising from the bursting of the machine a *safety-valve* is employed, which serves to indicate the existence of too large a pressure.

Various forms may be used, but the principle of the safety-valve is simply that of the uniform transmission of pressure in a fluid.



Thus if BC be one of the connecting tubes through which the fluid passes, and D a small tube opening out of BC , the pressure on a lid at the end of D will measure the fluid pressure within, and if the lid be of a suitable weight, it will be lifted when the pressure is greater than the machine is intended to bear. Suppose, for instance, the greatest permissible pressure of the fluid to be 500 lbs. on a square inch, and the sectional area of the tube D to be $\frac{1}{16}$ th of a square inch, then a weight of $\frac{500}{16}$ or $31\frac{1}{4}$ lbs. will be lifted when the pressure exceeds 500 lbs. The weight employed may be diminished if the lid be moveable about a hinge at A , and a weight w be placed at some little distance from A .

Ex. The tube D is circular, its diameter is one-fourth of an inch, and a weight of 4 lbs. is attached to the lid at a distance of two inches from the hinge; it is required to determine the greatest fluid pressure which will not lift the lid.

The resultant fluid pressure will evidently act at the centre of the circle, and therefore at a distance of $\frac{1}{8}$ th of an inch from A : hence if p be the greatest pressure

required, the forces $p \frac{\pi}{64}$ lbs. and 4 lbs. will balance about the point A , and therefore

$$p \frac{\pi}{64} \cdot \frac{1}{8} = 4 \times 2,$$

$$\text{or } p = \frac{64 \times 64}{\pi}.$$

Taking $\pi = 3$, we obtain roughly $p = 1365$ lbs.

Ex. 2. If the diameter be $\frac{1}{3}$ of an inch, and the distance AW $2\frac{1}{2}$ inches, find the weight which will indicate a pressure of 1000 lbs. on the square inch.

Answer. $5\frac{1}{2}$ lbs. approximately.

16. It will be seen that in Hydrostatic presses, as in all machines, the principle holds that *what is gained in power is lost in motion*.

Thus, if there be two apertures in a closed vessel, fig. art. 5, and the piston B be forced down through any given space, the piston A is forced upwards, if the fluid be incompressible, through a space which is less as the area of A is greater.

This is a simple case of the principle of virtual velocities which we proceed to demonstrate, as applied to incompressible fluids.

Let A, B, C, \dots be the areas of a number of pistons working in cylindrical pipes fitted into the sides of a closed vessel which is filled with fluid. Let the pistons be moved in any manner so that the fluid remains in contact with them, and a, b, c, \dots be the spaces through which they are moved, these quantities being positive or negative, as the pistons are pushed inwards or forced outwards.

Then, since the volume of fluid is the same as before, it follows that

$$Aa + Bb + Cc + \dots = 0,$$

the positive portions, that is, the volumes forced in, being balanced by the negative portions, or the volumes forced out.

But if P, Q, R, \dots be the forces on each piston,

$$P : Q : R : \dots = A : B : C : \dots$$

$$\therefore Pa + Qb + Rc + \dots = 0;$$

or the sum of the products of each force into the space through which its point of application is moved is equal to zero; and observing that a, b, c, \dots are proportional to the virtual velocities of the pistons, this is the equation of virtual velocities, or virtual work.

17. It is not to be imagined that there exists any substance in nature exactly fulfilling the definition which has been given of a fluid. Just as the ideas of a perfectly smooth surface and a perfectly rigid body are formed from observations of bodies of different degrees of rigidity, and surfaces of different degrees of smoothness, so the idea of perfect fluidity is suggested. Nevertheless in the cases of fluids at rest the theoretical properties of fluids derived from this definition will be found to agree with facts, and it is in cases of fluid motion that sensible discrepancies will be found. Thus, a cup of tea set rotating will gradually come to rest, proving the existence of a friction between the liquid and the tea-cup, and also between the particles of the liquid, since the dragging force is gradually communicated from the outer to the inner portions. The motion of water in inclined tubes also indicates the existence of a frictional action amongst the particles of water.

18. Recognizing the fact that all fluids possess, more or less, the characteristic of viscosity, we can give a definition which will include fluids of all degrees of viscosity.

A fluid is an aggregation of particles which yield to the slightest effort made to separate them from each other, if it be continued long enough.

It follows from this definition that in a fluid in equilibrium there can be no tangential action, or shearing stress, and therefore that the pressure on any surface in contact with the fluid is normal to that surface.

Hence all theorems relating to the equilibrium of fluids are true for fluids of any degree of viscosity.

EXAMINATION UPON CHAPTER I.

1. DISTINGUISH between elastic and inelastic fluids. Are any liquids absolutely inelastic?

2. State the property which is assumed as the basis of all reasonings upon fluid action.

3. Define the measure of fluid pressure.

4. It is found that the pressure is uniform over the whole of a square yard of a plane area in contact with fluid, and that the pressure on the area is 13608 lbs.; find the measure of the pressure at any point, 1st, when the unit of length is an inch, 2nd, when it is two inches.

5. The plane of a rectangle, in contact with fluid, is vertical, two of its sides are horizontal, and it is known that at all points of the same horizontal line the pressure is the same. The pressure on the rectangle, for all values of h , is $wbh(a+h)$ where b is the width and h the height of the rectangle; find the pressure at any point of the upper side. (Art. 4.)

6. A cylindrical pipe which is filled with water opens into another pipe the diameter of which is three times its own diameter: if a force of 20 lbs. be applied to the water in the smaller pipe, find the force on the open end of the larger pipe, which is necessary to keep the water at rest.

7. Account for the fact of the transmission of pressure through a liquid.

Mention any direct practical application of this principle.

8. In a Hydrostatic Bellows (Art. 12), the tube A is $\frac{1}{8}$ th of an inch in diameter, and the area B is a circle, the diameter of which is a yard. Find the weight which can be supported by a pressure of 1 lb. on the water in A .

9. A safety-valve consists of a heavy rectangular lid which is horizontal when it closes the aperture beneath it, and is moveable about one side. The aperture being a square which has one side coincident with the fixed side of the lid, find the maximum pressure marked by the valve.

10. Prove the principle of virtual velocities in the case of the sixth question.

11. A triangular area ABC is exposed to fluid pressure, and it is found that if any straight line PQ be drawn parallel to BC , and at a distance x from A , the pressure on the area APQ is px^2 ; find the pressure at A , and also at any point of the line BC .

12. A strong cylindrical tube, one foot in diameter inside, and ten feet in length, is filled with distilled water, and closed with a piston to which a pressure of 10000 lbs. is applied; shew that the resulting compression of the water will be nearly $\frac{1}{17}$ th of an inch.

CHAPTER II.

Density and Specific Gravity.

19. **I**N the classification of fluids the most prominent division is between gases and liquids, or elastic and non-elastic fluids, as they are sometimes termed, and under these two heads all fluids are naturally ranged. It has been remarked already that the term non-elastic is inaccurate, but no confusion will be produced by its use, as the compressibility of liquids is practically insensible, and for all ordinary purposes unimportant.

It will be found, however, that the theory of sound is partly dependent on this compressibility, and it is therefore of importance at once to recognize its existence.

There are many other characteristics which distinguish fluids from each other, such as colour, degree of transparency, chemical qualities, viscosity, &c., but in the theory of Hydrostatics and Hydrodynamics the only characteristic which it is necessary to consider is the *density* or *specific gravity* of a fluid.

It is not meant that density and specific gravity are synonymous terms, but that these terms have reference to the substance of a fluid.

Thus, a cubic inch of mercury and a cubic inch of water have different weights, the former being more than 13 times the latter, and it is inferred that the quantity of matter in the mercury is greater than in the water, or that the density of mercury is greater than that of water.

These remarks apply to both fluid and solid bodies, and the density and specific gravity of a fluid or solid must be measured respectively by reference to the density and specific gravity of some standard substance.

20. We may remark at this point that in one respect all fluids agree, whether elastic or not; they are all *ponderable* bodies, that is, they are all acted upon by gravity, and exhibit different degrees of intrinsic weight. The term density refers to the material of which bodies are composed, and the idea of a difference of density in two bodies does not involve the conception of weight, while the term specific gravity refers to the varying effect of the action of gravity on different bodies.

21. DEFINITION. *The measure of the density of a body is the number expressing the ratio which the mass of any volume of the body bears to the mass of an equal volume of the standard substance.*

For any given fluid let ρ be this number, and let unity represent the standard substance; ρ is then the density of the fluid measured in terms of the density of the standard substance.

It is clear that if a body be compressed into half its original volume, its density will be doubled, while its mass or the quantity of matter contained in it remains the same; and similarly, if it be compressed in any other proportion, its density will be increased in the same proportion. This is expressed by saying that

$$M \propto \rho V,$$

M representing the mass and V the volume.

Again, it is known that the weight of a body depends upon its position on the earth's surface, but that in all cases if g be the local acceleration due to the action of gravity, the weight of a given body, that is, a body of given mass, varies as g ,

$$\text{or } W \propto g,$$

and it is obvious that at all places

$$W \propto M.$$

Therefore generally $W \propto Mg$,

$$\text{or } W \propto g\rho V;$$

and we may suppose the units involved in these several symbols so chosen that the relations may be

$$M = \rho V, \quad W = Mg, \quad \text{and } \therefore W = g\rho V.$$

Absolute unit = force which gives to unit mass unit acceleration in unit time.

22. It must be observed that this formula defines one unit, if the rest be given.

Thus, if the units of space and time be a foot and a second, it is known that $g=32.2$, and making $\rho=1$, and $V=1$, i.e. a cubic foot, we obtain the weight of an unit of volume of the standard substance

$$=(32.2) \text{ unit of weight;}$$

and therefore the unit of weight is $\frac{1}{32.2}$ (the weight of a cubic foot of the standard substance).

Now, if distilled water at a temperature 60° F. be the standard substance, the weight of a cubic foot is about 1000 oz.; and therefore the formula $W=g\rho V$ implies that the unit of weight is $\frac{1000}{32.2}$ oz.;

$$\text{and therefore that } W=1000\rho V \text{ oz.}$$

Ex. Taking distilled water as the standard substance, find the weight of 12 cubic feet of a substance of which the density is 3.5.

$$\text{The weight} = g \times 3.5 \times 12 \times \frac{1000}{g} \text{ oz.} = 42000 \text{ oz.}$$

This example is more directly solved by observing, that the weight must be 3.5 times that of 12 cubic feet of water.

23. In the previous articles we have considered homogeneous bodies only; if the density be variable, or the bodies be heterogeneous, the density *at any point* of a body is that of a homogeneous mass which has the same density as the body about the proposed point.

If the density vary continuously from point to point we may determine it *at a point* by taking a small mass of the fluid containing the point, and comparing its weight with that of an equal volume of the standard substance, it being conceived that in a very small continuous mass the density will not vary sensibly throughout.

24. In order to render more clear the mathematical conception of a continuously varying substance, imagine a number of homogeneous strata of equal thickness t placed on each other, and suppose the density of the lowest stratum to be ρ and of the highest ρ' , and of the intermediate strata let the densities increase by successive additions from ρ' to ρ

Gravitation unit - force which will give to unit mass, the same acceleration as gravity in unit time.

If now we suppose the thickness of each stratum t to become indefinitely small, and the number of intermediate strata to become indefinitely large, while the densities of the extreme strata ρ, ρ' remain the same, the densities of the intermediate strata which are to increase from ρ' to ρ will differ from each other by infinitely small quantities, and we can thus form an idea of a continuously varying medium.

This mode of viewing continuity by means of discontinuity is necessary for the purposes of mathematical calculation.

The atmosphere in a state of rest is a case in point, as its density decreases continually as the height increases.

25. The density of a mixture may be determined by the previous formula $M = \rho V$.

Thus, if volumes V, V', V'', \dots of fluids whose densities are $\rho, \rho', \rho'', \dots$ be mixed together, and if the mixture form a homogeneous mass, and no change of volume occur from chemical action, the whole mass

$$= \rho V + \rho' V' + \rho'' V'' + \dots = \Sigma(\rho V),$$

$$\text{and the whole volume} = V + V' + V'' + \dots = \Sigma(V);$$

$$\therefore \text{the density of the mixture} = \frac{\Sigma(\rho V)}{\Sigma V}.$$

26. DEFINITION. *The measure of the specific gravity of a body is the number expressing the ratio which the weight of any volume of the body bears to the weight of an equal volume of the standard substance.*

This definition, it will be seen, makes the measure of specific gravity the same as that of its density, provided the standard substance be the same in both cases. The standard substance, however, is not necessarily the same.

If s be the specific gravity of a body or fluid, and W the weight of a volume V of the body or fluid, we have the relation

$$W = sV,$$

which means that if the unit of weight be the weight of an unit of volume of the standard substance, the weight is sV times that unit of weight.

Thus, if distilled water be the standard, and one foot be the unit of length, the weight of a volume V of a fluid, of which the specific gravity is s , $=sV$ times the weight of a cubic foot of water,

$$= 1000sV \text{ oz.}, \text{ or } \frac{1000}{16} sV \text{ lbs.}$$

27. *To find the specific gravity of a mixture of given volumes of any number of fluids, whose specific gravities are given.*

Let V, V', V'', \dots be the volumes of fluids of which the specific gravities are s, s', s'', \dots

Then the weight of the mixture is

$$sV + s'V' + s''V'' + \dots \text{ or } \Sigma(sV),$$

and the volume is $V + V' + V'' + \dots$ or $\Sigma(V)$,

and therefore if σ be the specific gravity of the mixture,

$$\sigma \Sigma(V) = \Sigma(sV),$$

$$\text{or } \sigma = \frac{\Sigma(sV)}{\Sigma(V)}.$$

If by any chemical action among the fluids the volume becomes U instead of $\Sigma(V)$, the specific gravity will be

$$\frac{\Sigma(sV)}{U}.$$

28. *To find the specific gravity of a mixture when the weights and specific gravities of the components are given.*

Let W, W', W'', \dots be the weights, and s, s', \dots the specific gravities of the respective fluids.

Their volumes are respectively $\frac{W}{s}, \frac{W'}{s'}, \dots$

$$\text{and the whole volume} = \frac{W}{s} + \frac{W'}{s'} + \dots = \Sigma\left(\frac{W}{s}\right),$$

$$\text{while the whole weight} = W + W' + \dots = \Sigma(W).$$

Hence if σ be the specific gravity of the mixture,

$$\sigma \cdot \Sigma\left(\frac{W}{s}\right) = \Sigma(W).$$

29. *To find the unit of specific gravity, or to determine the specific gravity of the standard substance when the units of length and weight are given.*

From the equation $W=sV$, which gives $W=1$ when $s=1$ and $V=1$, it appears that the standard substance is one of which the weight of an unit of volume is the unit of weight.

Thus if 1 lb. and 1 foot be the units, the standard is that substance of which a cubic foot weighs one pound.

Now a cubic foot of water weighs 1000 oz. ; therefore $\frac{1}{1000}$ of a cubic foot of water weighs 1 lb.

The standard substance therefore is such that a cubic foot of it weighs the same as $\frac{1}{1000}$ of a cubic foot of water.

Hence the density of the standard is to the density of water as 16 : 1000.

30. *Comparison of the equations $W=sV$, $W=g\rho V$.*

It appears from the definitions that, if the standard substances be the same, the measures of the density and specific gravity of any given fluid are the same, that is, the numbers s and ρ will be identical. The standards however not being necessarily the same, s and ρ will be in general different numbers.

From the equations $W=sV$, $W=g\rho V$, we infer that if the standard substances are the same and the units of length the same, the units of weight are different. In fact, the unit of weight in the first equation would be g times that in the second.

We also infer that if the units of weight and length are the same, the standard substances are different. Thus if s and ρ refer to a substance of which a volume V weighs W , then $s=g\rho$, and therefore the density of the standard to which s refers is *less* than that to which ρ refers in the ratio of $g : 1$.

In the equation $W=g\rho V$, the unit of time enters, the value of g depending upon it ; and, by a change in the unit of time, one or more of the other units, those namely of length, weight, and density, is necessarily changed.

31. The practical methods of determining the specific gravities of solids, liquids, and gases will be discussed in a future chapter.

For solids and liquids tables of specific gravity are usually given with reference to distilled water at 60° F. as the standard.

Gases and vapours are, however, generally referred to atmospheric air at the same temperature and under the same pressure as the gases themselves.

EXAMINATION UPON CHAPTER II.

1. EXPLAIN how density is measured.

What convention with regard to units is implied in the equation $W = g\rho V$?

2. Find the weight of a cubic foot of mercury, the specific gravity of which is 13.568.

3. If a cubic inch of a standard substance weigh .45 of a lb., what is the weight of a cubic yard of a substance of which the density is 5?

4. A mixture is formed of two fluids; the specific gravity of the mixture, the ratio, $m : 1$, of the volumes, and the ratio, $n : 1$, of the specific gravities are given; find the specific gravities of the fluids.

5. Equal weights of two fluids, of which the densities are ρ and 2ρ , are mixed together, and one-third of the whole volume is lost; find the density of the resulting fluid.

6. Taking water as the standard, find the weight of a cubic yard of a substance of which the specific gravity is .12.

7. A cubic inch of a substance weighs $\frac{1}{3} \frac{6}{11} \frac{2}{5}$ ths of a lb.; find its specific gravity referred to water.

8. A mixture is formed of equal volumes of three fluids; the densities of two are given and the density of the mixture is given; find the density of the third fluid.

9. Volumes V , V' of two fluids, the specific gravities of which are σ , σ' , are mixed together, and the specific gravity of the mixture is s ; find the change in volume.

10. Two fluids of equal volume, and of specific gravities s , $2s$, lose $\frac{1}{4}$ th of their whole volume when mixed together; find the specific gravity of the mixture.

EXAMPLES.

1. A mixture is formed of equal volumes of n fluids, the densities of which are in the ratio of the numbers 1, 2, 3, ... n ; find the density of the mixture. Also find the density of the mixture when the volumes are:—1st, in the ratio of the numbers 1, 2, 3, ... n , and 2nd, of the numbers n , $n-1$, ... 3, 2, 1.

2. Having given the specific gravity σ of a mixture formed of equal volumes of two fluids, and also the specific gravity σ' of a mixture formed by taking a quantity of one fluid double that of the other; find the specific gravities of the fluids.

3. If 25 oz. be the unit of weight, and a yard and a second the units of length and time, what is the density of the standard substance compared with that of water in the equation $W = g\rho V$?

4. If 3 cwt. be the unit of weight, and 4 feet the unit of length, find the density of the standard compared with that of water, the unit of time being one second.

5. If the units of weight, length, and time be 1 lb., one foot, and one second, compare the standards in the formulæ $W = sV$, $W = g\rho V$.

6. If the units of weight and length be the same, the latter being one foot, find the unit of time in order that the standard may be the same in both formulæ.

7. If the units of weight, length, and time be 1 lb., one yard, and half a second, compare the standards in the equations $W = g\rho V$, $W = sV$.

8. If the standards be the same, and also the units of weight, find the unit of length in $W = g\rho V$, the unit of time being $2\sqrt{2}$ seconds, and the unit of length in $W = sV$ being one foot.

9. If the units of length and time be 3 yards and 4 seconds, and if the units of weight be also the same in both equations, compare the densities of the standard substances.

10. When a vessel is filled by means of equal volumes of two fluids, the specific gravity of the compound is $\frac{4}{3}$ of what it would have been if the vessel had been filled by means of equal weights of the fluids. Compare the specific gravities of the two fluids.

NOTE. In the above examples assume that $g = 32$ when a foot and a second are units.

Slip

CHAPTER III.

Pressure at different points of a liquid at rest, Surface of a liquid, Liquids maintaining their level, Liquids in a bent tube, Pressures on Plane Surfaces, Whole Pressure, Centre of Pressure.

32. **T**HE pressure of a liquid at rest is the same at all points of the same horizontal plane.

Take a thin cylindrical portion AB of the liquid, having its axis horizontal, and its ends A, B vertical, and



imagine this portion to become solid. We have then a solid body AB kept at rest by the fluid pressures on its curved surface, all of which are perpendicular to the axis of the cylinder, by the pressures on the two ends, which are horizontal, and by the weight of the solid.

If p and p' be the measures of the pressures at A and B , and a the area of each end, which is taken to be very small in order that the pressure may be sensibly uniform over the whole of either end, the pressures on the ends are pa and $p'a$, and since these balance each other, we have


$$p = p'.$$

This proof also holds good for the case of elastic fluids, or of heterogeneous incompressible fluids.

33. *To find the pressure at any given depth in a heavy homogeneous liquid at rest.*

Taking any point P in the fluid, draw PA vertically to the surface, and describing a thin cylinder about PA with its base horizontal, imagine it to become solid.

Then the solid body PA is kept at rest by the fluid pressure on the end P , the weight of the solid, and the fluid pressures on the curved surface, which are all horizontal.



Hence the fluid pressure on P must be equal to the weight, and therefore, if a be the area of the base, w the weight of an unit of volume, and p the pressure at P ,

$$pa = wa \cdot AP,$$

$$\text{or } p = w \cdot AP;$$

that is, the pressure at any depth varies as the depth below the surface.


Similarly, if P and Q be any two points in the same vertical line, by solidifying a cylinder PQ , it will be seen that the difference of the pressures on the ends P and Q of the cylinder must be equal to the weight of the cylinder.

Hence if p, p' be the pressures at P and Q ,

$$p'a - pa = wa \cdot PQ,$$

$$\text{or } p' - p = w \cdot PQ;$$

that is, *the difference of the pressures at any two points varies as the vertical distance between the points.*



If ρ be the density of the liquid, the weight of AP is $g\rho AP$, and therefore, if $AP = z$,

$$p = g\rho z.$$

34. The form, $g\rho z$, of the expression for p is the one which we shall generally employ, and a few remarks upon it will be useful.

The symbol p represents the pressure on an unit of area, and therefore its numerical value depends on the unit of length which

is adopted. Further, the numerical value of g depends on the unit of time as well as the unit of length, and the value of ρ depends on the standard to which the liquid is referred. Hence the numerical value of p depends on all these elements.

Thus, if water be the standard, and a foot and a second be units of space and time, we obtain, at a depth of one foot in water, putting $\rho=1$ and $z=1$, $p=32$. Now we know that at this depth the pressure is really 1000 oz. per square foot, and therefore p being 32, the unit of weight must be $\frac{1000}{32}$ oz.

Hence with these units the actual pressure at a depth z is $1000\rho z$ oz.

Again, if 1 lb. be the unit of weight, one second the unit of time, and water the standard, we shall obtain

$$x^3 \frac{1000}{16} = g = \frac{32}{x},$$

x feet being the unit of length, and

$$\therefore x^4 = \frac{32 \times 16}{1000} \text{ of a foot.}$$

In all these cases it will be seen that we must have given some fact relating to the weight or density of the standard substance.

35. Let the cylinder of which AP is the axis be bounded at P by a plane not horizontal, and let a' be its area and θ its inclination to the horizon.

Then for the equilibrium of the cylinder, taking p' as the pressure at P upon a' , we have by resolving vertically,


$$p' a' \cos \theta = waAP,$$

$$\text{but } a = a' \cos \theta;$$

$$\therefore p' = w \cdot AP, \text{ which is independent of } \theta.$$

We thus have another proof of the proposition that the pressure at any point is the same in all directions.

It may be perhaps objected to the proof of Art. 33 that the surface at A is assumed to be horizontal. By making the cylinder AP a very thin cylinder, that is, of very small radius, it will be seen that its weight is sensibly $g\rho aAP$, and therefore that the proof does not depend on any assumption as to the form of the surface.



Or, to reason more strictly, draw two horizontal planes through the highest and lowest points B , A of the small portion AB of the surface intercepted by the cylinder.


Then, if the radius of the cylinder be indefinitely diminished, these two planes will coalesce.

If z and z' be the heights above P of these planes, the weight of the cylinder lies between

$$gpaz \text{ and } gpaz',$$

and therefore p lies between gpz and gpz' , and ultimately when the planes coalesce,

$$p = gpz.$$




36. *Difference of pressures at any two levels in an elastic fluid.*

We have already mentioned in Art. 20, that gases are heavy bodies; hence, by the same reasoning as in Art. (33), if P and Q be two units of area in an elastic fluid, P being vertically above Q , the difference of the pressures at P and Q is equal to the weight of the column of air PQ . This column is not of uniform density, and hence the law of variation of the pressure at different levels in an elastic fluid does not present itself in a simple form. Further information will be found in Chapter V; at this point we need only call attention to the fact that the pressure decreases as we ascend in an elastic fluid.

37. *The surface of a liquid at rest is a horizontal plane.*

Take two points P , Q , in the same horizontal plane, within the liquid, and draw PA , QB vertically to the surface.

Then pressure at $P = w \cdot AP$,
 pressure at $Q = w \cdot BQ$,
 and these are equal; therefore AP and BQ are equal, and A , B are in the same horizontal



plane. Similarly any other point in the surface can be proved to be in the same horizontal plane with A or B .

Or we might have argued that, since the pressures are equal at all points of the same horizontal plane, conversely, all points at which the pressures are equal are in the same horizontal plane, and therefore all points in the surface, at which the pressure is either zero, or equal to the atmospheric pressure, must be in the same horizontal plane.

38. The pressure of the atmosphere is found to be about 14.73 lbs. to a square inch, or very nearly 15 lbs. We can hence calculate the pressure upon any given area, and, if Π be the atmospheric pressure on the unit of area, the pressure at a depth z of a fluid, the surface of which is exposed to atmospheric pressure, will be


$$g\rho z + \Pi.$$

39. *Illustration.* Take a hollow glass cylinder open at both ends; in contact with the lower end, and closing that end, place a heavy flat disc supported by a string passing up the cylinder.

Holding the string, depress the cylinder in a vessel of water, and it will be found that, at a certain depth, the string may be loosened, and the disc will remain in contact with the cylinder, being supported by the pressure of the water beneath.

If w be the weight of the disc and r the radius of the cylinder, the requisite depth (x) of the disc is given by the equation

$$w = g\rho x\pi r^2.$$




The presence or absence of the atmosphere will not affect this depth, since the pressure of the atmosphere downwards on the disc would be counteracted by the pressure upwards, transmitted from the surface of the water.

40. If in Art. (32) the line AB do not lie entirely

within the fluid, we can still prove the truth of the proposition by the aid of Art. (33).


For A and B can be connected by horizontal and vertical lines as AC , CD , DB , and



$$\begin{aligned} \text{pressure at } B &= \text{pressure at } D - w \cdot BD \\ &= \text{pressure at } C - w \cdot AC \\ &= \text{pressure at } A. \end{aligned}$$

41. Hence it appears that all points on the surface of a liquid, at which the pressure is either zero or is equal to the constant atmospheric pressure, must be in the same horizontal plane, and that this is true even though the continuity of the surface be interrupted by the immersion of solid bodies, or in any other way.

This sometimes appears under the form of the assertion that *liquids maintain their level*, and an experimental illustration may be employed as in the figure.




A number of glass vessels of different forms, all open into a closed tube or vessel AB , and it is found that if water be poured into any one of the tubes, it will, after filling the tube AB , rise to exactly the same vertical height in every one of the tubes, and if any portion be withdrawn from any of the vessels, that the water will sink to its new position of rest through the same vertical height in each.

An important practical illustration of this principle is seen in the construction by which towns are supplied with water. A reservoir is placed on a height, and pipes leading from it carry the water to the tops of houses or to any point which is not higher than the surface of the water in the reservoir, and these pipes may be carried under ground or over a road, provided that no portion of a pipe is above the original level.

42. *The common surface of two liquids that do not mix is a horizontal plane.*

Take two points P , Q in the lower fluid, both in the same horizontal plane, and let vertical lines PA , QB to the surface of the upper fluid meet the common surface of the fluids in C and D .



Then if w' be the weight of an unit of volume of the lower fluid, and w of the upper,

pressure at $P = w' \cdot CP +$ pressure at C

$$= w' \cdot CP + w \cdot CA,$$

$$\text{and at } Q = w' \cdot QD + w \cdot DB;$$

$$\therefore w' \cdot CP + w \cdot CA = w' \cdot QD + w \cdot DB.$$

Also AB is horizontal, and therefore

$$CP + CA = QD + DB;$$

\therefore multiplying by w and subtracting,

$$(w' - w) CP = (w' - w) QD,$$

or $CP = QD$, and therefore CD is horizontal.

43. *If two liquids that do not mix together meet in a bent tube, the heights of their upper surfaces above their common surface will be inversely proportional to their densities.*


Let A and B be the two surfaces, C the common sur-

face, and ρ, ρ' the densities of BC and CA .

Let horizontal planes through A, B , and C meet a vertical line in a, b , and c , and take C' in the denser fluid in the same horizontal plane as C .

The pressure at $C = g\rho.bc$, and at $C' = g\rho'.ac$, and these are equal, by Art. 32;


$$\begin{aligned} \therefore \rho bc &= \rho' ac, \\ \text{or } bc : ac &= \frac{1}{\rho} : \frac{1}{\rho'}. \end{aligned}$$



44. *Two fluids that do not mix are contained in the same vessel; it is required to find the pressure at a given depth in the lower fluid.*

Let P be the point in the lower fluid, PBA a vertical line meeting the common surface in B . Describe a small cylinder about AP , and suppose it solidified.

Then, if p be the pressure at P and a the sectional area of the cylinder,



$$pa = \text{weight of } ABP = g\rho ABa + g\rho'BP a.$$

ρ and ρ' being the densities,

$$\text{or } p = g\rho AB + g\rho'BP.$$

This might have been at once inferred from the equation

$$p = g\rho'BP + \text{pressure at } B,$$

for the pressure at $B = g\rho AB$.

And in the same manner the pressure at any point of a mass of fluid containing any number of strata of different densities can be determined.

If the surface A be subject to the atmospheric pressure Π ,

$$\text{the pressure at } P = g\rho'BP + g\rho AB + \Pi.$$

45. We now proceed to consider two simple cases of the pressure of a fluid on plane surfaces.

PROP. *The pressure of a liquid on any horizontal area is equal to the weight of a column of the liquid of which the area is the base and of which the height is equal to the depth of the area below the surface.*


For, if z be the depth, the pressure at every point is wz or gpz ;

\therefore if κ be the area, the pressure upon it $= wz\kappa$,
and $z\kappa$ is the volume of the column described.

It will be seen that this is independent of the form of the vessel containing the fluid.

This result may also be obtained in the following manner.

Draw through the boundary of K vertical lines to the surface, and suppose the portion of fluid enclosed to be-




come solid. The pressure of the surrounding fluid is entirely horizontal, and therefore the pressure on the base must be equal to the weight of the solid.

If the vessel be of the form indicated in the dotted line so that the actual surface does not extend over the area K , we may suppose the fluid extended over K by enlarging the vessel, and the pressure at any point of K will not be changed. Hence the above reasoning is applicable to this case also.

Thus if a hollow cone, vertex upwards, be filled with water, and if r be the radius of the base and h the height of the cone, the pressure on the base $= w\pi r^2 h$, or $gp\pi r^2 h$, that is, the weight of the cylinder of fluid on the same base as the cone, and of the same height.

46. A plane area in the form of a rectangle is just immersed in liquid with one edge in the surface, and its plane inclined at an angle θ to the vertical; it is required to find the pressure upon it.

Let the figure be a vertical section perpendicular to



the side b of the rectangle in the surface, $AB (= a)$ being the section of the rectangle.

Draw a vertical plane BC through the lower side B , and suppose the fluid in ABC to become solid; then its weight is supported by the plane AB , since the pressure on BC is horizontal.

Hence if R be the pressure on AB , perpendicular to its plane,

$$\begin{aligned} R \sin \theta &= \text{weight of } ABC = \frac{1}{2} w \cdot AC \cdot BC \cdot b \\ &= \frac{1}{2} w b a^2 \sin \theta \cos \theta; \\ \therefore R &= \frac{1}{2} w b a^2 \cos \theta = w b a \cdot \frac{1}{2} a \cos \theta, \end{aligned}$$

that is, the pressure is the weight of a column of fluid of which the rectangle is the base, and the height is equal to the depth of the middle point of AB below the surface.

Since the direction of R makes an angle θ with the horizon, it follows that the horizontal component of R is

$$\frac{1}{2} w b a^2 \cos^2 \theta.$$

Now the solid ABC is kept at rest by the horizontal pressure on BC , by its weight, and by the reaction R .

$$\begin{aligned} \text{Hence the pressure on } BC &= R \cos \theta = \frac{1}{2} w b a^2 \cos^2 \theta \\ &= w \cdot b a \cos \theta \cdot \frac{1}{2} a \cos \theta \end{aligned}$$

$= w \cdot (\text{area } BC) (\text{depth of middle point of } BC)$,
the same law as for AB .

This also appears from the value of R by putting $\theta = 0$.

The results thus obtained are generalized in the following article in which a different method is adopted.

Whole Pressure.

47. DEF. *The whole pressure of a fluid on any surface is the sum of all the normal pressures exerted by the fluid on every portion of the surface.*

In the case of a plane, the pressure at every point is in the same direction and the whole pressure is the same as the resultant pressure. In the case of curved surfaces, the whole pressure is merely the arithmetical sum of all the pressures acting in various directions over the surface.

48. PROP. *The whole pressure of a liquid on a surface is equal to the weight of a column of liquid of which the base is equal to the area of the surface, and the height is equal to the depth of its centre of gravity below the surface of the liquid.*

Let the surface be divided into a great number of very small areas a_1, a_2, a_3, \dots and let z_1, z_2, z_3, \dots be the depths below the surface of the centres of gravity of these areas. By making the areas very small, each may be considered plane, and the pressures upon them will be respectively

$$w a_1 z_1, w a_2 z_2, \dots$$

taking the pressure over each area to be uniform.

Hence the whole pressure = $w \Sigma (az)$.

But, if \bar{z} be the depth of the centre of gravity of the surface,

$$\bar{z} = \frac{\Sigma (az)}{\Sigma (a)}^* ;$$

\therefore whole pressure = $w \bar{z} \Sigma (a)$

= $w \bar{z} S$, if S be the area of the surface,

and $\bar{z} S$ is the volume of the column described.

If ρ be the density of the liquid, the expression for the whole pressure is $g \rho \bar{z} S$.

EX. 1. A rectangle is immersed with two sides horizontal, the upper one at a given depth (c), and its plane inclined at a given angle (θ) to the vertical.

Let a be the horizontal side, b the other side.

* See Goodwin's *Statics*, or Parkinson's *Mechanics*, Art. 71.

The depth of the centre of gravity $= \frac{1}{2}(2c + b \cos \theta)$, and the whole pressure $= \frac{1}{2}w(2c + b \cos \theta)ab$.

Ex. 2. A vertical cylinder, radius r and height h , is filled with fluid.

The surface $= 2\pi r h$ and the whole pressure $= w\pi r h^2$.

Ex. 3. A hollow cone, vertex downwards, is filled with water.

Let r be the radius, and h the height of the cone.

By cutting the cone down a generating line and unrolling it into a plane, its surface forms the sector of a circle, of which the slant side is the radius and the perimeter of the base is the arc.


But the area of a sector $= \frac{1}{2}(\text{arc})(\text{radius})$;

$$\therefore \text{the surface} = \pi r \sqrt{r^2 + h^2}.$$

Again, the surface of a cone is the ultimate form of the surface of a pyramid formed by triangles, having the vertex of the cone as their common vertex, and having for their bases the sides of a polygon inscribed in the circle, and since the centre of gravity of each triangle is at a depth $\frac{1}{3}h$ below the surface of the fluid, it follows that $\frac{1}{3}h$ is the depth of the centre of gravity of the surface.

Hence the whole pressure $= \frac{1}{3}w\pi r h \sqrt{r^2 + h^2}$.

Ex. 4. The cylinder in Ex. 2, closed at both ends, is just filled with liquid, and its axis is inclined at an angle θ to the vertical.



The surface of the fluid is a horizontal plane through the highest point of the cylinder, and the depth of G

$$= \frac{h}{2} \cos \theta + r \sin \theta.$$

Hence the whole pressure on the curved surface is

$$w\pi rh(h \cos \theta + 2r \sin \theta),$$

and the whole pressure including the plane ends is

$$w(\pi rh + \pi r^2)(h \cos \theta + 2r \sin \theta).$$

Ex. 5. A cubical vessel is filled with two liquids of given densities, the volume of each being the same, it is required to find the pressure on the base and on any side of the vessel.


Let a be a side of the vessel, ρ, ρ' the densities of the upper and lower liquids, ρ' being taken greater than ρ .

The pressure on the base = the weight of the whole fluid

$$= g\rho' \frac{a^3}{2} + g\rho \frac{a^3}{2}.$$

The pressure on the portion BC

$$= g\rho \frac{a^2}{2} \cdot \frac{a}{4} = \frac{1}{8} g\rho a^3.$$



To find the pressure on AC , replace the liquid DC by an equal weight of the lower liquid. This change will not affect the pressure at any point of CA .

If $B'D$ be its surface,

$$\rho' CB' = \rho CB = \rho \frac{a}{2},$$

and the depth of the centre of gravity of AC below B'

$$= B'C + \frac{a}{4} = \frac{a}{4} \left(1 + \frac{2\rho}{\rho'} \right);$$

$$\therefore \text{the pressure on } AC = g\rho' \frac{a^2}{2} \cdot \frac{a}{4} \left(1 + \frac{2\rho}{\rho'} \right)$$

$$= \frac{1}{8} g\rho' a^3 \left(1 + \frac{2\rho}{\rho'} \right).$$


Centre of Pressure.

49. DEF. The centre of pressure of a plane area is the point of action of the resultant fluid pressure upon the plane area.

As a simple case, suppose a rectangle immersed in a liquid with one side in the surface.

Divide the area into a number of very small equal parts by equidistant horizontal lines.

The pressure on each part will act at its middle point and will be proportional to the depth below the surface, and we have to find the centre of a system of parallel forces acting perpendicularly to the plane at equidistant points of the line EF and proportional to the distance from E .




This is evidently the same as finding the centre of gravity of a triangle of which E is the vertex and F the middle point of the base. The centre of pressure therefore divides EF in the ratio 2 : 1.

It will be seen that this result is independent of the inclination of the plane of the rectangle to the vertical.

If a triangular area be immersed with its vertex in the surface and its base horizontal, and be divided by equidistant horizontal lines, the pressure on each strip will act at its middle point and be proportional to the square of the distance of that point from the vertex E .

Hence if F be the middle point of the base, the centre of pressure will be the same as the centre of gravity of a solid cone, vertex E and axis EF , and therefore divides EF in the ratio 3 : 1.

If a triangular area be immersed with its base in the surface, the pressure on a strip will be proportional to the product $EN \cdot NF$, and consequently proportional to the square of the ordinate NP of a semi-circle described upon EF as diameter.



The centre of pressure will therefore be the middle point of *EF*.

We may also give the following general method, applicable to the case of a plane area immersed in any position.

Through the boundary line of the plane area draw vertical lines to the surface and consider the equilibrium of the liquid so enclosed ; the reaction of the plane resolved vertically, is equal to the weight of the liquid, which acts in the vertical line through its centre of gravity ; and the point in which this line meets the plane is the centre of pressure.

50. The student will now be able to appreciate more clearly the nature of fluid pressures, and to see that the action of a fluid does not depend upon its quantity, but upon the position and arrangement of its continuous portions. It must be carefully borne in mind that the surface of an inelastic fluid or liquid is always the horizontal plane drawn through the highest point or points of the fluid, and that the pressure depends only on the depth below that horizontal plane.

Thus in the construction of dock-gates, or canal-locks, it is not the expanse of sea outside which will affect the pressure, but the height of the surface ; and, in considering the strength required in the construction, the greatest height of the surface due to tides must also be taken into account. Any violent action due to rapid tides or storms is of course a subject for separate consideration.

The same principle shews that in the construction of dikes, or the maintenance of river-banks, the strength must be proportional to the depth below the surface.

EXAMINATION UPON CHAPTER III.

1. To what extent is the pressure on the base of a vessel affected by pouring in more liquid?
2. Find the pressure at a depth of 100 feet in a lake, 1st, neglecting, 2nd, taking account of the atmospheric pressure.
3. Explain the statement that liquids maintain their level.
4. A reservoir of water is 200 feet above the level of the ground-floor of a house; find the pressure of the water in a pipe at a height of 30 feet above the ground-floor.
5. Three liquids that do not mix are contained in a vessel; prove that their common surfaces are horizontal, and find the pressure at any depth in the lowest liquid.
6. An equilateral triangular area is immersed in water with a side 1 ft. in length in the surface; find the pressure upon it in lbs.
7. Distinguish between whole pressure and resultant pressure.
8. A hollow cone, vertex upwards, is just filled with liquid; find the whole pressure on its curved surface.
9. Prove that the depth of the centre of pressure of a plane area is greater than the depth of the centre of gravity of the area.
10. Find the centre of pressure of a rectangular area immersed, with plane vertical and two sides horizontal.
11. A rectangle has one side in the surface of a liquid; divide it by a horizontal line into two parts on which the pressures are equal.
12. Divide the same rectangle by horizontal lines into n parts on which the pressures are equal.
13. A triangle has its base horizontal and its vertex in the surface; divide it by a horizontal line into two parts on which the pressures are equal.

EXAMPLES.

1. Two equal vertical cylinders standing on a horizontal table are connected together by a pipe passing close to the table, and are partially filled with water. In contact with and

above the water in one cylinder is a closely-fitting piston of given weight; find its position of equilibrium.

2. The upper surface of a vessel filled with water is a square whose side is 2 feet 6 inches, and a pipe communicating with the interior is filled with water to a height of 8 feet; find the weight (in lbs.) which must be placed on the lid of the vessel to prevent the water from escaping, the weight of a cubic foot of water being 1000 oz.

3. A parallelogram is immersed in a liquid with one side in the surface; shew how to draw a line from one extremity of this side dividing the parallelogram into two parts on which the pressures are equal.

4. A fine tube ABC is bent so that the portions AB , BC are straight and perpendicular to each other; the tube is placed so that each branch is equally inclined to the vertical, and equal quantities of two liquids, the densities of which are in the ratio of 2 : 1, are poured into the respective branches; find the height above B of their common surface.

5. A smooth vertical cylinder one foot in height and one foot in diameter is filled with water, and closed by a heavy piston weighing 4 lbs; find the whole pressure on its curved surface.

6. If a ball, weighing 1 lb. in water, be suspended in the water by a string fastened to the piston, and if the specific gravity of the metal be to that of water as 7 to 2, find the pressure at any depth and the whole pressure on the curved surface.

7. A cylindrical vessel standing on a table contains water, and a piece of lead of given size supported by a string is dipped into the water; how will the pressure on the base be affected, (1) when the vessel is full, (2) when it is not full? and in the second case, what is the amount of the change?

8. A hollow cylinder closed at both ends is just filled with water and held with its axis horizontal: if the whole pressure on its surface, including the plane ends, be three times the weight of the fluid, compare the height and diameter of the cylinder.

9. A triangle ABC is immersed vertically in a liquid with the angle C in the surface and the sides AC , BC equally inclined to the surface; shew that the vertical through C divides the triangle into two others, the fluid pressures upon which are as $b^3 + 3ab^2 : a^3 + 3a^2b$.

10. A triangle is immersed in a fluid with one of its sides in the surface; find the position of a point within the triangle, such that, if it be joined to the angular points, the triangle shall be divided into three others, the fluid pressures upon which are equal.

11. The side AB of a triangle ABC is in the surface of a fluid, and a point D is taken in AC , such that the pressures on the triangles BAD , BDC , are equal; find the ratio $AD : DC$.

12. The lighter of two fluids, whose specific gravities are as 2.3, rests on the heavier, to a depth of four inches. A square is immersed in a vertical position with one side in the upper surface; determine the side of the square in order that the pressures on the portions in the two fluids may be equal.

13. A vertical cylinder contains equal portions of three inelastic fluids, the densities of which are ρ , 2ρ , and 3ρ , respectively, the lighter fluid being uppermost, and the heavier fluid lowest; compare the whole pressures on the portions of the curved surface of the cylinder in contact with the several fluids.

14. A fine tube, which is bent into the form of a circle, contains given quantities of two different liquids; if the two together occupy half the tube, determine the position of equilibrium.

15. The inclinations of the axis of a submerged solid cylinder to the vertical in two different positions are complementary to each other; P is the difference between the pressures on the two ends in the one, and P' in the other position: prove that the weight of the displaced fluid is equal to

$$(P^2 + P'^2)\dagger.$$

16. A vertical cylinder contains a quantity of fluid, whose depth equals a diameter of the circular base. A sphere of four times the specific gravity of the fluid and of the same radius as the cylinder is placed upon the fluid and is supported by it: find the increase of pressure sustained by the curved surface of the cylinder, the sphere fitting it exactly.

17. Three fluids whose densities are in arithmetic progression, fill a semicircular tube whose bounding diameter is horizontal. Prove that the depth of one of the common surfaces is double that of the other.

18. A small cylindrical tube is bent into a semicircle, and placed with the diameter horizontal; within the tube is placed a small stop which can slide freely up and down: two fluids of densities ρ and ρ' are poured into the respective ends of the tube;

if when the stop is vertical the surface of the lower fluid is $\frac{a}{\sqrt{2}}$ below the horizontal diameter, find the distance of the higher.

19. Prove that, as a plane area is lowered vertically in a liquid, the centre of pressure approaches to, and ultimately coincides with, the centre of gravity.

20. A lamina in the shape of a quadrilateral $ABCD$ has the side CD in the surface, and the sides AD , BC vertical and of lengths α , β , respectively. Prove that the depth of the centre of pressure is

$$\frac{1}{2} \cdot \left(\frac{\alpha^3 + \alpha^2\beta + \alpha\beta^2 + \beta^3}{\alpha^2 + \alpha\beta + \beta^2} \right).$$

21. A vessel contains two liquids whose densities are in the ratio of 1 to 14. A triangle is immersed vertically in the liquids so that its base is in the surface of the upper liquid. If the pressures on the portions in the two liquids be equal, prove that the areas of those portions are as 8 to 1.

22. The depth of the water on one side of a rectangular vertical floodgate is double that on the other. Supposing the gate to be fastened at the angular points, find the pressures at these points.

23. A vertical cylinder contains equal quantities of two liquids; compare their densities when the whole pressures of the two liquids on the curved surface of the cylinder are in the ratio 1 : 3.

24. If one second be the unit of time, what must be the unit of length in order that the formula $p = gpz$ may give the pressure in pounds, supposing the unit of volume of the standard substance to weigh 16 lbs. ?

25. If the density of distilled water be the unit of density, and 1 foot per second the unit of velocity, find the units of space and time, in order that the formula, $p = gpz$, may give the pressure in ounces.

26. If one yard be the unit of length, what must be the unit of time in order that the formula, $p = gpz$, may give the pressure in pounds, the weight of an unit of volume of the standard substance being 1000 lbs. ?

27. A sphere of 6 inches radius lies at the bottom of a pail of water, whose depth is 2 feet; find the numerical value of

the pressure on its surface, a foot being the unit of length, the density of water the unit of density, and one quarter of a second the unit of time.

28. A solid triangular prism, the faces of which include angles α , β , γ , is completely immersed in water with its edges horizontal; if P , Q , R , be the pressures on the three faces, which are respectively opposite to the angles α , β , γ , prove that

$$P \operatorname{cosec} \alpha + Q \operatorname{cosec} \beta + R \operatorname{cosec} \gamma$$

is invariable so long as the depth of the centre of gravity of the prism is unchanged.

29. A cubical vessel, standing on a horizontal plane, has one of its vertical sides loose, which is capable of revolving about a hinge at the bottom. If a portion of fluid equal in volume to one-fourth of the cube be poured into the vessel, the loose side will rest at an inclination of 45° to the horizon: compare the weight of the side with the weight of the fluid in the vessel.

30. A cubical box, filled with water, has a close fitting heavy lid fixed by smooth hinges to one edge; compare the tangents of the angles through which the box must be tilted about the several edges of its base, in order that the water may just begin to escape.


31. A cylindrical tumbler, containing water, is filled up with wine; after a time half the wine is floating on the top, half the water remains pure at the bottom, and the middle of the tumbler is occupied by wine and water completely mixed, the common surfaces being horizontal planes; if the weight of the wine be two-thirds of that of the water, and their densities be in the ratio of 11 : 12, prove that in this position the whole pressure of the pure water on the curved surface of the tumbler is equal to the whole pressure of the remainder of the liquid on the tumbler.

CHAPTER IV.

Resultant Vertical and Horizontal Pressure on any Surface, resultant Pressure on the Surface of an immersed Solid, Conditions of Equilibrium of a Floating Body, the Camel, Method of removing Wooden Piles, Stability of Equilibrium, Metacentre, Bodies floating in Air, the Balloon.

51. PROP. *To find the resultant vertical pressure of a liquid on any surface.*

Let PQ be a portion of surface in contact with a liquid




at rest, and through the boundary line of PQ draw vertical lines to the surface AB , thus enclosing a mass of the liquid.

The pressure of the surrounding liquid on this mass is entirely horizontal, and it is therefore clear that the weight of the mass is entirely supported by the reaction of the surface PQ .

Hence the vertical component of this reaction must be equal to the weight of the mass $ABQP$.

By the previous Chapter this is true whether the curve AB be really in the liquid, or only in the horizontal plane through the highest point of the liquid, as in the figure.




Hence it follows that *the resultant vertical pressure is the weight of the superincumbent liquid.*

52. There are other cases which it is requisite to consider.

Thus the liquid may press *upwards* on the surface.

In this case, let AB as before be the curve formed by vertical lines round PQ , and imagine the liquid within to be removed and the outside of PQ to be under the pressure of a fluid of which AB is the surface. It will be seen that the pressure at any point of PQ is the same as before in magnitude, but opposite in direction, and the resultant vertical pressure is therefore the same, only that it is now downwards, and by the previous article it is equal to the weight of $ABQP$.




Hence the resultant vertical pressure upwards on PQ is as before equal to the weight of the liquid above it, that is, between PQ and the surface.

Or the pressure may be partly upwards and partly downwards, as on PEQ .

Draw QQ' vertical, and consider the pressures on QEQ' and QP separately.


By the same reasoning the vertical pressure on QEQ' is downwards and equal to the weight of the liquid contained between the surface and the vertical plane QQ' , and the difference between this and the upward vertical pressure on PQ' is the resultant vertical pressure on the surface PQ .



In all cases the line of action of the resultant vertical pressure is the vertical through the centre of gravity of the superincumbent liquid.

53. PROP. *To find the resultant horizontal pressure in a given direction of a liquid on any surface.*

Take a fixed vertical plane perpendicular to the given direction, and draw horizontal lines through the boundary of the surface PQ , meeting the vertical plane in the curve AB . Considering the liquid thus enclosed as a solid body, its equilibrium is maintained by its own weight, by the fluid pressures on its curved surface which are all parallel to the vertical plane, and by the fluid pressures on the surfaces AB and PQ .




Hence the horizontal component of the reaction of PQ must be equal to the pressure on AB , which can be found from previous investigations, and the line of action will be the horizontal line through the centre of pressure of AB .

54. We are now in a position to determine the resultant pressure in direction and magnitude of a liquid on any surface; for we can obtain separately the vertical and horizontal pressures, and hence, by the principles of Statics, determine the magnitude and direction of the resultant.

Ex. 1. A vessel in the form of an open semi-cylinder with its ends vertical, is filled with water; it is required to find the resultant pressure on either of the portions into which it is divided by a vertical plane through the axis of the cylinder.

Let h be the length of the cylinder and a its radius, and let the figure be a vertical section through the middle point O of its length.



The resultant vertical pressure on AB

$$\begin{aligned} &= \text{the weight of the fluid } OAB \\ &= wh \frac{\pi a^2}{4}, \end{aligned}$$

if w be the weight of an unit of volume.

The resultant horizontal pressure on AB = the pressure on the vertical section perpendicular to the plane of the paper, that is, on a rectangle of which the sides are a and h ,

$$= wa h \frac{a}{2} = \frac{1}{2} wa^2 h.$$

Hence the angle θ , at which the direction of the resultant pressure is inclined to the horizon, is given by the equation

$$\tan \theta = \frac{\frac{1}{4} w \pi a^2 h}{\frac{1}{2} wa^2 h} = \frac{\pi}{2}.$$

Moreover, since the pressure at any point acts in a direction passing through the axis of the cylinder, the resultant pressure acts in a line through O , and, if $POB = \tan^{-1} \left(\frac{\pi}{2} \right)$, the point P is the centre of pressure of the curvilinear surface.

Ex. 2. A hollow cone filled with water is held with its vertex downwards; it is required to determine the resultant pressure on either of the portions into which it is divided by a vertical plane through its axis.

Let a be the radius of the base and $2a$ the vertical angle.


$$\text{The volume} = \frac{1}{3} \pi a^3 \cot \alpha.$$

The resultant vertical pressure on the portion $AEVB$

$$= \frac{1}{2} \text{ the weight of the fluid}$$

$$= \frac{1}{6} w \pi a^3 \cot \alpha,$$

if w be the weight of an unit of volume.



The resultant horizontal pressure

$$= \text{the pressure on the triangle } AVB$$

$$= w \cdot a^2 \cot \alpha \cdot \frac{1}{3} a \cot \alpha$$

$$= \frac{1}{3} w a^3 \cot^2 \alpha;$$

therefore the resultant pressure

$$= \frac{w}{3} a^3 \cot \alpha \sqrt{\frac{\pi^2}{4} + \cot^2 \alpha},$$

and if θ be the angle at which its direction is inclined to the horizon,

$$\tan \theta = \frac{\frac{1}{6} \pi}{\frac{1}{3} \cot \alpha} = \frac{\pi}{2} \tan \alpha.$$

In general the determination of the line of action can only be effected by means of the Integral Calculus, but in the first example we were able to infer at once the position of the line of action, and in some cases it may be determined by special geometrical contrivances.

As an example, the position of the line of action in this last case will be obtained in the appendix by the help of such a contrivance.

55. *To find the resultant pressure of a liquid on the surface of a solid either wholly or partially immersed.*

Imagine the solid removed, and the space it occupied in the liquid to be filled with the liquid, and suppose this liquid to be solidified. It is clear that the resultant pressure on this solidified liquid is the same as on the original solid. The weight of the liquid is entirely supported by the pressure of the surrounding liquid, and therefore the resultant pressure is equal to the weight of the displaced liquid, and acts vertically upwards in a line passing through its centre of gravity.

This is sometimes expressed by saying that *a solid immersed in fluid loses as much of its weight as is equal to the weight of the fluid it displaces*, observing that the above reasoning is equally applicable to the case of a body immersed in elastic fluid.

56. *To find the conditions of equilibrium of a floating body.*

By the previous article the resultant pressure is equal to the weight of the displaced liquid. It follows therefore that, the body being supported entirely by the liquid, the weight of the displaced liquid must be equal to the weight of the body, and the centres of gravity of both must be in the same vertical line.

These conditions also hold good when the body floats partly immersed in two or more liquids, and are, for such cases, established by precisely the same reasoning.

57. *If a homogeneous body float in a liquid, its volume will bear to the volume immersed the inverse ratio of the specific gravities of the solid and liquid.*

For if V , V' be the volumes, and s , s' the specific gravities,

$$\begin{aligned} Vs &= \text{the weight of the body} \\ &= \text{the weight of the displaced fluid} \\ &= V' s'; \end{aligned}$$

$$\therefore V : V' = s' : s.$$

58. *To find the conditions of equilibrium of a solid floating in liquid and partly supported by a string.*


First, let the solid be homogeneous and wholly immersed; then the centres of gravity of the solid and of the liquid displaced will be the same, and the direction of the string must be the vertical through the centre of gravity. Also

$$\begin{aligned} \text{the tension} &= \text{the weight of the body} - \text{the weight lost} \\ &= V(s - s') \end{aligned}$$

if s, s' be the specific gravities of the solid and fluid.

Secondly, let the solid be homogeneous and partly immersed.

Let V' be the part immersed, H its centre of gravity, and G the centre of gravity of the body.



Draw vertical lines through H and G meeting the surface in C and A , and let the direction of the string meet the surface in B .

Then, if T be the tension, the three forces, T, Vs , and $V' s'$ acting at B, A , and C will balance each other;

$$\therefore Vs = T + V' s',$$

and $Vs \cdot AB = V' s' \cdot CB.$

The second equation is the condition of equilibrium, and the first gives the requisite tension.

The case in which a heterogeneous body is partly supported by a string may be left for the consideration of the student.

59. *The Camel.* This is an apparatus for carrying a ship over the bar of a river. It consists of four, or a greater number of, watertight chests, which are filled with water, placed in pairs on opposite sides of the ship, and attached to the ship, or attached to each other by chains passing under the keel. If the water be then pumped out, the vessel will be lifted, and may be towed over the bar into deep water. The lifting power of the camel is the weight of water displaced by the chests, diminished by the weight of the whole apparatus.

60. *Removing wooden Piles.* It is sometimes necessary to remove entirely piles which have been driven down in deep water; for instance, the piles employed to keep out water during the construction of a dock. After the water has been allowed to flow within the piles, they are sawn off to a convenient depth, and a barge is floated over them and filled with water. The barge is then attached by chains to a pile, and the water pumped out; as the pumping proceeds the barge is lifted, and the pile is forcibly drawn out. If the operation take place in the sea, a great advantage is gained by fastening the barge to the pile at low tide. The rise of the tide will sometimes draw out the pile, but, if necessary, additional force must be gained by pumping water out of the barge.

61. We now proceed to exemplify the preceding propositions by their application to some particular cases.

Ex. 1. A man, whose weight is 150 lbs., and specific gravity 1.1, just floats in water, the specific gravity of which is 1, by the help of a quantity of cork. The specific gravity of cork being .24, find its volume in cubic feet.

Let V be the volume of the cork, and V' of the man, in cubic feet.

$$\begin{aligned} \text{Then } V (.24) + V' (1.1) &= \text{the weight of the water displaced} \\ &= V + V'; \end{aligned}$$

$$\text{or } V (.76) = V' (.1).$$

$$\begin{aligned} \text{But } V' (1.1) \frac{1000}{16} \text{ lbs.} &= \text{the weight of the man} \\ &= 150 \text{ lbs.}; \end{aligned}$$

$$\therefore V' = \frac{24}{11},$$

$$\text{and } V = \frac{.1}{.76} \times \frac{24}{11} = \frac{60}{209} \text{ ths of a cubic foot.}$$

Ex. 2. A cylindrical piece of wood floats in water with its axis vertical; find how much it will be depressed by placing a given weight on the top of it.

If w be the weight placed on the top, it will be depressed through such a space that the additional amount of displaced fluid has its weight equal to w .

Now, if W be the weight of the cylinder, it is also the weight of the fluid displaced by the cylinder, and therefore, if h be the depth of the base of the cylinder originally, and x the depression,

$$w : W :: x : h;$$

$$\therefore x = \frac{w}{W} h.$$

If this value of x exceed the height of the cylinder originally above the surface, it will be entirely immersed, and the possibility of equilibrium will then depend on the density of w .

Ex. 3. An isosceles triangular lamina floats in water with its base horizontal: it is required to find the position of equilibrium when the base is above the surface.

Take ρ' and ρ as the densities of the lamina and of water, h as the height of the triangle, and x the depth to which it is immersed.


Then ρ' (volume of lamina) = ρ (volume of fluid displaced); and therefore, similar triangles being proportional to the squares of homologous sides, we have

$$\rho' h^2 = \rho x^2, \quad \text{and } x = h \sqrt{\frac{\rho'}{\rho}}.$$

The second condition is obviously satisfied in this and the preceding example.

Ex. 4. Can an isosceles triangular lamina float with its base vertical in a liquid of twice its density?

The first condition requires that half the triangle should be immersed, and therefore its vertex A is in the surface.



Also, if G , H be the centres of gravity,

$$AG = \frac{2}{3} AE, \text{ and } AH = \frac{2}{3} AF, \text{ } F \text{ being the middle point of } EC;$$

$$\therefore AG : AH :: AE : AF.$$

Hence GH is parallel to EF , is therefore vertical, and both conditions are satisfied.

Ex. 5. A cylinder floats with its axis vertical, partly immersed in two liquids, the densities of the upper and lower liquids being respectively ρ and 2ρ , and the density of the cylinder $\frac{3\rho}{4}$; find the position of equilibrium of the cylinder, its length being twice the depth of the upper fluid.

Let x be the length immersed in the lower fluid, k the area of either end, and $2h$ the whole length.

$$\text{Then.} \quad \frac{3\rho}{4} k \cdot 2h = \rho kh + 2\rho kx;$$

$$\therefore x = \frac{1}{4} h.$$

If the cylinder were just immersed, its density ρ' would be such that

$$2\rho'h = \rho h + 2\rho h;$$

$$\text{or } \rho' = \frac{3\rho}{2},$$

and x would then be equal to h .

Ex. 6. A cubical box, the volume of which is one cubic foot, is three fourths filled with water, and a leaden ball, the volume of which is 72 cubic inches, is lowered into the water by

a string; it is required to find the increase of pressure on the base and on a side of the box.

The complete immersion of the lead will raise the surface $\frac{1}{4}$ an inch, since 144 square inches is the area of the surface.

The pressure on the base is therefore increased by the weight of 72 cubic inches of water, i. e. by $\frac{72}{1728} 1000$ oz., or $41\frac{2}{3}$ oz.

The area of a side originally in contact with the fluid was $\frac{3}{4}$ of a square foot,

and the pressure was $1000 \times \frac{3}{4} \times \frac{3}{8}$ oz., or $281\frac{1}{4}$ oz;

$\frac{3}{8}$ ths of a foot being the depth of the centre of gravity.

The new area is $\frac{3}{4} + \frac{1}{24}$, or $\frac{19}{24}$ of a square foot;


$$\therefore \text{the new pressure} = 1000 \times \frac{19}{24} \times \frac{19}{48} \text{ oz.}$$

$$= 313\frac{5}{144} \text{ oz.}$$

The increase is therefore a little more than 32 oz.

Ex. 7. A solid hemisphere is moveable about the centre of its plane base which is fixed in the surface of a liquid; if the density of the liquid be twice that of the solid, any position of the solid will be one of rest.

Hold the solid in the position *ADB*, *DE* being the surface of the liquid; continue the sphere to the surface *E* of the liquid, and imagine the portion of liquid within *CBE* to become solid and to be attached to the hemisphere. Make the angle *DCF* equal to *ECB*, the figure being a vertical section through the centre *C* of the hemisphere perpendicular to its plane base.



The wedge or lune *FCB* would be of itself in equilibrium; and, without knowing the position of the centre of gravity of a wedge, it is easily seen that the horizontal distance from *C* of

the centre of gravity of ECB is equal to that of the centre of gravity of ACF ; hence the moment about C of the weight of ECB is equal to the moment about C of the weight of ACF . Moreover the fluid pressures on the surface DBE all act in lines through C , and therefore, releasing the hemisphere, and restoring CBE to its liquid condition, the solid remains at rest.

The result of this problem has been practically employed in the construction of an oil-lamp, called Cecil's Lamp, such that the surface of the oil supplying the wick is always the same. DEB is a hemispherical vessel containing oil, and ADB a hemisphere, the specific gravity of which is half that of the oil; as the oil is consumed, ADB turns round C , and CE is always the surface of the oil.

Ex. 8. A solid hemisphere, completely immersed in liquid, of density ρ , is held so that the centre of its base is at a depth c below the surface, and the plane of its base inclined at an angle θ to the vertical; it is required to determine the resultant horizontal and vertical pressures on its curved surface.

Taking a for the radius, the resultant vertical pressure on its whole surface = the weight of the fluid displaced,

$$= \frac{2}{3} g \rho \pi a^3.$$

This resultant is the difference between the resultant vertical pressures on the curved surface and the plane base; but the pressure on the base = $g \rho \pi a^2 c$, in a direction inclined to an angle θ to the horizontal; and therefore the resultant vertical pressure on the base = $g \rho \pi a^2 c \sin \theta$.

Hence, if the base be turned upwards, the resultant vertical pressure on the curved surface

$$= \frac{2}{3} g \rho \pi a^3 + g \rho \pi a^2 c \sin \theta.$$

If the base be turned downwards, the vertical pressure on the curved surface

$$= g \rho \pi a^2 c \sin \theta - \frac{2}{3} g \rho \pi a^3.$$

Also the horizontal pressure on the curved surface

= the horizontal pressure on the base

$$= g \rho \pi a^2 c \cos \theta.$$

Hence the actual resultant pressure on the curved surface


$$= g\rho\pi a^2 \sqrt{c^2 \pm \frac{4}{3} ac \sin \theta + \frac{4}{9} a^2}.$$

It will be seen that the method of this example can be applied to find the resultant pressure on the surface of any solid bounded by a plane of known area, if the volume of the solid be known.

Stability of Equilibrium.

62. Imagine a floating body to be slightly displaced from its position of equilibrium by turning it round so that the line joining its centre of gravity with that of the fluid displaced shall be inclined to the vertical. If the body on being released return to its original position its equilibrium is stable; if, on the other hand, it fall away from that position its original position is said to be one of unstable equilibrium.

Metacentre. In the figure let G , H be the centres



of gravity of the body and of the fluid originally displaced, H' the centre of gravity of the fluid displaced in the new position, and M the point where a vertical through H' meets HG .

The resistance of the fluid acts vertically upwards in the line $H'M$, and it is therefore evident that, if M be above G , the action of the fluid will tend to restore the body to its original position; but, if M be below G , to turn the body farther from its original position.

The position of the point M will in general depend on the extent of displacement. If the displacement be very

small, that is, if the angle between GH and the vertical be very small, the point M is called the *metacentre*, and the question of stability is now reduced to the determination of this point.


One of the most important problems in naval architecture is to secure the ascendancy under all circumstances of the metacentre over the centre of gravity.

This is effected by a proper form of the midship sections, so as to raise the metacentre as much as possible, and by ballasting so as to lower the centre of gravity, and the greater the distance between the points G and M , the greater is the steadiness of the vessel.

Moreover, the naval architect must have in view the probability of large displacements, due to the rolling of the vessel, and not merely the small movement which is considered in the determination of the metacentre.

63. In particular cases the metacentre can be sometimes found by elementary methods, but its general determination involves the application of the Integral Calculus.

In one case however its position is obvious. Let the lower portion of the solid be spherical in form; then as long as the portion immersed is spherical, the pressure of the water at every point acts in the direction of the centre of the sphere, and therefore the resultant pressure must act in the vertical line through the centre (E) of the sphere.



Now in the original position the centre of gravity of

the fluid displaced is evidently in the vertical through *E*, and therefore the centre of gravity of the body is in the vertical through *E*.

Hence the point *E* is the metacentre.

Thus if any portion whatever be cut from a solid spheroid it will float in stable equilibrium with its curved surface partly immersed.

64. *Bodies floating in air.*

The fact that air is heavy enables us to extend to bodies, floating either wholly or partly in air, the laws of equilibrium which have been established for bodies floating in liquids.

Taking one case, if a body, lighter than water, float on its surface, it displaces a certain quantity of water and also a certain quantity of air; if we remove the body and suppose its place filled by air and water, it is clear that the weight of the displaced air and water is supported by the resultant vertical pressures of the air and water around it.

Hence the weight of the body must be equal to the weight of the air and water it displaces, and the centre of gravity of the air and water displaced must be in the same vertical line with the centre of gravity of the body.

In a similar manner, if a body float in air alone, its weight must be equal to the weight of the air it displaces.


65. *The Balloon.* The ascent of a balloon depends on the principle of the previous article. A balloon is a very large envelope, made of silk, or some strong and light substance, and filled with a gas of less density than the air, usually coal gas. A car is attached in which the aeronauts are seated, and the weight of air displaced being greater than the whole weight of the balloon and car, the balloon ascends, and will continue ascending until the air around is not of sufficient density to support its weight. In order to descend, a valve is opened and a portion of the gas allowed to escape.

The ascensional force on a balloon is the weight of the

air it displaces diminished by the weight of the balloon itself.

66. If a body float in a liquid, the centre of gravity of the liquid displaced is called the *Centre of Buoyancy*.

If the body be moved about, so that the volume of liquid displaced remains unchanged, the locus of the centre of the displaced liquid is called the *Surface of Buoyancy*.



Taking a simple case, suppose a triangular lamina immersed with its plane vertical, and vertex beneath the surface, and let the area APQ be constant. Through H the centre of gravity of the area APQ , draw pHq parallel to PQ ; then the area Apq is constant, and therefore pq always touches, at its middle point H , an hyperbola of which AB and AC are the asymptotes. This hyperbola is the curve of buoyancy. Now in the position of equilibrium, GH is vertical, and is consequently perpendicular to pq .

The position of equilibrium is therefore determined by drawing normals from G to the curve of buoyancy.

67. It is a general theorem that *the positions of equilibrium of a floating body are determined by drawing normals from the centre of gravity of the body to the surface of buoyancy*.

We give a proof for the case of a lamina with its plane vertical, or, which is the same thing, of a prismatic or cylindrical body with its flat ends vertical.

EXAMINATION UPON CHAPTER IV.

1. Shew how to find the resultant vertical pressure of a liquid on a surface; 1st, when it acts upwards, 2nd, when it acts downwards.

2. Apply the preceding to find the resultant pressure on a solid completely immersed.

3. A solid cone of metal, completely immersed in liquid, is supported by a string; find the tension of the string.

4. State the conditions of equilibrium of a floating body.

X 5. A wooden plank floats in water, and a weight is placed at one end of the plank; find the weight which, placed at a given distance from the other end, will keep the plank in a horizontal position.

6. Describe a method of removing piles in deep water.

7. A cylinder floats vertically in a fluid with 8 feet of its length above the fluid; find the whole length of the cylinder, the specific gravity of the fluid being three times that of the cylinder.

8. A body floats in one fluid with $\frac{3}{4}$ ths of its volume immersed, and in another with $\frac{4}{5}$ ths immersed; compare the specific gravities of the two fluids.

9. A cylinder of wood 3 feet in length floats with its axis vertical in a fluid of twice its specific gravity; compare the forces required to raise it 6 inches and to depress it 6 inches.

10. Three equal rods are jointed together so as to form an equilateral triangle, and the system floats in a liquid of twice the density of the rods, with one rod horizontal and above the surface; find the position of equilibrium.

11. Explain what is meant by stability of equilibrium, and define the metacentre.

12. A small iron nail is driven into a wooden sphere, and the weight of the sphere is then half that of an equal volume of water; find its positions of equilibrium in water, and examine the stability of the equilibrium.

13. A block of wood, the volume of which is 4 cubic feet, floats half immersed in water; find the volume of a piece of metal, the specific gravity of which is 7 times that of the wood, which, when attached to the lower portion of the wood, will just cause it to sink.

14. A cylindrical block of wood is placed with its axis vertical in a cylindrical vessel whose base is plane, and water is then poured in to twice the height of the cylinder; find the pressure of the wood on the base of the vessel.

15. Two cylindrical vessels, containing different fluids, and standing near each other on a horizontal plane, are connected by a fine tube, which is close to the horizontal plane; when the communication is opened between them, determine which of the fluids will flow from its own vessel into the other, and find the condition that the equilibrium may not be disturbed.

16. Two bodies of given size and given specific gravities are connected by a string passing over a pulley, and rest completely immersed in water; find the condition of equilibrium.

NOTE ON CHAPTER IV.

The Principle of Archimedes. The enunciation and proof of the proposition of Article (56) are due to Archimedes, and it is a remarkable fact in the history of science, that no further progress was made in Hydrostatics for 1800 years, and until the times of Stevinus, Galileo, and Torricelli, the clear idea of fluid action thus expounded by Archimedes remained barren of results.

An anecdote is told of Archimedes, which practically illustrates the accuracy of his conceptions. Hiero, king of Syracuse, had a certain quantity of gold made into a crown, and suspecting that the workman had abstracted some of the gold and used a portion of alloy of the same weight in its place, applied to Archimedes to solve the difficulty. Archimedes, while reflecting over this problem in his bath, observed the water running over the sides of the bath, and it occurred to him that he was displacing a quantity of water equal to his own bulk, and therefore that a quantity of pure gold equal in weight to the crown, would displace less water than the crown, the volume of any weight of alloy being greater than that of an

equal weight of gold. It is related that he immediately ran out into the streets, crying out *εὑρηκα! εὑρηκα!*

The two books of Archimedes which have come down to us, "*De iis quæ in humido vehuntur*" were first found in an old Latin MS. by Nicholas Tartaglia, and edited by him in 1537. In the first of these books it is shewn that the surface of water at rest must be a sphere of which the centre is at the earth's centre, and various problems are then solved relating to the equilibrium of portions of spherical bodies. The second book contains the proposition of Art. (56), and the solutions of a number of problems on the equilibrium of paraboloids, some of which involve complicated geometrical constructions.

The authenticity of these books is confirmed by the fact that they are referred to by Strabo, who not only mentions their title, but also quotes the second proposition of the first book.

Stevinus and Galileo. The Treatise of Stevinus on Statics and Hydrostatics, about 1585, follows that of Archimedes in the order of thought. In this he shewed how to determine the pressure of a liquid on the base and sides of a vessel containing it.

Galileo, in his Treatise on Floating Bodies, published in 1612, states the Hydrostatic paradox, and explains why the floating of bodies does not depend on their form.

EXAMPLES.

1. A uniform solid floats freely in a fluid of specific gravity twice as great as its own; prove that it will also float in equilibrium, if its position be inverted.

2. A block of ice, the volume of which is a cubic yard, is observed to float with $\frac{2}{5}$ ths of its volume above the surface, and a small piece of granite is seen embedded in the ice; find the size of the stone, the specific gravities of ice and granite being respectively .918 and 2.65.

3. An isosceles triangular lamina floats with its base horizontal and beneath the surface of a liquid of twice its density; find the position of equilibrium.

4. A solid cone floats with its axis vertical in a liquid the density of which is twice that of the cone; compare the portions of the axis immersed, 1st, when the vertex is upwards, 2nd, when it is downwards.

5. If w_1, w_2, w_3 be the apparent weight of a body in three liquids, the specific gravities of which are s_1, s_2, s_3 , prove that

$$w_1 (s_2 - s_3) + w_2 (s_3 - s_1) + w_3 (s_1 - s_2) = 0.$$

6. An equilateral triangular lamina suspended freely from A , rests with the side AB vertical, and the side AC bisected by the surface of a heavy fluid; prove that the density of the lamina is to that of the fluid :: 15 : 16.

7. A vertical cylinder of density $\frac{7\rho}{4}$ floats in two liquids, the density of the upper liquid being ρ , and of the lower 2ρ ; if the length of the cylinder be twice the depth of the upper liquid, find its position of rest.

8. A wooden rod is tipped with lead at one end; find the density of a liquid in which it will float at any inclination to the vertical; the weight of the lead being half that of the rod, and its size being neglected.

9. The weight of the unimmersed portion of a body floating in water being given, find the specific gravity of the body, in order that its volume may be the least possible.

10. A cylindrical glass cup weighs 8 oz., its external radius is $1\frac{1}{2}$ inches, and its height $4\frac{1}{2}$ inches; if it be allowed to float in water with its axis vertical, find what additional weight must be placed in it, in order that it may sink.

11. A vessel in the form of half the above cylinder with both its ends closed, floats in water, with its ends vertical; find the additional weight which being placed in the centre of the vessel will cause it to be totally immersed.

12. A uniform rod, whose weight is W , floats in water in a position inclined to the vertical with a particle, of weight W , attached to its lower end; shew that, if the density of the water be four times that of the rod, half the length of the rod will be immersed.

13. A uniform rod floats partly immersed in water, and supported at one end by a string; prove that, if the length immersed remain unaltered, the tension of the string is independent of the inclination of the rod to the vertical.

14. A spherical shell, the internal and external radii of which are given, floats half immersed in water; find its density compared with the density of water.

15. A heavy hollow right cone, closed by a base without weight, is completely immersed in a fluid, find the force that will sustain it with its axis horizontal.

16. Find the position of equilibrium of a solid cone, floating with its axis vertical and vertex upwards, in a fluid of which the density bears to the density of the cone the ratio 27 : 19.

17. A rectangular lamina $ABCD$ has a weight attached to the point B , and floats in water with its plane vertical and the diagonal AC in the surface; prove that the specific gravity of the fluid is three times that of the lamina.

18. A solid paraboloid floats in a liquid with its axis vertical and vertex downwards; having given the densities of the paraboloid and the liquid, find the depth to which the vertex is submerged.

19. A ship sailing from the sea into a river sinks two inches, but after discharging 40 tons of her cargo, rises an inch and a half; determine the weight of the ship and cargo together, the specific gravity of sea-water being 1.025, and the horizontal section of the ship for two inches above the sea being invariable.

20. A cylindrical vessel of radius r and height h is three-fourths filled with water; find the largest cylinder of radius r' and specific gravity .5 which can be placed in the water without causing it to run over, the axes of the cylinders being vertical and r' less than r .

21. A hollow cylinder is just filled with water, and closed, and is then held with its axis horizontal; find the direction and magnitude of the resultant pressure on the lower half of the curved surface. Also, if the cylinder be held with its axis vertical, find the direction and magnitude of the resultant pressure on the same surface.

22. A solid cylinder, one end of which is rounded off in the form of a hemisphere, floats with the spherical surface partly immersed: find the greatest height of the cylinder which is consistent with stability of equilibrium.

23. A body floating on an inelastic fluid is observed to have volumes P_1, P_2, P_3 respectively above the surface at times when the density of the surrounding air is ρ_1, ρ_2, ρ_3 ; shew that

$$\frac{\rho_2 - \rho_3}{P_1} + \frac{\rho_3 - \rho_1}{P_2} + \frac{\rho_1 - \rho_2}{P_3} = 0.$$

24. A frustum of a right circular cone, cut off by a plane bisecting the axis perpendicularly, floats with its smaller end in a fluid and its axis just half immersed; compare the densities of the cone and fluid.

25. A solid cone and a solid hemisphere, which have their bases equal, are united together, base to base, and the solid thus formed floats in water with its spherical surface partly immersed; find the height of the cone in order that the equilibrium may be neutral.

26. Three uniform rods, joined so as to form three sides of a square, have one of their free extremities attached to a hinge in the surface of a fluid, and rest in a vertical plane with half the opposite side out of the fluid; shew that the specific gravity of the rods is to that of the fluid as 31 : 40.

27. A triangle ABC floats in a fluid with its plane vertical, the angle B being in the surface of the fluid and the angle A not immersed. Shew that the density of fluid : density of the triangle :: $\sin B$: $\sin A \cos C$.

28. A solid cone floats with its axis vertical and vertex downwards in an inelastic fluid; prove that, whatever be the density of the fluid, supposing it greater than that of the solid, the whole pressure on its curved surface is the same.

29. Two fluids are in equilibrium, one upon the other, the lower fluid having the greater specific gravity, and a solid cylinder, the height of which is equal to the depth of the upper fluid, is immersed with its axis vertical: the specific gravity of the cylinder being greater than that of the upper fluid, find the position of equilibrium.

What would be the effect of an increase in the density of the upper fluid? Will the equilibrium be stable or unstable for a vertical displacement?

30. Two equal uniform rods AB , BC are freely jointed at B , and are capable of motion about A , which is fixed at a given depth below the surface of a uniform heavy fluid. Find the position in which both rods will rest partly immersed, and shew that, in order that such a position may be possible, the ratio of the density of the rods to the density of the fluid must be less than $\frac{5}{9}$.

31. An equilateral triangle, ABC , of weight W and specific

gravity σ , is moveable about a hinge at A , and is in equilibrium when the angle C is immersed in water and the side AB is horizontal. It is then turned about A in its own plane until the whole of the side BC is in the water and horizontal; prove that the pressure on the hinge in this position

$$= 2 \frac{1 - \sqrt{\sigma}}{\sqrt{\sigma}} W.$$

32. A solid hemisphere is completely immersed with the centre of its base at a given depth; if W be the weight of fluid it displaces, P the resultant vertical pressure, and Q the resultant horizontal pressure, on its curved surface, prove that for all positions of the solid $(W - P)^2 + Q^2$ is constant.

33. A hollow cone, filled with water and closed, is held with its axis horizontal; find the resultant vertical pressure on the upper half of its curved surface.

34. A solid cylinder which is completely immersed in water has its centre of gravity at a given depth below the surface, and its axis inclined at a given angle to the vertical; determine the resultant horizontal and resultant vertical pressures upon its curved surface, and the direction and magnitude of the resultant pressure on the curved surface.

35. The vertical angle of a solid cone is 60° ; prove that it can float in a liquid with its vertex above the surface and its base touching the surface, if the densities of the cone and the liquid are in the ratio of $2\sqrt{2} - 1 : 2\sqrt{2}$.

36. A thin hollow cone closed by an equally thin base will remain wherever it is placed entirely within a liquid; prove that its vertical angle is $2 \sin^{-1} \frac{1}{3}$.

37. The base of a vessel containing water is a horizontal plane, and a sphere of less density than water is kept totally immersed by a string fastened to the centre of a circular disc, which lies in contact with the base. Find the greatest sphere of given density, and also the sphere of given size and least density, which will not raise the disc.

CHAPTER V.

ON AIR AND GASES.

Elasticity of Air, Effect of Heat, Thermometers, Torricelli's Experiment, Weight of Air, the Barometer and its Graduation, the Relations between Pressure, Density, and Temperature, Determination of Heights by the Barometer, the Siphon, Graduation of a Thermometer, the Differential Thermometer.

68. **T**HE pressure of an elastic fluid is measured exactly in the same way as the pressure of a liquid, and it has been mentioned before that the properties of equality of pressure in all directions and of transmission of pressure are equally true of liquids and gases.

There is however this difference between a gas and a liquid, that the pressure of the latter is entirely due to its weight, or to the application of some external pressure, while the pressure of a gas, although modified by the action of gravity, depends in chief upon its volume and temperature.

The action of a common syringe will serve to illustrate the elasticity of atmospheric air. If the syringe be drawn out and its open end then closed, a considerable effort will be required to force in the piston to more than a small fraction of the length of its range, and if the syringe be air-tight, and strong enough, it will require the application of very great power to force down the piston through nearly the whole of its range. Moreover, this experiment with a syringe shews that the pressure increases with the

compression, the air within the syringe acting as an elastic cushion. If the piston after being forced in be let go, it will be driven back, the air within expanding to its original volume.

Another simple illustration may be obtained by immersing carefully in water an inverted glass cylinder. Holding the cylinder vertical, fig. Art. 94, Ex. (2), it may be pressed down in the water without much loss of air, and it will be seen that the surface of the water within the vessel is below the surface of the water outside. It is evident that the pressure of the air within is equal to the pressure of the water at its surface within the cylinder, which, as we have shewn before, is equal to the pressure at the outside surface, increased by the pressure due to the depth of the inner surface; hence the air within, which has a diminished volume, has an increased pressure.

69. *Effect of heat.* It is found that if the temperature be increased, the elastic force of a quantity of air or gas which cannot change its volume is increased, but that if the air can expand, while its pressure remains the same, its volume will be increased.

To illustrate this, imagine an air-tight piston in a vertical cylinder containing air, and let it be in equilibrium, the weight of the piston being supported by the cushion of air beneath.

Raise the temperature of the air in the cylinder; the piston will then rise, or, if it be not allowed to rise, the force required to keep it down will increase with the increase of temperature.

70. *Thermometer.* As a general rule bodies expand under the action of heat, and contract under that of cold, and the only method of measuring temperatures is by observing the extent of the expansion or contraction of some known substance.

For all ordinary temperatures mercury is employed, but for very high temperatures a metal of some sort is the most useful, and for very low temperatures, at which mercury freezes, alcohol must be employed.

71. *The Mercurial Thermometer* is formed of a thin glass tube terminating in a bulb, and having its upper end hermetically sealed. The bulb contains mercury which also extends partly up the tube, and the space between the mercury and the top of the tube is a vacuum.

It must be observed that, as the glass expands with an increase of temperature, as well as the mercury, the apparent expansion is the difference between the actual expansion and the expansion of the glass.

In the Centigrade Thermometer the freezing point is marked 0° , and the boiling point 100° , the space between being divided into 100 equal parts, called degrees.

In Fahrenheit's Thermometer the freezing point is marked 32° , and the boiling point 212° ; and in Reaumur's the freezing point is 0° , and the boiling point 80° .



72. *To compare the scales of these Thermometers.*

Let C , F and R be the numbers of degrees marking the same temperature on the respective thermometers; then, since the space between the boiling and freezing points must in each case be divided in the same proportion by the mark of any given temperature, we must have

$$C : F - 32 : R :: 100 : 180 : 80$$

$$:: 5 : 9 : 4,$$


$$\text{or } \frac{C}{5} = \frac{F - 32}{9} = \frac{R}{4},$$

it being taken for granted that the temperature indicated by the boiling point is the same in all.

The method of filling the thermometer, and the definitions of the freezing and boiling points, will be given at the end of the chapter.

73. *Pressure of the Atmosphere. Torricelli's Experiment.*

The action of the atmosphere was distinctly ascertained by the experiment of Torricelli. Taking a glass tube *AB*,



32 or more inches in length, open at the end *A* and closed at the end *B*, he filled it with mercury, and then, closing the end *A*, inverted the tube, immersed the end *A* in a cup of mercury, and then opened the end *A*. The mercury was observed to descend through a certain space, leaving a vacuum at the top of the tube, but resting with its surface at a height of about 29 or 30 inches above the surface of the mercury in the cup.

It thus appears that the atmospheric pressure, acting on the surface of the mercury in the cup, and transmitted, as we have shewn that such pressures must be transmitted, supports the column of mercury in the tube, and provides us with the means of directly measuring the amount of the atmospheric pressure.

In fact, the weight of the column of mercury in the

tube above the surface in the cup, is exactly equivalent to the atmospheric pressure on an area equal to that of the section of the tube. This is about 15lbs. on a square inch.

74. *Air has weight.* This may be directly proved by weighing a flask filled with air; and afterwards weighing it, when the air has been withdrawn by means of an air-pump. The difference of the weights is the weight of the air contained by the flask.

We are now in a position to account for the fact of atmospheric pressure. The earth is surrounded by a quantity of air, the height of which is limited, as may be proved by dynamical and other considerations; and if, above any horizontal area, we suppose a cylindrical column extending to the surface of the atmosphere, the weight of the column of air must be entirely supported by the horizontal area upon which it rests, and the pressure upon the area is therefore equal to the weight of the column of air.

According to this theory the pressure of the air must diminish as the height above the earth's surface increases, and, from experiments in balloons, and in mountain ascents, this is found to be the case. As before, taking Π for the pressure at any given place, and ρ as the density of the air, the pressure at a height z will be

$$\Pi - g\rho z,$$

if we assume that the density of the air is sensibly the same through the height z .

75. It has been mentioned that the pressure of a gas depends chiefly upon its volume and temperature, but it is implied in that statement that the gas is confined within a limited space, for without such a restriction the effect of its elasticity might be the unlimited expansion and ultimate dispersion of the gas.

The action of gravity is equivalent to the effect of a compression of the gas, and it is thus seen that the pressure of a gas is in fact due to its weight, as in the case of a liquid.

76. It may be shewn in the same manner as for air that any other gas has weight, and that the intrinsic weight is in general different for different gases.

Carbonic acid gas, for instance, is heavier than air, and this is illustrated by the fact that it can be poured, as if it were liquid, from one jar to another.

The Barometer.

77. This instrument, which is employed for measuring the pressure of the atmosphere, consists of a bent tube ABC , closed at A , and having the end C open.

The height of the portion AB is usually about 32 or 33 inches, and the portion BC is generally for convenience of much larger diameter than AB . The tube contains a quantity of mercury, and the portion AP above the mercury is a vacuum.

If the plane of the surface in BC intersect AB in Q , it is clear, since the pressure at all points of a horizontal plane is the same, that the pressure at Q is the same as the atmospheric pressure, which is transmitted from the surface at C to Q , and therefore the atmospheric pressure supports the column of mercury PQ . Hence the height of this column is a measure of the atmospheric pressure, and if σ be the density of mercury, and Π the atmospheric pressure,

$$\Pi = g\sigma PQ.$$

The density of mercury diminishes with an increase of temperature, and it is an experimental result that, for an increase of 1° centigrade, the expansion of mercury is $\frac{1}{5550}$ th, or .00018018 of its volume; and therefore if σ_t be the density at a temperature t , and σ_0 at a temperature 0° ,

$$\sigma_0 = \sigma_t(1 + .00018018t),$$

$$\text{or, if } \theta = .00018018, \sigma_0 = \sigma_t(1 + \theta t);$$

$$\text{and } \therefore \Pi = g\sigma_t \cdot PQ = g\sigma_0(1 - \theta t)PQ.$$

78. The *average height* of the barometric column at the level of the sea is found to vary with the latitude, but it is generally between $29\frac{1}{2}$ and 30 inches. This height is however subject to continuous variations; during any one day there is an oscillation in the column, and the mean



height for one day is itself subject to an annual oscillation, independently of irregular and rapid oscillations due to high winds and stormy weather. Usually the height of the column is a maximum about 9 in the morning; it then descends until 3 P.M., and again attains a maximum at 9 in the evening.

79. *The Water Barometer.* Any kind of liquid will serve to measure the atmospheric pressure, but the great density of mercury renders it the most convenient for the purpose. If water were employed, it would be necessary to have a tube of great length; in fact, as the density of mercury is about 13.568 times that of water, the height of the column of water would be about $33\frac{1}{2}$ feet.

80. *Graduation of the Barometer.* Suppose the column of mercury to rise above P (fig. Art. 77); then it is clear that it descends below C in BC , and that the variation in the height of the column is the sum of these two changes.

Let k , K be the sectional areas of the tubes, and x the ascent above P , or the apparent variation; then the descent below C is $\frac{kx}{K}$, and the true variation is

$$x + \frac{kx}{K}, \text{ or } \left(1 + \frac{k}{K}\right)x.$$

Hence in graduation the distances actually measured from the zero point must be marked larger in the ratio of

$$1 + \frac{k}{K} : 1.$$

81. *To find the atmospheric pressure on a square inch.*

This we can determine at once by observing that it is the weight of a cylindrical column of mercury of which the base is a square inch and the height equal to that of the barometric column.

The specific gravity of mercury is 13.568 times that of water; hence the atmospheric pressure on a square inch, taking 30 inches as the height of the barometer at the sea level,

$$\begin{aligned} &= 30 \times 13.568 \times 1000 \div 1728 \text{ oz.} \\ &= 14.7 \text{ lbs.} \end{aligned}$$

This pressure varies from time to time, but is generally between $14\frac{1}{2}$ and 15 lbs.

82. *The height of the homogeneous atmosphere.*

If the density of the atmosphere were the same throughout the whole vertical column as it is at the sea level, its height would be less than 5 miles.


To prove this, let σ , ρ be the densities of mercury and of air, each referred to water; then, if h be the height of the barometer, the atmospheric pressure $= \sigma h$. Hence the height of the atmospheric column would be $\frac{\sigma}{\rho} h$. Now, it has been found that the ratio $\sigma : \rho$ is about 10462 : 1, and if we take h to be 30 inches, we shall find that $\frac{\sigma}{\rho} h$ is a little less than 5 miles.

83. *The pressure of a given quantity of air, at a given temperature, varies inversely as the space it occupies.*

The experimental proof of this law, due to Boyle and Marriotte, is as follows.

A bent glass tube, the shorter branch of which can have its end closed, is fixed to a graduated stand. Both ends being open, a little mercury is poured in, which rests with its surfaces P, P in the same horizontal plane. The end A is now closed and more mercury is poured in at B ; the effect is a compression of the air in AP , the mercury rising to a height Q , which is however below the surface R of the mercury in BP .

After closing the end A the pressure of the air is equal to the atmospheric pressure, and when more mercury has been poured in, the pressure of the air in AQ is equal to that of the mercury at Q , the same level in the longer branch. This latter pres-



sure is due to the atmospheric pressure on the surface R , and the weight of the column RQ .

If now the spaces AQ , AP be compared, which may be effected by comparing the weights of the mercury they would contain, and if the height h of the barometer be observed, it will be found that

$$\frac{\text{space } AP}{\text{space } AQ} = \frac{h + QR}{h}.$$


But, taking Π as the original pressure of the air in AP , and Π' as its pressure when compressed,

$$\Pi = g\sigma h, \text{ and } \Pi' = \Pi + g\sigma RQ = g\sigma (h + RQ);$$

$$\therefore \Pi' : \Pi :: \text{space } AP : \text{space } AQ,$$

and this proves the law for a *compression* of air.

For a dilatation, employ a bent glass tube, of which both branches are long, and pour in mercury to a height P ; then close the end A and withdraw some of the mercury from the branch B ; let Q and R be the new surfaces.



It will now be found that

$$\frac{\text{space } AP}{\text{space } AQ} = \frac{h - QR}{h}.$$

But if Π'' be the pressure of the air when dilated,

$$\begin{aligned} \Pi'' &= \text{pressure at } R - g\sigma QR \\ &= g\sigma (h - QR); \end{aligned}$$

And $\therefore \Pi'' : \Pi :: \text{space } AP : \text{space } AQ.$

In each case care must be taken to have the temperatures the same at the beginning and at the conclusion of the experiment.

Hence it follows that, since the density of a given quantity of air varies inversely as its volume, the pressure varies directly as the density. If p be the pressure, and ρ the density, this is expressed by the equation

$$p = k\rho,$$


where k is a quantity to be determined by experiment.

84. *Effect of a change of temperature.*

If the pressure remain constant, an increase of temperature of 1° centigrade, produces in a given mass of air an expansion .003665 of its volume.

This experimental law, combined with the preceding, enables us to express the relation between the pressure, density, and temperature of a given mass of air or gas.

Imagine a quantity of air confined in a cylinder by a piston to which a given force is applied, and let the



temperature be 0° C. Raise the temperature to t° : the piston will then be forced out until the original volume (V_0) is increased by .003665 $t \cdot V_0$ or $a t V_0$, designating the decimal by a . If V be the new volume, we have

$$V = V_0(1 + at),$$

and therefore, if ρ , ρ_0 be the densities at the temperatures t° , 0° , $\rho_0 = \rho(1 + at)$.

Hence,
$$p = k\rho_0 = k\rho(1 + at).$$

85. *Absolute Temperature.*

If we can imagine the temperature of a gas lowered until its pressure vanishes, without any change of volume, we arrive at what is called the absolute zero of temperature.

Assuming t_0 to be this temperature on the centigrade scale, we have

$$1 + at_0 = 0,$$

or

$$t_0 = -273^{\circ}.$$

In Fahrenheit's scale this is -459° .

Hence $p = k\rho(1 + at) = k\rho a(t - t_0) = k\rho a T$,
if T be the absolute temperature.

Taking V as the volume of the gas, ρV is constant, and therefore $\frac{pV}{T}$ is constant; *i.e.* the product of the pressure and volume is proportional to the absolute temperature.

The *air Thermometer* is a long straight tube of uniform bore closed at its lower end, open at the upper end, and containing air or some other gas, which is separated from the external air by a short column of liquid.

This thermometer is very sensitive, but it has the disadvantage that, as the atmospheric pressure is variable, no estimation can be made of the temperature without at the same time taking account of the height of the barometer*.

86. *Illustration.* The effect of heat in the expansion of air may be illustrated by a simple experiment.

Take a glass tube, open at one end, and ending in a bulb at the other; immerse the open end in water, and then apply the heat of a lamp to the bulb. The air in the bulb will expand, and will drive out a portion of the water in the tube.



If the lamp be removed, the air within will be cooled, and the water will then rise in the tube.

87. *Determination of heights by the barometer.*

It is found both from theory and from observation, that the height of the barometric column depends on its altitude above the sea level, and we are thus provided with a means of directly inferring from observation the height of any given station above the level of the sea.

For this purpose it is necessary to construct a formula which shall connect the height of the barometer with the height of its position above a given level, such as the sea level.

A general formula would be somewhat complicated, and difficult to obtain without the aid of the Integral Calculus, since the atmospheric pressure depends on the temperature and density of the air, which both vary with the height, and also on the intensity of gravity, which diminishes with an increase of height.

We shall however construct a formula on the supposition that the temperature and the force of gravity re-

* See Chapter II. of Maxwell's Heat.

main constant: this will be practically useful for the determination of comparatively small differences of altitude.

88. *If a series of heights be taken in arithmetic progression, the densities of the air decrease in geometric progression.*

Take a vertical column of the atmosphere of a given height z , and let it be divided into n horizontal layers of the same thickness, i. e. $\frac{z}{n}$, and suppose that $\rho_1, \rho_2, \rho_3 \dots \rho_n$ represent the densities of the successive layers, measuring upwards.

These layers may be supposed each of the same density throughout, and, if we take the temperature the same in all, the pressures on the upper sides of the layers will be

$$k\rho_1, k\rho_2, \dots, k\rho_n,$$

k being the constant of variation for the particular temperature.

The difference between any two consecutive pressures must be equal to the weight of the air between them, and therefore, taking the $(r-1)^{\text{th}}$ and r^{th} pressures,

$$k\rho_{r-1} - k\rho_r = g\rho_r \frac{z}{n},$$

$$\text{or } k\rho_{r-1} = \left(k + g\frac{z}{n}\right)\rho_r;$$

$$\therefore \frac{\rho_{r-1}}{\rho_r} = 1 + \frac{gz}{kn},$$

that is, the densities diminish in geometric progression.

89. *To find an expression for the difference of the altitudes of two stations.*

If z be this difference, we have from the preceding article, putting γ for

$$1 + \frac{gz}{kn},$$

and ρ_0 for the density immediately beneath the lowest layer of air,

$$\rho_{n-1} = \gamma \rho_n, \quad \rho_{n-2} = \gamma \rho_{n-1} \dots \rho_1 = \gamma \rho_2, \quad \rho_0 = \gamma \rho_1,$$

and therefore,

$$\rho_0 = \gamma^n \rho_n.$$

Hence, if p' , p be the corresponding pressures

$$p = \gamma^n p'.$$

Let h' , h be the observed altitudes of the barometer at the higher and lower stations respectively.

Then
$$\frac{h}{h'} = \frac{p}{p'} = \gamma^n = \left(1 + \frac{gz}{kn}\right)^n.$$

And
$$\begin{aligned} \log_e \frac{h}{h'} &= n \log \left(1 + \frac{gz}{kn}\right) \\ &= n \left(\frac{gz}{kn} - \frac{1}{2} \frac{g^2 z^2}{k^2 n^2} + \dots \right) \\ &= \left(\frac{gz}{k} - \frac{1}{2} \frac{g^2 z^2}{k'n} + \dots \right). \end{aligned}$$

Now the larger we make n , the more nearly our hypothetical case approaches to the continuous variation of the actual density of the air, and by taking n very large, we obtain the approximate expression,

$$z = \frac{k}{g} \log \frac{h}{h'},$$


observing that h' is less than h , and that the temperature and the force of gravity are supposed constant throughout the height z .

The Siphon.

90. The action of a siphon is an important practical illustration of atmospheric pressure.

It is simply a bent tube ABC , which is open at both ends. When filled with water, the ends are closed and the siphon is then inverted, and one end C placed in water, the other end A being below the surface of the water.

If the end *C* be opened, it is clear that the pressure at *A* is greater than the pressure at *Q*, which is equal to the pressure at *P*, and therefore to the atmospheric pressure.



Hence, if the end *A* be unclosed, the water at *A* will begin to flow out, and by so doing diminish the pressure in the tube, and tend to form a vacuum in the upper portion of the tube. But if the height of *B* above the surface of the water be less than the height *h* of the water-barometer, the atmospheric pressure will force the water up the tube, and maintain a continuous flow through the end *A*, until either the surface has fallen below *C*, or, if the siphon be long enough, until it has descended so far that its depth below *B* is greater than *h*.

91. *Methods of filling and graduating a Thermometer.*

To fill the Thermometer with mercury a paper funnel is fastened to the open end, and mercury poured into it; the bulb is then heated over a spirit-lamp, a portion of the air in the tube is thereby expelled, and if the bulb be cooled the mercury descends in the tube. This process is repeated until the air is completely expelled, and when the tube is quite full and the mercury overflowing, the upper end is hermetically sealed by means of a blow-pipe; during the subsequent cooling the mercury contracts and descends, leaving the vacuum at the top of the tube*.

The freezing and boiling points are now to be determined.

The freezing point is obtained by immersing the bulb and the lower portion of the tube in melting snow, and

* This so-called vacuum is filled with the vapour of mercury.

marking the tube outside at the end of the mercurial column.

The boiling point is obtained by immersing the bulb in the vapour of water boiling under a given atmospheric pressure, and marking the tube as before.

The temperature of steam depends on the atmospheric pressure, and it is therefore necessary to fix on some standard pressure, and to define the boiling point as the temperature of steam at that pressure. A barometric column of 30 inches at the level of the sea is the usual standard.

For the Centigrade Thermometer, the boiling point, 100° , is the temperature of steam when the height of the barometric column is 29.9218 inches at the level of the sea in latitude 45° .

For some time after boiling the height of the mercury at the freezing temperature is gradually increased, and it has been found that it takes 4 or 5 years for the zero to attain its permanent position after boiling.

92. *Use of the Mercurial Thermometer limited.*

Mercury freezes at a temperature of -40° C, and boils at a temperature of about 350° C; it is therefore necessary for very high or very low temperatures to employ different substances.


For very low temperatures spirit of wine is used, and this liquid is generally employed in the construction of *minimum* Thermometers.

High temperatures are compared by observing the expansion of bars of metal or other solid substances, and various instruments, called *pyrometers*, have been constructed for this purpose.

93. *The Differential Thermometer* is constructed in two different forms. In one form, of which the figure is a section, a horizontal tube branches upwards into two short vertical tubes ending in bulbs of equal size.

These bulbs contain air, and in the horizontal tube is a small portion of some coloured liquid, by which the air in one bulb is separated from the air in the other. The

quantities of air are equal, so that when the bulbs have the same temperature the bubble of liquid rests at the



middle of the tube: if however the temperatures be different, the liquid will rest in a position nearer to the bulb of lower temperature than to the other, since the air-pressure within it will be less than that in the other.

In the other form of the differential thermometer, the vertical portions, *A*, *B*, of the tube extend to a much greater height, and the liquid fills the whole of the horizontal portion of the tube, and also partly fills the vertical portion of the tube.

The principle of the construction is the same, and the difference consists in the graduation of the vertical portions, instead of the horizontal portion of the tube.

On account of their great sensibility these thermometers are extremely useful in detecting small differences of temperature.

In graduating the second of these instruments, allowance must be made for the weight of the liquid, which is contained in the vertical tubes.

94. Ex. 1. *The same quantities of atmospheric air are contained in two hollow spheres; the internal radii being r , r' and the temperatures t , t' respectively, compare the whole pressures on the surfaces.*

Taking ρ , ρ' as the densities, we have, since the masses are equal, and the volumes in the ratio of $r^3 : r'^3$,

$$\rho r^3 = \rho' r'^3.$$

If p, p' be the corresponding pressures,

$$p = k\rho(1+at), \quad p' = k\rho'(1+at'),$$

and the pressures on the surfaces are


$$4\pi r^2 p, \text{ and } 4\pi r'^2 p',$$

which are in the ratio

$$r^2 \rho (1+at) : r'^2 \rho' (1+at'),$$

$$\text{or } r' (1+at) : r (1+at').$$

Ex. 2. *A hollow cylinder, open at the top, is inverted, and partly immersed in water; it is required to find the height of the surface of the water within the cylinder.*



Take b for the length of the cylinder, and a for the length not immersed.

Let x be the depth of the surface within below the surface without, Π, Π' the pressures of the atmospheric air and of the compressed air in EC .

Then $\Pi' : \Pi :: b : a+x$, Art. 83,

and $\Pi' =$ pressure of the water at the level $C = \Pi + g\rho x$;

$$\therefore \frac{\Pi + g\rho x}{\Pi} = \frac{b}{a+x}.$$

If h be the height of the water-barometer, $\Pi = g\rho h$, and

$$\frac{h+x}{h} = \frac{b}{a+x},$$

$$\text{or } x^2 + (a+h)x = (b-a)h.$$

This equation gives two values for x , one positive and the other negative, the positive value being the one which belongs to the problem before us. The negative value is the result of another problem, the algebraical statement of which leads to the same quadratic equation.

Ex. 3. *A small quantity of air is left in the upper part of a barometer-tube; it is required to determine the effect on the height of the column.*

Let a be the length of the upper part of the tube which the air would occupy if its density were the same as that of the external air, and x the space it actually occupies, when the height of a true barometer is h .

If Π be the pressure of the external air, and Π' of the air in the space x ,

$$\frac{\Pi'}{\Pi} = \frac{a}{x}.$$

Let h' be the height of the faulty barometer, then

$$\Pi = g\sigma h, \text{ and } \Pi' + g\sigma h' = \Pi;$$

$$\therefore \frac{h-h'}{h} = \frac{a}{x}. \quad (1).$$

The column is therefore depressed $\frac{ah}{x}$ inches,

$$\text{or, since } \frac{h'}{h} = 1 - \frac{a}{x}, \quad \frac{ah'}{x-a} \text{ inches.}$$

Hence, if a be known, and h' and x be observed, the height of a true barometer can be inferred.

If a be unknown, it can be found from the equation (1) by taking simultaneous observations of h' , x , and the height h of a true barometer.

EXAMINATION ON CHAPTER V.

1. WHAT is the effect of heat on the elastic force of air or gas?

2. If Fahrenheit's Thermometer mark 40° , what are the corresponding marks of Reaumur's and the Centigrade?

3. Describe a method of shewing that air is a ponderable body.

4. When the mercurial barometer stands at 30 inches, what is the height of the barometer formed of a liquid of which the specific gravity is 5.6?

5. The air contained in a cubical vessel, the edge of which is one foot, is compressed into a cubical vessel of which the edge is one inch; compare the pressures on a side of each vessel.

6. State the relation between the pressure, density, and temperature of a gas.

The air in a spherical globe, one foot in diameter, is compressed into another globe, 6 inches in diameter, and the temperature is raised by t° ; compare the pressures of the air under the two conditions. Also compare the pressures on the surfaces of the globes.

7. Describe the siphon and its action. What would be the effect of making a small aperture at the highest point of a siphon?

8. Explain how the boiling point in a thermometer is defined.

9. If a barometer be held in a position not vertical, what would be the effect on the length of the column of mercury?

10. If the sum of the readings on Fahrenheit's and the centigrade thermometer be zero for the same temperature, find the reading of each thermometer.

11. At the top of a mountain the barometer stands at 25 inches; what would be the effect on the action of a siphon carried to the top?

12. A siphon is filled with mercury, and held with its legs pointing downwards, and the ends closed; what will be the effect of opening the ends, 1st, when they are, and 2ndly, when they are not, in the same horizontal plane?

13. A cylindrical vessel contains water; how will a change in the height of the barometer affect the pressures on the base and curved surfaces of the cylinder, and to what extent?

14. A block of wood weighs, in air, exactly the same as a block of iron; which is really the heavier?

15. Examine the effects of making a small aperture, 1st,

in the longer branch, 2ndly, in the shorter branch of the tube of a barometer?

16. Explain the uses, 1st, of the small hole which is made in the lid of a teapot, 2ndly, of a vent-peg.

17. Supposing the air half exhausted in a pair of Magdeburgh hemispheres, $1\frac{1}{2}$ ft. in diameter, find the force required to separate them, taking 15 lbs. as the atmospheric pressure on a square inch.

18. If a piece of glass float in the mercury within a barometer, will the mercury stand higher or lower in consequence?

19. Will any change in the action of a siphon be in any case coincident with a fall in the barometer?

20. A weight, suspended by a string from a fixed point, is partially immersed in water; will the tension of the string be increased or diminished as the barometer rises?

21. A bladder $\frac{1}{8}$ th filled with atmospheric air is placed under the receiver of an air-pump; the capacity of the receiver being twice that of the barrel. Shew that it will be fully distended before the completion of the 6th stroke.

NOTES ON CHAPTER V.

Thermometers were first constructed about the end of the sixteenth century, but the name of the inventor is not certainly known.


The various scales were formed in the early part of the 18th century; Fahrenheit's in 1714, at Dantzic; Reaumur's in 1731; and the Centigrade by Celsius, a Swede, somewhat later.

The Aneroid Barometer. This instrument was invented by Vidi, and is exceedingly useful in mountain ascents on account of its small size and weight. Its construction depends on the varying effect of the atmospheric pressure on a thin metallic plate closing an exhausted chamber. A small metallic chamber, cylindrical in form, about an inch in height, and 2 or 3 inches in diameter, and closed by an elastic metal plate, is exhausted; this is placed in a larger cylinder and the top of the elastic plate is connected by a system of levers with the hand of a graduated dial-face, so that any slight change of elevation or

depression at the centre of the metallic plate is magnified and rendered visible by the motion of the hand.

Bourdon's Metallic Barometer, invented in 1850, is another instrument of a similar kind*.

It consists of an elastic flattened tube, *ABC*, of metal, ex-



hausted of air, and bent very nearly into a circular form; the middle part *B* is fixed and the rest of the tube is free. The section of the tube is like an ellipse, *D*, and it is found that if the atmospheric pressure increase, the tube becomes more curved, and the ends *A*, *C* approach each other; and if it diminish, that the ends *A*, *C* separate. Hence if these ends be connected with the hand of a dial-face, the motion of the hand will mark the changes of atmospheric pressure.

If the tube *ABC*, instead of being a vacuum, be connected by a pipe with the boiler of a steam engine, or with any vessel containing air or gas, it becomes a very convenient *manometer*, (see Art. 114), and is in fact sometimes used for this purpose on the engines of locomotives.

The Siphon. The general use of the siphon is to transfer liquids from one vessel to another without moving either vessel. It is useful in many other operations, such as draining a flooded field; and lately large siphons, 140 feet in length and $3\frac{1}{2}$ feet in diameter, have been constructed for the purpose of draining the lands flooded by the inundation which occurred during the year 1862 on the eastern coast. These siphons were set working successfully. *The Times*, Oct. 1, 1862.

The Magdeburgh Hemispheres. A practical demonstration of the fact of atmospheric pressure was given by Otto von Guericke in 1654, who constructed this apparatus.

* The term Aneroid is sometimes applied to this instrument.

It consists of two hollow hemispheres of brass, fitting each other very accurately. A tube out of one of the hemispheres is screwed on the plate of an air-pump, and, when the two have been fitted together and the air exhausted, the stop-cock is turned, the apparatus removed from the air-pump, and a handle screwed on. Supposing the diameter of the hemispheres to be 3 or 4 inches, it will be found that a force of from 100 to 180 lbs. will be necessary to separate them. The inventor employed hemispheres of nearly a foot in diameter, and shewed that a strain of more than 1500 lbs. was required to force them asunder.



Taking the diameter as one foot, we can calculate the requisite force. The resultant pressure on one hemisphere is equal to the air-pressure on a circle one foot in diameter, that is, upon an area of 36π square inches. Making allowance for the fact that a perfect vacuum cannot be obtained, we may take 14 lbs. as approximately the pressure on a square inch, and the pressure is 504π lbs., or nearly 1583 lbs.

Weight of the Air. Galileo measured the weight of the air by filling a globe with compressed air, and then weighing the globe. He employed a syringe to force the air into the globe; and, in order to find the quantity of air, he placed the globe in an inverted glass receiver filled with water, then opened it, and observed the amount of water displaced.

Torricelli and Pascal. The experiment of Torricelli, described in Art. (73), was made in the year 1643, one year after the death of Galileo, who had remarked the fact that a pump would not raise water to a greater height than 32 or 33 feet, but was unable to account for it. It was reserved for his pupil and successor, Torricelli, to explain the real cause of the phenomenon, and his experiment was repeated and its consequences were extended by Pascal a few years later.

Torricelli shewed that the pressure of the air supports the column of mercury in a barometric tube; Pascal demonstrated that the weight of the air is the cause of the pressure. Amongst various experiments, Pascal had a water-barometer constructed, but his most valuable idea was a suggestion that the heights of a barometer, at the foot and at the top of a mountain, should be compared. This was effected by his friend Perier in 1648, who ascended the Puy de Dome in Auvergne, and ascertained the fact of a fall of nearly 4 inches in the barometer at the top

of the mountain. The observations were repeated in various ways, on the roofs of houses, and in cellars, and it was thus rendered clear that the weight of the air is the immediate cause of the existence of the barometric column.

The two treatises of Pascal, *De l'équilibre des liqueurs, et de la pesanteur de la masse de l'air*, contain the theory of the pressure of fluids, and give complete explanations of the actions of siphons and pumps, and of many common phenomena; the main object however of these treatises is to demonstrate the unphilosophical character of the old explanation that the abhorrence of nature to a vacuum accounted for the rise of water in a pump, and that this abhorrence did not exist beyond a rise of 32 feet.

It appears that Des Cartes was acquainted with the fact that air has weight, and indeed he made a suggestion that the reason why water will not rise beyond a certain height is *the weight of the water which counterbalances that of the air*.

Balloon Ascents. The fall of the barometer in balloon ascents is a means of determining the altitude attained.

In a balloon ascent by De Luc, the barometer at the greatest height stood at 12 inches; but in a late balloon ascent by Mr Glaisher, the column was seen to descend to less than 10 inches, implying a height of nearly six miles; and it is probable, as the observations were interrupted by the severity of the cold, and the rarity of the air, that an altitude of more than six miles was attained. *The Times*, Sept. 9, 1862.

EXAMPLES.

1. THE temperature of the air in an extensible spherical envelope is gradually raised t° , and the envelope is allowed to expand till its radius is n times its original length; compare the pressure of the air in the two cases.

2. A volume of air of any magnitude, free from the action of force, and of variable temperature, is at rest: if the temperatures at a series of points within it be in arithmetical progression, prove that the densities at these points are in harmonical progression.

3. A given weight of heavy elastic fluid of uniform temperature is confined in a smooth vertical cylinder by a piston of given weight; shew how to find the volume of the fluid.

4. A mass of air at a temperature t is contained in a cylinder which has an air-tight piston fitting into it, and it is found that the air exerts a pressure P on the piston: the air being suddenly compressed into $\frac{1}{n}$ th of its former volume, and the temperature changed to t' , find the pressure on the piston.

5. A piston moves freely in a closed air-tight cylinder, the axis of which is vertical. When the piston is in the middle of the cylinder, the air above and the air below are of the same density. Find the position of equilibrium of the piston.

6. A vertical closed cylinder is half filled with water, the other half being occupied by air of a given density and temperature; if the temperature be raised t° , find the increase of the whole pressure on the base, and on the curved surface of the cylinder.

7. Find the greatest height over which a liquid of density ρ can be carried by means of a siphon when the height of the barometer is h .

8. If h, h' be the heights of the surface of the mercury in the tube of a barometer above the surface of mercury in the cistern at two different times, compare the densities of the air at those times, the temperature being supposed unaltered.

9. A vertical cylinder, containing air, is closed by a piston, which is tied by an elastic string fastened to its central point, and also to the base of the cylinder. If when the piston is in equilibrium the string have its natural length, determine the effect on the length of the string of increasing the temperature of the air in the cylinder, by a given number of degrees.

10. *How does elastic force of a stretched string vary?*
If under an exhausted receiver a cylinder sinks to a depth equal to three-fourths of its axis; find the alteration in the depth of immersion when the air (specific gravity = .0013) is admitted.

× 11. A body is floating in a fluid; a hollow vessel is inverted over it and depressed: what effect will be produced in the position of the body, (1) with reference to the surface of the fluid within the vessel, (2) with reference to the surface of the fluid outside?

12. A pipe 15 feet long, closed at the upper extremity, is placed vertically in a tank of the same height; the tank is then filled with water; shew that, if the height of the water-barometer be 33 feet, 9 inches, the water will rise 3 feet, 9 inches in the pipe.

13. A vessel, in the form of a prism, whose base is a regular hexagon, is filled with air; prove that, if every rectangular face of the prism be capable of turning freely about its edges, and the prism be then compressed so that its base becomes an equilateral triangle, the pressure of the air within it will be increased in the ratio of 3 to 2.

14. A conical wine-glass is immersed, mouth downwards, in water; how far must it be depressed in order that the water within the glass may rise half way up it?

* 15. A jar contains water in which a hollow rigid envelope open at the bottom and partially filled with air just floats; the top of the jar is closed by an elastic membrane, and a small space between it and the water is filled with air; on pressing the membrane inwards the envelope sinks; explain this.

16. A barometer is held suspended in a vessel of water by a string attached to its upper end, so that a portion of the string is immersed; find the height of the mercury and the tension of the string. If more water be poured into the vessel, how will the tension of the string be affected?

17. A piston, the weight of which is equal to the atmospheric pressure on one of its ends, is placed in the middle of a hollow cylinder which it exactly fits, so as to leave a length a at each end filled with atmospheric air. The ends of the cylinder are then closed, and the cylinder is placed with its axis inclined at an angle α to the vertical; shew that the piston will rest at a distance $a\{(1 + \sec^2 \alpha)^{\frac{1}{2}} - \sec \alpha\}$ from its former position.

18. A cylinder, open at both ends, is partly immersed in water, its axis being vertical; the upper end is then closed, and the cylinder is raised until its lower end is very near the surface of the water outside; find the height to which the water rises inside.

19. Two barometers of the same length and transverse section each contain a small quantity of air; their readings at one time are h, k , and at another time h', k' ; compare the quantities of air in them.

20. Taking the figure of Art. 93, determine the position of the bubble when the temperatures of the bulbs are respectively t and t' .

21. In the second of the differential thermometers, calculate the difference of the altitudes of the liquid in the vertical tubes when the temperatures of the bulbs are t and t' .

CHAPTER VI.

The Diving Bell, Common Pump, Lifting Pump, Forcing Pump, Fire Engine, Bramah's Press, Air-Pumps, Barometer Gauge, Siphon Gauge, Condenser, Manometers, Barker's Mill, Piezometer, Hydraulic Ram, and Steam Engine.

The Diving Bell.

95. **T**HIS is a large bell-shaped vessel made of iron, open at the bottom, and containing seats for several persons. Its weight is greater than that of the water it would contain, and, when lowered by a chain into the water, the air within it is compressed, but will prevent the water from rising high in the bell, and the persons seated within are thus enabled to descend in safety to considerable depths.

When the surface of the water within the bell is at a depth of 33 feet below the outer surface the bell will be half filled with water, and the compression of the air would of course increase with the depth, but the difficulty arising from this compression is overcome by forcing fresh air from above through a flexible tube opening under the mouth of the bell. There are also contrivances for the expulsion of the air when rendered impure.


Tension of the Chain. This is equal to the weight of the bell diminished by the weight of water displaced by

the bell and the air within. It is therefore evident that unless fresh air is forced in from above the tension of the chain will increase as the bell descends.

96. *Supposing the bell cylindrical, and that no air is supplied from above, it is required to find the height to which the water rises in the bell.*

If the bell be partially immersed, we fall upon a case already considered, Ex. 2. Ch. v.

If the bell be wholly immersed, let b represent the length of



the cylinder, α the depth of its top, and x the length occupied by air.

$$\begin{aligned} \text{The pressure of the air within} &= \Pi \frac{b}{x} \\ &= \Pi + g\rho(\alpha + x); \end{aligned}$$

$$\text{and } \therefore \text{ if } \Pi = g\rho h,$$

$$hb = (h + \alpha)x + x^2,$$

and as before the positive value of x is the one required.

If A be the area of the top of the bell, and if we neglect its thickness, the volume of water displaced is Ax , and the tension of the chain

$$= \text{weight of bell} - g\rho Ax.$$

The Common Pump.

97. The Pump most commonly in use is a Suction-pump, of which the figure is a vertical section.

AB , BC are two cylinders having a common axis, M is a piston moveable over the space AB by means of a vertical rod, connected with a handle, D is a spout a little above A , and C the surface of the water in which the lower part of the pump is immersed: also in the piston, and at B , are valves opening upwards.


Action of the Pump. Suppose the piston at B and the pump filled with ordinary atmospheric air; raising the piston, the air in BC will open the valve B , and then, expanding as the piston rises, its pressure will be less than that of the atmosphere at C outside the pump; hence the atmospheric pressure on the surface of the water outside will force water up the tube BC , until the pressure at C is equal to the atmospheric pressure.

As the piston rises the water will rise in BC , the pressure of the air above M keeping the valve M closed. When the piston descends, the valve B closes, and the air in MB becoming compressed will open the valve M , and escape through it.

This process being repeated, the water will at length ascend through the valve B , and at the next descent of the piston will be forced through the valve M and be then lifted to the spout D , through which it will flow.

The height BC must be less than the height (h) of the water-barometer, or else the water will never rise to the valve B .

It is not essential to the construction that there should be two cylinders; a single cylinder, with a valve somewhere below the lowest point of the piston-range will be



sufficient, provided the lowest point of the range be less than 33 feet above the surface in the reservoir.

In each case the height above the water in the reservoir of the piston-range should be considerably less than 33 feet; otherwise the quantity of water lifted by the piston at each stroke will be small.

In the figure the tubes are represented as straight tubes; this is not necessary to the working of a pump, and the tube below the piston-range may be of any shape, and may enter the reservoir at any horizontal distance from the upper portion of the pump.

98. *Tension of the Piston-rod.* If the water in BC has risen to P when the piston is at M , the pressure Π' of the air in $MP =$ pressure of water at $P =$ pressure at $C - g\rho.PC$

$$= \Pi - g\rho.PC.$$

But if A be the area of the piston, the tension of the rod is the difference between the atmospheric pressure above and the pressure $\Pi'A$ below, i.e. $(\Pi - \Pi')A$, or $g\rho PC.A$.

If one inch be taken as the unit of length, and h be the height in inches of the water-barometer, $g\rho h = 15$ lbs. nearly, and the tension $= 15 \frac{PC.A}{h}$ lbs.

99. *To find the height through which the water rises during one stroke of the piston.*

Let P and Q be the surfaces of the water at the beginning and end of an upward stroke of the piston, that is, while the piston is raised from B to A .

The air which at the beginning of the stroke occupied the space BP occupies at the end of it the space AQ ; but the pressures are respectively, if $\Pi = g\rho h$,

$$g\rho(h - PC), \quad g\rho(h - QC).$$

Hence $h - PC : h - QC :: \text{vol. } AQ : \text{vol. } BP$.

If r , R be the radii of the cylinders, (Fig. Art. 97),

$$\text{vol. } AQ = \pi R^2 AB + \pi r^2 BQ = \pi R^2 AB + \pi r^2 (BC - QC),$$

$$\text{vol. } BP = \pi r^2 BP = \pi r^2 (BC - PC),$$

$$\frac{h - PC}{h - QC} = \frac{R^2 AB + r^2 (BC - QC)}{r^2 (BC - PC)},$$

and for any given value of PC this equation determines QC .

100. If the range of the piston be less than AB , as for instance AE , then EC must be less than h . Moreover, a limitation exists with regard to the position of E .

For, if P be the surface of the water when the piston M is at A , then as the piston descends, the valve B will close, but the valve M will not be opened until the pressure of the air in MB is greater than the atmospheric pressure.

When M is at A the pressure of the air $= g\rho (h - PC)$, and, unless the valve is opened before M arrives at E , the pressure of the air in $EB = g\rho (h - PC) \frac{AB}{EB}$, which must be greater than $g\rho h$, and therefore $h \cdot AE$ must be greater than $AB \cdot PC$. Hence, to ensure the opening of the valve while the surface is below B , we must have

$$h \cdot AE > AB \cdot BC;$$

i.e. AE must be at least the same fraction of AB that BC is of h .

This condition, although in all cases necessary, may not be sufficient.


For, suppose that when M is at A , the surface of the water is at Q , in which case the pressure of the air in $AQ = g\rho (h - QC)$.

When the piston descends to E , the pressure in EQ

$$= g\rho (h - QC) \frac{AQ}{EQ},$$

which must be greater than $g\rho h$,

$$\text{and } \therefore h \cdot AE > AQ \cdot QC.$$



The greatest value of $AQ \cdot QC$ is $\frac{1}{4} AC^2$, and \therefore we must have

$$h \cdot AE > \frac{1}{4} AC^2.$$

Since $\frac{1}{4} AC^2 > AB \cdot BC$, unless B is the middle point of AC , it follows that this latter condition includes the preceding, which is therefore in general insufficient.

These conditions must be also satisfied in the case of the pump with a single cylinder.


101. *Tension of the rod when the pump is in full action.*

In the figure of the previous Article, let $CD = h$; then it will be seen that, at each stroke, the volume DE of water is lifted, and therefore the tension of the rod when the piston is ascending will be $gpA(h + ED)$ until the water begins to flow through the spout.

If A be on a level with the spout, all the water lifted will be discharged, and, as the piston descends, the tension of the rod will be $gpAh$.

The Lifting Pump.

102. By means of this instrument, water can be lifted to any height. It consists of two cylinders, in the upper of which a piston M is moveable; the piston-rod works through an air-tight collar, and a valve opens outwards at D leading into a vertical tube. When the piston ascends, lifting water, the valve D opens and water ascends in the tube; when the piston descends the valve D closes, and every successive stroke increases the quantity of water in the tube. The only limitation to the height to which water can be lifted is that which depends on the strength of the instrument, and the power by which the piston is raised.




Tension of the rod. If $CK=h$ the piston lifts the volume BK at each stroke, and, as the air is expelled before the machine is in full action, the tension $=gpA \cdot KB$, until the water is lifted to the valve D . The power applied to the piston-rod must be then increased until the pressure of the water opens the valve D , that is, until the pressure $=gp(h+FD)$, F being the surface of the water in the tube. The water will then be forced up the tube, the tension of the rod increasing as the surface F ascends.


The Forcing Pump.

103. In this pump the piston M is solid, and ranges over the space AE . At B and D are valves opening upwards, DF being a tube leading out of AB .

When this pump is first set in action, it works as a common pump, the air at each descent of the piston being driven through D , and the water rising in BC . When however the water has risen through B , the piston, descending, forces it through D , and when the piston ascends, the valve D closes and more water rises through B . The next descent forces more water through D , and it is obvious that water can be thus forced upwards to any height consistent with the strength of the instrument.




The stream which flows from the top of the tube will be intermittent, but a continuous stream can be obtained by employing a strong air-vessel DL , out of which the vertical tube passes upwards. The air in the upper part of the vessel is condensed, and exerts a varying, but continuous pressure on the surface of the water within the vessel, and if the size of the vessel be suitable to that of the pump, and to the rate of working it, the air pressure will not have lost its force before a new compression is applied to



it, and thus a continuous, although varying, vertical flow will be maintained.

The Fire Engine.

104. The Fire Engine is only a modification of the Forcing-pump with an air-vessel, as just described.



Two cylinders are connected with the air-vessel, and the pistons are worked by means of a lever GEG' , so that while one ascends the other descends. The vertical tube out of the air-vessel has a flexible tube of leather attached to it, by means of which the stream can be thrown in any direction.


Bramah's Press.

105. This instrument is a practical application of the principle of the transmission of fluid pressures.

In the figure, which represents a vertical section of the instrument, A and C are two solid cylinders working in air-tight collars; EB and FD are strong hollow cylinders connected by a pipe BD ; at B is a valve opening inwards, and at D a valve opening upwards, a pipe from D communicating with a reservoir of water. M is a moveable platform, on which the substance to be pressed is placed, and N is the top of a strong frame; HKL is the lever working the cylinder C , H being the fulcrum, and L the handle.

Action of the Press. Suppose the spaces EB , FD filled with water, and C in its lowest position; on raising C , the atmospheric pressure forces water from the reservoir into FD , and when C is afterwards forced down, the valve D closes, the valve B is opened, a portion of the water in FD is driven into EB , and the cylinder A is

then made to ascend. A continued repetition of this process will produce any required compression of the substance between *M* and *N*.



At *G* there is a plug which can be unscrewed when the compression is completed.

The Force produced. If *P* be the power applied at the handle *L*, the force on *C* downwards is $P \frac{HL}{HK}$. Let *r*, *R* be the radii of the cylinders *C* and *A*, and *p* the pressure of the water,

then
$$\pi r^2 p = P \frac{HL}{HK},$$

and the pressure on *A* = $\pi R^2 p = P \frac{HL}{HK} \cdot \frac{R^2}{r^2}.$


It is obvious that by increasing the ratio of *R* to *r*, any amount of pressure may be produced.

We have taken for granted in describing the action of the press that the cylinders at first were full of water. If this is not the case the water will be pumped up from the reservoir by the action of the cylinder *C*, and whatever air there may be within will be compressed until its pressure is the same as that of the water.

Presses of this kind were employed in lifting into its place the Britannia Bridge over the Menai Straits.

106. The portion *C* of the instrument is sometimes called a *Plunger Pole Pump*, and an important part of the machine is the construction of the water-tight collars at *E* and *F*, as without these water under great pressure would force its way between the pole and the hollow cylinder in which it works.


A circular aperture *DE* is made in the side of the cylinder, and a piece of leather is doubled over a metal ring within it. The figure is a vertical section of the cylinder and collar, and it will be seen that the water pressing on the under side of the leather keeps it in close contact with the side of the cylinder, and the greater the pressure the closer the contact, so that no escape of water can possibly take place, unless the leather be torn.



Hawksbee's Air-Pump.

107. Two cylinders, *AB*, *A'B'*, are connected by pipes leading from *B* and *B'* through *C* with a receiver. Pistons *MM'* are worked in the cylinders by means of a toothed wheel, and at *B*, *B'* and in the pistons are valves opening upwards.

Suppose *M* at its highest and *M'* at its lowest position, and turn the wheel so that *M* descends and *M'* ascends; the valve *B* closes and the air in *MB* being compressed flows through the valve *M*, while the valve *M'* closes, and air from the receiver flows through *B'* into *M'B'*.



When the wheel is turned and M' descends, the valve B' closes and the air in $M'B'$ flows through M' , while the valve M closes and air from the receiver flows through B . At every stroke of the piston a portion of the air in the receiver is withdrawn, and it is evident that a degree of exhaustion may be thus obtained, limited only by the weight of the valves which must be lifted by the pressure of the air beneath.

Let A be the volume of the receiver, and B of either cylinder; ρ the density of atmospheric air and $\rho_1, \rho_2, \dots, \rho_n$ the densities in the receiver after 1, 2, ... n descents of the pistons.

After the first stroke the air which occupied the space A will occupy the space $A + B$, and therefore

$$\rho_1(A + B) = \rho A,$$

$$\text{similarly } \rho_2(A + B) = \rho_1 A;$$

$$\therefore \rho_2(A + B)^2 = \rho A^2,$$

and after n strokes

$$\rho_n(A + B)^n = \rho A^n.$$

Hence if Π_n be the pressure of the air in the receiver after n strokes, and Π of the atmospheric air,


$$\frac{\Pi_n}{\Pi} = \frac{\rho_n}{\rho} = \left(\frac{A}{A + B} \right)^n.$$

In working the instrument, the force required is that which will overcome the friction, together with the difference of the pressures on the under surfaces of the pistons, the pressures on their upper surfaces being the same.

It will be seen that a perfect vacuum cannot be obtained by this instrument, but, since the density decreases in geometric progression as the number of strokes increases, a very large proportion of the air can be withdrawn if the instrument be constructed with sufficient care.

Smeaton's Air-Pump.

108. This instrument consists of a cylinder AB in which a piston is worked by a rod passing through an air-tight collar at the top; a tube from B leads into a glass receiver C , and at A and B , and in the piston there are valves opening upwards.



Supposing the receiver and cylinder to be filled with atmospheric air, and the piston at B ; raising the piston, the air in AM is compressed, opens the valve A , and flows out through it, while at the same time a portion of the air in C flows through the valve B , so that when the piston arrives at A , the air which at first occupied C now fills both the receiver and the cylinder. When the piston descends, the valves B and A close and the valve M opens; the air in AB passes above the piston, and as the piston rises is forced through A , which is opened as soon as the pressure in M becomes greater than the atmospheric pressure. Thus at every stroke a portion of the air in the receiver is forced out through A .


If ρ be the density of atmospheric air, ρ_n the density in the receiver after n strokes of the piston, and A, B the volumes of the receiver and cylinder respectively, then, as in the previous article,

$$\rho_n(A + B)^n = \rho A^n,$$

observing that the volume of the connecting tube is neglected.

An advantage of this instrument is that, the upper end of the cylinder being closed, when the piston descends, the valve *A* is closed by the external pressure, and the valve *M* is then opened easily by the air beneath. Moreover, the labour of working is diminished by the removal, during the greater part of the stroke, of the atmospheric pressure on *M*, which is only exerted while the valve *A* is open during the latter part of the ascent of the piston.

A greater degree of exhaustion may be obtained by making the *B* aperture in the side of the cylinder without a valve, and working the piston, a solid one with or without a valve, below the aperture *B*. The limitation arising from the weight of the valve at *B* is thus removed, and the only limitations left are those which arise from the weight of the valve at *A*, and the exact fitting of the piston and receiver.




The Barometer Gauge.

109. The density of the air in the receiver of an air-pump at any moment is shewn by this instrument.

It is simply a barometric tube, the upper end of which communicates with the receiver, while the lower end is immersed in a cup of mercury, so that, as the pressure in the receiver diminishes, the mercury will rise in the tube.

If x be the altitude, PQ , of the mercury in the gauge, and h the height of the barometer, the pressure of the air in the receiver $= g\sigma h - g\sigma x$, if σ be the density of mercury.



Hence the density in the receiver is to the density of atmospheric air $:: h - x : h$.

It is important to use this gauge for experiments requiring strict accuracy, but for less important experiments a siphon-gauge may be used.

The Siphon Gauge.

110. This is a glass tube *ABCD*, the end *D* of which can be screwed on a pipe communicating with the receiver.

The end *A* is closed and the portion *AB* completely filled with mercury, which also fills a small part *BP* of *BC*.



If *AP* be not more than 28 inches in length, the tube *AB* will at first remain completely filled, but as the exhaustion proceeds, the mercury will sink in *AB* and rise in *BC*, and if at any time *x* be the difference of the heights in *AB* and *BC*, $g\sigma x$ will be the pressure in the receiver, and the density will


therefore be $\rho \frac{x}{h}$.

The Condenser.

111. This instrument is employed in the compression of air.

A hollow cylinder *AB* has one end screwed into the neck of a strong receiver *C*; at *B* is a valve opening inwards, and a piston *M* also has a valve opening inwards.

Suppose the cylinder and receiver filled with atmospheric air and the piston to be at *A*; forcing the piston down, the air in *MB* is compressed, and, opening the valve *B*, is forced into the receiver. When the piston is drawn back, the valve *B* is closed by the air in the receiver, and the valve *M* is opened by the outer air which flows in and fills the cylinder: this air is forced into the receiver at the next stroke, and at every succeeding stroke the same quantity of air is added to the receiver.




After n strokes, the volume of air of density ρ , forced into the receiver, is $A+nB$, A being the volume of the receiver and B of the cylinder; hence, if ρ_n be its density,

$$\rho_n A = \rho (A + nB),$$

$$\text{or } \frac{\rho_n}{\rho} = 1 + n \frac{B}{A}.$$

Gauge of a Condenser. A glass tube AB , closed at the end B , and connected with the condenser at the end




A contains atmospheric air in the portion BC , which is separated from the air in the condenser by a drop of mercury at C . As the condensation proceeds, the drop of mercury is forced towards B , until the density in BC is the same as the density in the condenser. Thus when the mercury is at D the density $= \rho \frac{BC}{BD}$.

Manometer.

112. The term manometer is applied to any instrument for measuring the pressure of condensed air or gas of any kind, when its elastic force is greater than that of the atmosphere. The gauge of a condenser, for instance, is a manometer. The term however is sometimes applied to any instrument, such as the barometer-gauge, for measuring the elastic force of air or gas under any circumstances.

The annexed figure represents a manometer, the principle of which is nearly the same as that of the gauge of a condenser.

AB is a vertical glass tube, closed at the end A and containing dry air in the part AP ; the tube ends in a strong bulb B containing mercury, and from this bulb a tube BC pro-



ceeds, leading to the vessel which contains the condensed air or gas. When the air in the tube C is ordinary atmospheric air at a given pressure, the mercury stands at the same level CC' in both tubes, but when the tube BC is connected with air or gas at a higher pressure the mercury rises in $C'A$, compressing the air above it, until the pressure in PA is equal to the pressure in EC diminished by the pressure due to the column PE' of mercury.

113. To find the relation between the pressure to be measured and the height of the mercury.

Let Π' represent the pressure in EC , and Π'' the pressure in PA ;

$$\text{then } \Pi'' = \Pi \cdot \frac{AC'}{AP}, \text{ and } \Pi'' + g\sigma PE' = \Pi';$$

$$\therefore \Pi' = g\sigma \cdot PE' + \Pi \frac{AC'}{AP}.$$

Let k, K be the sectional areas of the tubes AC', CE ;

$$\therefore \text{if } PC' = x, \quad CE = \frac{xk}{K},$$

$$\text{and } \Pi' = g\sigma x \left(1 + \frac{k}{K}\right) + \Pi \frac{a}{a-x} \text{ where } a = AC',$$

or if $\Pi = g\sigma h, \quad \Pi' = g\sigma h',$

$$\frac{h'}{h} = \frac{x}{h} \left(1 + \frac{k}{K}\right) + \frac{a}{a-x}.$$

This equation gives the ratio of the pressure required to the atmospheric pressure.

The graduation of the instrument depends on the solution of the equation; thus, making $h' = 2h, 3h, \&c.$, the successive proper values of x mark the altitudes for pressures of 2, 3, ... atmospheres.

114. The *Siphon Manometer* is a long glass tube ABC , open at the end A , and communicating at the end C

with the gas or vapour, the pressure of which is to be measured.

The tube contains mercury, and the height of the mercury in AB above its equilibrium level measures the excess of the pressure in the part BC of the tube above the atmospheric pressure.


Then if the mercury ascend to P in AB , and descend to E in CB , CC' being the original level, $CE = C'P$, and therefore, if $C'P = x$, and $\Pi' =$ pressure in CB ,

$$\begin{aligned}\Pi' &= \Pi + g\sigma 2x, \\ \text{or } \Pi' - \Pi &\propto x.\end{aligned}$$

A graduated scale is attached to the tube AB , and, from the equation above, it is seen that the length of $C'P$ corresponding to a pressure of n atmospheres is $\frac{n-1}{2}h$, if h be the height of the ba-

rometer. Hence by giving successive integral or fractional values to n , the graduation of the scale can be effected.


The manometers we have now described are constructed on purely hydrostatic principles, but there are others, depending on different mechanical principles, and a very useful one, from its portability, is *Bourdon's Metallic Manometer*, which has the additional advantage of not being fragile. The construction of this instrument is briefly explained in the notes appended to Chapter v.



Barker's Mill.

115. ACB is a tube, capable of revolving about its axis which is vertical, and having two or more horizontal tubes BE , BD connected with it. C is a cup through which water can be poured down the tube, and at D and E , in the sides of BD and BE , orifices are made which open in opposite directions. Suppose a stream of water to flow into C and through the tubes; as the water flows

through *BD* the pressures on the sides balance each other except at *D*, at which part of the tube there is an uncompensated pressure on the side opposite the orifice, the effect of which is to turn the tube *CD* round. The




same effect is produced by the water issuing at *E*, and a continued rotation of the instrument is thus produced. By means of a toothed wheel at *A* the instrument may be employed in communicating and maintaining motion in other machines.

The Piezometer.

116. This is an instrument for measuring the compressibility of liquids.

A thermometer tube *CD*, open at the end *C*, is enclosed in a strong glass vessel, which also contains a condenser-gauge *EF*. (See Art. 111.)


The liquid to be examined is poured into *CD*, and a drop of mercury is then introduced into *CD* so as to isolate the liquid, and the vessel is filled with water and closed by a piston. The piston *A* is moveable in the neck of the vessel, and, by means of a screw *B*, any required pressure can be produced. The gauge *EF* measures the pressures, and the compression of the liquid is obtained by observing the space through which the drop of mercury *P* is forced.



The area of a section of *CD* and the volume of the bulb are found by weighing the quantities of mercury contained by the bulb and a portion of the tube.

The Hydraulic Ram.

117. The fall of water from a small height produces a momentum which by means of the Hydraulic Ram* is utilized and made to produce the ascent of a column of water to a much greater height.



The figure is a vertical section of the machine, *AB* being the descending and *FG* the ascending column of water, which is supplied from a reservoir at *A*. *E* is an air-vessel with a valve at *C*, opening upwards; at *D* is a valve opening downwards, and *H* is a small auxiliary air-vessel with a valve *K* opening inwards.

The action of the Machine. The valve *D* will at first be open in its lowest position, and if water descend from *A*, a portion will flow through *D*, but the action on the valve will soon close it, and the sudden check thus produced increases the pressure; the valve *C* is lifted and water flows into the vessel *E*, and condenses the air within; the reaction of the air thus condensed forces water up the tube *FG*.

During this process the pressure of the water in the large tube diminishes, and the valves *C* and *D* both fall; the fall of the latter produces a rush of water through the


* Invented by Montgolfier.

opening D , followed by an increased flow down AB , the result of which is again the closing of D , and a repetition of the process just described, the water ascending higher in FG , and finally flowing through G .

The action of the machine is assisted by the air-vessel H in two ways, first, by the reaction of the air in H which is compressed by the descending water, and secondly by the valve K which affords supplies of fresh air. When the water rises through C , the air in H suddenly expands, and its pressure becoming less than that of the outer air, the valve K opens, and a supply flows in, which compensates for the loss of the air absorbed by the water and taken up the column FG , or wasted through D . About a third of the water employed is wasted, but the machine once set in motion will continue in action for a long time provided the supply in the reservoir be maintained.

The Atmospheric Steam Engine.

118. This instrument, constructed by Newcomen soon after the year 1700, was the first in which the oscillation of a beam was maintained by the elastic force of steam.



A solid beam EGF , which is moveable about G , has its ends arched; to these ends chains are attached which are connected with the rod of a piston in a cylinder AB , and with a rod supporting a weight P , this weight being less than the atmospheric pressure on the piston. C is a

pipe connected with a boiler, *B* a pipe opening by a stop-cock, and *D* is a pipe connected with a cistern of cold water.

This engine was first used for working the pumps of mines, and a rod *Q* attached to *P* is connected with the piston-rod of a pump.

The stop-cocks at *C* and *D* are connected with the beam, so that when *M* is at *A*, *C* is closed, and *D* opens, and when *M* is at *B*, *C* opens and *D* is closed. The stop-cock at *B* is made to open when *M* descends to *B*, and to close immediately after.

Action of the Engine. The pressure of the steam in the boiler is a little greater than that of the atmosphere, and when *M* is at *B*, *C* is open, and steam rushes into *MB*; hence the weight *P* will cause the piston to ascend. When *M* reaches *A*, *C* is closed, *D* is opened, and a jet of cold water is thrown in, condensing the steam, and thereby producing very nearly a vacuum below *M*. The pressure of the air on the piston being greater than the weight *P* forces the piston down, and when it has descended, *C* again opens, and an oscillation of the piston is thus maintained.

As *B* opens when *M* descends to the lowest point of its range the water flows out before the ascent.

In the actual engine constructed by Newcomen the stop-cocks were turned by hand, but an attendant, left to work them, invented the machinery by which the engine became self-acting.


The Single-acting Steam Engine.

119. In the atmospheric engine, the cooling of the cylinder at each stroke of the piston causes a great loss of power, for the steam on first entering the cylinder is partially condensed, and its elastic force is therefore diminished. One of Watt's first improvements was to produce the condensation in a separate vessel. The tube *D* was made to communicate with a vessel containing cold water, the space above the water being a vacuum. This vacuum could be produced by filling the vessel with steam and then

condensing it by cooling the vessel. When the piston is at *A*, the stop-cock opens and the steam rushes into the vacuum, and is therefore condensed by the cold water. A pump from the condensing vessel was connected with the beam, so that the overplus of water arising from the condensed steam would be drawn off as soon as formed. These two changes in the atmospheric engine constitute the single-acting engine, but the additional change of making the steam drive the piston downwards as well as upwards, leads to the double-acting engine, the type of most of the steam engines now in actual use.

Watt's Double-acting Steam Engine.

120. The cylinder *AB*, in which the piston works, is closed at both ends, the piston ranging from *a* to *b*. The end of the piston-rod is connected by means of a jointed



parallelogram with the end E of the beam EGF , and the end F of the beam is attached to the crank of the fly-wheel. At C and D there are stop-cocks which are connected with the fly-wheel, so that when M arrives at a , the steam flows from the boiler through C into AM , and when M arrives at b , the steam flows through D into BM . In each case the steam is shut off when M has passed over about one-third of its range.

K , the condenser, is surrounded with cold water, and L is a pump connected with it; a tube from K , not drawn in the figure, is connected with C and D so that when steam from the boiler flows into AM , the steam from MB flows into K , and when steam from the boiler flows into MB , the steam from AM escapes into K .

Supposing M to be at a , steam enters AM from the boiler and forces the piston down, its expansive force being sufficient to complete the piston-range after it is cut off; on arriving at b , the steam in AM escapes into K and is condensed, and fresh steam from the boiler enters MB , drives the piston upwards, and then escapes into K and is condensed. The continued accumulation of water in K is prevented by the pump L , by which it is drawn off at every stroke.

The use of the fly-wheel is to maintain a continuous motion, and prevent the irregularity which would arise from the intermittent action of the piston.

Parallel motion. The parallelogram $EQRS$ represents a system of jointed rods, invented by Watt for the purpose of making the end Q of the piston-rod move very nearly in a vertical line. The point R is connected with a fixed centre at P , and, by a proper adjustment of the lengths of the rods, it is found that the point Q deviates very slightly from the vertical during its motion.

A full account of the various contrivances for parallel motion will be found in Professor Willis's *Mechanism*.

The High-Pressure Engine.

121. In the double-acting engine the pressure of the steam need not be greater than the atmospheric pressure.

In the high-pressure engine it is many times greater, and the steam instead of being condensed is let off into the open air at each stroke. The condenser and air-pump are thus rendered unnecessary, and the engine simplified. The engines of locomotives on railways are high-pressure engines.

These descriptions give the main principles on which the constructions of steam engines depend, but for the various forms in which these principles are developed, and the innumerable details of the mechanism connected with them, the reader must consult special treatises on the subject, such as Dr Lardner's in Weale's series, Bourne's works on the Steam engine, or the excellent article in the *Encyclopædia Britannica*.

EXAMINATION UPON CHAPTER VI.

1. A DIVING-BELL is lowered until the surface of the water within is 66 feet below the outer surface; state approximately how much the air is compressed.

2. If a small hole be made in the top of a Diving-bell, will the water flow in, or the air flow out?

3. Describe the action of a common Pump.

To what height could mercury be raised by a pump?

4. Distinguish between a Lifting Pump and a Forcing Pump, and state the principle of the construction of a Fire-engine.

5. In a Bramah's Press, HK is one inch, HL is 4 inches, the diameter of A is 4 inches, and that of C is half an inch; find the force on A produced by a force of 2 lbs. applied at L .

6. If the receiver be 4 times as large as the barrel of an air-pump, find after how many strokes the density of the air is diminished one half.

7. State any limitations which exist to the degree of exhaustion producible by an air-pump.

8. Describe the Siphon-gauge, and its use.

9. What is a Manometer? Describe any such instrument.

What must be the height of a Siphon Manometer that it may mark a pressure of 60 lbs. on a square inch?

10. Describe the difference between the Atmospheric Steam engine and Watt's Double-Acting Engine.


11. The diameter of the piston of a Lifting pump is 1 foot, the piston-range is $2\frac{1}{2}$ feet, and it makes 8 strokes per minute; find the weight of water discharged per minute, supposing that the highest level of the piston-range is less than 33 feet above the surface in the reservoir, and that 33 feet is the height of the water-barometer.

12. If, in working the same pump, the lower level of the piston-range be $31\frac{1}{2}$ feet above the surface in the reservoir, find the weight discharged per minute.

NOTES ON CHAPTER VI.

Archimedes' Screw.

This instrument, one of the earliest hydraulic machines on record, is employed for raising water, and depends for its action only on the weight and mobility of the particles of water.



Let $ABCD$ be a metal tube, bent into the form of a cork-screw, and then held so that its axis is inclined to the vertical, and let it be moveable about its axis. The axis is to be inclined so much to the vertical, that a stone, inserted at A , will fall to B , and after oscillating rest at B . In the figure the tube is drawn as if wound round a cylinder moveable about its axis.


If we turn the cylinder in direction of the arrows, B will ascend, and the portions of the tube from B to C will successively take the same positions as B relative to the axis of the cylinder; as they do so, the stone at B will fall into those positions, and thus be gradually passed along the tube. Instead of the stone, suppose water poured in at A ; the turning of the

instrument will gradually raise the water until it flows out at the upper end. If the end A be immersed in water, a continued stream will ascend and flow out above.

Tradition assigns to Archimedes the credit of the invention of this instrument, and it is certain that its use dates at least as far back as the time of Archimedes. It was employed in Egypt in draining the land after an inundation of the Nile.

The point B at which the stone will rest is not underneath the cylinder but on one side, the ascending side, and between the middle and the under part of the surface of the cylinder: this can be seen experimentally.

Speaking strictly, the point B lies between the lowest generating line of the cylinder, and the generating line which lies halfway between the highest and lowest generating lines.



The machine will not act unless the inclination of the axis of the cylinder to the vertical be greater than the pitch of the screw, i. e. the inclination of the thread of the screw to a circular section of the cylinder. If these inclinations be equal, the point B is on the side of the cylinder, on the middle generating line, and the descending tangent BT is directed downwards at all other points. To make this clear, take a cylinder, of which BF is a diameter; let the dotted line represent a portion of the thread of a screw, BT being the tangent at B , and turn the cylinder round BF , which is supposed to be horizontal, until BT is horizontal: the inclination of the axis to the vertical is then equal to the pitch of the screw.

Turn the cylinder further, and if the screw mark the direction of a tube, it is an Archimedes' screw, in a position to work freely in raising water.

The Piezometer.

In the *Annales de Chimie et de Physique*, Vol. xxxr., 1851, a full account is given, by M. Grassi, of experiments with this instrument on the compressibility of water and some other liquids, and also on the compressibility of glass: these experiments were a continuation of M. Regnault's on the compressibility of water and mercury.

The apparatus employed by M. Grassi is identical in principle with the piezometer of the text, but differs in details. In one particular point the difference is of practical importance;

instead of producing pressure by a screw, the pressure on the surface of the water is produced by means of condensed air. The advantages gained are that the pressure can be measured with greater precision, and that it can be adjusted more easily, and changed more gradually.

The following are Grassi's final conclusions with regard to water:

(1) The compressibility of distilled water, deprived of air, varies with the temperature, and diminishes as the temperature increases.

(2) For distilled water, the compression due to one atmosphere is the same whatever be the pressure, provided the temperature remain constant.

EXAMPLES.

1. If the receiver and the barrel of an air-pump are in the proportion of 4 to 1, find how much has been pumped out at the end of the fifth stroke.

* 2. How would the tension of the rope of a Diving-bell be affected by opening a bottle of soda-water in the bell?

3. If P be the weight of a Diving-bell, P' of a mass of water the bulk of which is equal to that of the material of the bell, and W of a mass of water the bulk of which is equal to that of the interior of the bell, prove that, supposing the bell to be too light to sink without force, it will be in a position of unstable equilibrium, if pushed down until the pressure of the enclosed air is to that of the atmosphere as W to $P - P'$.*

4. If a cylindrical Diving-bell, height 5 feet, be let down till the depth of its top is 55 feet, find the space occupied by air, the water-barometer standing at 33 feet.

Also find how much air must be forced in to expel the water completely.

* 5. After a very great number of strokes of the piston of an air-pump the mercury stands at 30 inches in the barometer-gauge, the capacity of the barrel being one-third that of the receiver, prove that after 3 strokes the height of the mercury is very nearly $12\frac{2}{3}$ inches. $17\frac{1}{3}$ in

6. A fine tube of glass, closed at the upper end, is inverted and its open end is immersed in a basin of mercury, within the

* See Appendix, Example 4.

receiver of a condenser; the length of the tube is 15 inches, and it is observed that after 3 descents of the piston the mercury has risen 5 inches; how far will it have risen after four descents*?

7. If a cylindrical diving-bell, whose capacity is V cubic feet, be sunk to such a depth that the water stands at $\frac{1}{m}$ th of its height, and be then lowered at the uniform rate of n feet per second, prove that the number of cubic feet of air at the atmospheric pressure which must be pumped in per second in order that the water may always remain at the same height will be $\left(1 - \frac{1}{m}\right) \frac{n}{h} V$, where h is the height of the water-barometer in feet.

8. The length of the lower pipe of a common pump above the surface of the water is 10 feet, and the area of the upper pipe is 4 times that of the lower: taking 33 feet as the height of the water-barometer, prove that if at the end of the first stroke the water just rise into the upper pipe, the length of the stroke must be very nearly 3 feet 7 inches.

9. If the receiver of an air-pump be over a fluid, on which a solid is floating, shew how to calculate the density of air in the receiver after one stroke of the piston.

10. A cylindrical diving-bell, of height a , is furnished with a barometer and lowered into a fluid: the heights of the mercury in the barometer before and after immersion being h and h' respectively, shew that the depth of the bottom of the bell below the surface of the fluid is equal to $\left(\frac{\sigma}{\rho} + \frac{a}{h'}\right)(h' - h)$, where σ is the specific gravity of mercury, and ρ that of the fluid.

11. A bent tube, the arms of which are vertical, and which is open at one end and closed at the other, is partially filled with mercury, the density of the air between the mercury and the closed end of the tube being initially equal to that of the external air. If this tube be placed within the receiver of an air-pump, investigate a formula for determining the difference of heights of the mercury, in the two arms of the tube, after n strokes of the piston.

* Take 30 inches as the height of the barometer.

12. If the highest level to which the piston of a common pump ranges be below the spout, find the greatest tension of the piston-rod.

13. The valve in the piston of an air-pump being of given size and weight, find at what point of the n^{th} descent the valve will be raised.

14. If h be the range of the piston of an air-pump, a its distance from the top of the barrel in its highest position, β its distance from the bottom in its lowest position, and ρ the density of the atmosphere; prove that the limiting density of the air in the receiver will be $\frac{a\beta}{(h+a)(h+\beta)} \rho$.

15. In the $(n+1)^{\text{th}}$ ascent of the piston of a Smeaton's air-pump, find the position of the piston when the highest valve (whose weight may be neglected) begins to open; and shew that then the tension of the piston rod : the pressure of the atmosphere on the piston :: $1 - \left(\frac{A}{A+B}\right)^n : 1 - \left(\frac{A}{A+B}\right)^n \frac{B}{A+B}$.

16. A cylindrical diving-bell of internal volume v , is filled with air at atmospheric pressure Π and absolute temperature t , and is lowered to a certain depth below the surface of water. Shew that if a small rise (x) in the temperature and increase (y) in the atmospheric pressure now take place, the apparent weight of the bell will be unaltered provided $\frac{x}{t} = \frac{yv'}{\Pi v}$, v' being the volume of the air in the bell.

CHAPTER VII.

Method of Determining Specific Gravities. Specific Gravities of Air and Water, the Hydrostatic Balance, the Common Hydrometer, Sikes's, Nicholson's, and Hare's Hydrometers, the Stereometer.

To compare the specific gravities of air and water.

122. TAKE a large flask, which can be completely closed by a stop-cock, and exhaust it by means of an air-pump.

Weigh the flask, and then permit the air to enter, and weigh the flask again. Finally find the weight of the flask when filled with water.

Let w be the weight of the exhausted flask, w' , w'' its weights when filled with air and water ;

$\therefore w' - w =$ weight of the air contained by the flask,
and $w'' - w =$ water.....

Hence $w' - w$ and $w'' - w$ being the weights of equal volumes of air and water,

specific gravity of water : that of air :: $w'' - w$: $w' - w$.

In the same manner the specific gravity of any gas can be compared with that of water.

The specific gravity of water at 20.5° is about 768 times that of air at 0° under the pressure of 29.9 inches of mercury at 0° .

To compare the specific gravities of two fluids by weighing the same volume of each.

Let w be the weight of a flask, w' its weight when filled with one fluid (A), and w'' its weight when filled with the other fluid (B).

Then

$w' - w =$ weight of the fluid A contained in the flask,

$w'' - w = \dots\dots\dots B \dots\dots\dots$;

\therefore specific gravity of A : that of B :: $w' - w$: $w'' - w$

If the flask be not exhausted when its weight is determined, then, for strict accuracy, w must be diminished by the weight of the air which the flask contains.

123. *To find the specific gravity of a solid broken into small fragments.*

Put the broken pieces in a flask, fill the flask with water and let its weight be then w'' ; let w be the weight of the flask when filled with water, and w' the weight of the solid in air.

Then

$w'' - w =$ weight of solid pieces - weight of the water they displace;

$= w' -$ weight of water displaced ;

therefore

$w' + w - w'' =$ weight of water displaced,

and $\frac{\text{specific gravity of solid}}{\text{that of water}} = \frac{w'}{w' + w - w''}$.

If we take account of the air displaced by the solid, its real weight is greater than w' by the weight of air displaced. This weight must therefore be added to w' .


The Hydrostatic Balance.

124. The hydrostatic balance is an ordinary balance, having one of the scale-pans smaller than the other, and at a less distance from the beam, so that weights immersed in water may be suspended from it.

The following cases are examples of its use.

(1) *To compare the specific gravities of a solid and a liquid.*

Let w be the weight of the solid in air.



Place the liquid in a vessel, as in the figure, and suspend the solid from the scale-pan.

Let w' be the weight of the solid in the liquid,

$\therefore w - w'$ is the weight lost by the solid, and is therefore the weight of the liquid displaced by the solid, Art. (39);

and w , $w - w'$ are the weights of equal volumes of the solid and liquid.

Hence,

specific gravity of solid : that of liquid $:: w : w - w'$.

If we take account of the air displaced by the solid, we must add to w the weight of the air it displaces, since its true weight is diminished by exactly the weight of air.

This remark applies also to the next two articles.

125. We have tacitly supposed the solid to be specifically heavier than the liquid. If it be lighter it must be attached to a heavy body of sufficient size and weight to make the two together sink in the liquid.

Let w = the weight of the solid in air,

x = the weight in air of the heavy body attached to it,

x' = the weight in the liquid of the heavy body,

w' = the weight in the liquid of the two together.

$w + x - w'$ = the weight of liquid displaced by the two together, since it is the weight lost.

$x - x'$ = weight of liquid displaced by the heavy body.

Hence

$w + x' - w'$ = weight of liquid displaced by the solid,

and therefore $\frac{\text{specific gravity of solid}}{\text{specific gravity of liquid}} = \frac{w}{w + x' - w'}$.

126. (2) *To compare the specific gravities of two liquids.*

Take a solid which is specifically heavier than either liquid, and let w be its weight in air.

Let w' = weight of solid in one liquid (A),

and w'' =the other liquid (B);

$\therefore w - w'$ = weight of liquid A displaced by the solid,

$w - w''$ = B;

\therefore specific gravity of A : that of B :: $w - w'$: $w - w''$.

The Common Hydrometer.

127. The common hydrometer consists of a straight stem ending in two hollow spheres B and C .

This hydrometer is usually made of glass, and the sphere C is loaded so that the instrument will float with the stem vertical.

When the hydrometer is immersed and allowed to float in a liquid, it displaces its own weight of the liquid, and by observing the positions of equilibrium in two liquids, the volumes displaced are inferred, and the specific gravities of the liquids can be compared.



Let κ be the area of a section of the stem, v the volume, and w the weight of the hydrometer.

Suppose that when floating in a liquid (A) the level D

of the stem is in the surface, and that in liquid (*B*) the level *E* is in the surface.

Then, if *s*, *s'* be the specific gravities of *A* and *B* respectively,

$$w = s (v - \kappa \cdot AD)$$

$$\text{and } w = s' (v - \kappa \cdot AE);$$

$$\therefore \frac{s}{s'} = \frac{v - \kappa \cdot AE}{v - \kappa \cdot AD}.$$

Sikes's Hydrometer.

128. This instrument differs from the common hydrometer in the shape of the stem, which is a flat bar and very thin, so that it is exceedingly sensitive. It is generally constructed of brass, and is accompanied by a series of small weights *F*, which can be slipped over the stem above *C* so as to rest on *C*.

The use of the weights is to compensate for the great sensitiveness of the instrument, which would without the weights render it applicable only to liquids of very nearly the same density.

Suppose the instrument floating in a liquid (*A*), with the level *D* of the stem in the surface, and that *w'* is the weight on *C*. In a liquid (*B*) let *E* be in the surface, and *w''* the weight at *C*.

Let *w* be the weight of the instrument, *v* its volume, κ the section of the stem, *v'*, *v''* the volumes of *w'*, *w''*, and *s'*, *s''* the specific gravities of the liquids.

Then $w + w' =$ weight of fluid *A* displaced,

$$v + v' - \kappa \cdot AD = \text{volume of } A \text{ displaced};$$

$$\therefore w + w' = s' (v + v' - \kappa \cdot AD).$$

Similarly $w + w'' = s'' (v + v'' - \kappa \cdot AE);$

and therefore $\frac{s'}{s''} = \frac{w + w'}{w + w''} \cdot \frac{v + v'' - \kappa \cdot AE}{v + v' - \kappa \cdot AD}.$



If the liquid (*B*) be the standard liquid, $s'' = 1$, and s' , the specific gravity of (*A*) is at once determined.

Nicholson's Hydrometer.

129. The two hydrometers just described are used for comparing the specific gravities of fluids; Nicholson's hydrometer can be also employed in comparing the specific gravities of a solid and a fluid.

It consists of a hollow vessel *B*, generally of brass, supporting a cup *A* by a very thin stem, which is often a steel wire, and having attached to it a heavy cup *C*: on the stem connecting *A* and *B* a well-defined mark *D* is made.

We proceed to explain the use of the instrument in the two cases.

(1) *To compare the specific gravities of two liquids.*

If w be the weight of the hydrometer, w' the weight which must be placed in *A* in order to sink the instrument to the point *D* in a liquid of specific gravity s' , and w'' the weight for a liquid of specific gravity s'' , the weights of the liquids displaced are respectively

$$w + w' \text{ and } w + w''.$$

Therefore, the volumes displaced being the same,

$$s' : s'' :: w + w' : w + w''.$$

(2) *To compare the specific gravities of a solid and a liquid.*

Let w be the weight which, placed in *A*, causes the instrument to sink to *D* in the liquid.

Place the solid in *A*, and let w' be the weight, placed in *A*, which sinks the instrument to *D*.

Then place the solid in *C*, and let the weight w'' , placed in *A*, sink the instrument to *D*.

Hence weight of solid = $w - w'$,

and its weight in the liquid = $w - w''$.



Hence the weight lost, which is the weight of the liquid displaced by the solid, = $w'' - w'$, and

\therefore spec. gravity of solid : that of liquid :: $w - w' : w'' - w'$.

If we take account of the air, we must, as before, add to $w - w'$ the weight of the air displaced by the solid.

Hare's Hydrometer.

130. This instrument is an application of the principle of the barometer; it consists of two vertical glass tubes leading out of a hollow vessel *A*, which can be connected with an air-pump.

B and *C* are two cups in which the lower ends of the tubes are immersed, and which contain the two fluids to be compared.

Let the air in *A* be partially withdrawn, so that its pressure is diminished from Π the atmospheric pressure to Π' .


Then if *D*, *E* be the surfaces of the liquids in the tubes, and *F*, *G* in the cups, and if ρ , ρ' be the specific gravities,

$$\Pi = \Pi' + g\rho DF, \text{ and } \Pi = \Pi' + g\rho' EG;$$

$$\therefore \rho DF = \rho' EG,$$

$$\text{and } \rho : \rho' = EG : DF.$$

There is no absolute necessity for an air-pump, as a partial vacuum may be obtained in several other ways.



The Stereometer.

131. The name stereometer* has been given to a modified form, by Professor Miller, of Say's instrument for measuring the volumes of small solids.

It consists of two glass tubes, *PQ*, *DB*, of equal dia-

* Prof. Miller, *Phil. Trans.* Part III. 1856.

meter, cemented into cylindrical cavities communicating with each other at their lower ends in a piece of iron *G*.

Two apertures lead out of *PQ* and *DB*, the one, *K*, stopped with a screw and the other, *L*, having a stop-cock.

The upper end of *PQ* opens into a cup *F*, the rim of which is ground plane, so that it can be closed and made airtight by a plate of glass *E*, smeared with lard. The tube *PQ* is graduated by lines traced on the glass, and measured downwards from a fixed point *P*.

The solid to be examined being placed in *F*, mercury is poured into *D*, till its surface rises to *P*, and the cup is then closed by the plate of glass.

The stop-cock *L* is then opened and the mercury allowed to escape till the difference of the heights of the mercury in the tubes is nearly equal to half the height of the mercury in the barometer. Let *M* and *C* mark the height in the tubes; and let *u* be the volume of the air in *F* before the solid was placed in it, *v* the volume of the solid, and *h* the height of the barometer.

The pressure at *C* = $g\rho h$,


and \therefore at *M* = $g\rho (h - MC)$.

Hence, if *K* be the section of either tube, since the volume varies inversely as the pressure,

$$\frac{u - v + K \cdot PM}{u - v} = \frac{h}{h - MC},$$

$$\text{and } v = u - \frac{h - MC}{MC} K \cdot PM.$$

The volume *u* can be found by a similar process, the cup *F* being empty, and *K* is found by weighing the mercury contained in a given length of the tube.



If the weight w of the solid v be determined, its specific gravity s is given by the relation $w = sv$.

132. The screw K is used in the process of finding K . To do this, the cup is taken off and the tube PQ closed; the tubes are then inverted, the screw K taken out, and mercury is poured in through a slender glass tube inserted in K ; this precaution is taken in order to prevent the formation of air-bubbles in PQ .

The end P is then opened and the mercury allowed to run into a glass jar in which it is weighed.

A cubic inch of mercury at 16° weighs nearly $3429\frac{1}{2}$ grains, and therefore if w be the weight of a column of mercury a inches in length,

$$w = 3429\frac{1}{2} \cdot Ka,$$

from which K is determined in square inches.

Say's instrument consisted of one tube PQ , the lower end being open, so that it could be immersed in a cylindrical vessel of mercury.

The instrument was invented for the purpose of determining the specific gravity of gunpowder: it can be employed in finding the specific gravities of powders or soluble substances, for which the methods which require immersion in water are inapplicable.

EXAMINATION UPON CHAPTER VII.

1. A SOLID, which is lighter than water, weighs 5lbs., and when the solid is attached to a piece of metal, the whole weighs 7 lbs. in water; the weight of the metal in water being 9 lbs., compare the specific gravities of the solid and of water.

2. A solid weighing 25lbs., weighs 16 lbs. in a liquid A , and 18 lbs. in a liquid B ; compare the specific gravities of A and B .

3. The whole volume of a hydrometer is 5 cubic inches, and its stem is one-eighth of an inch in diameter; the hydrometer floats in a liquid A with one inch of the stem above the surface, and in a liquid B with two inches above the surface; compare the specific gravities of A and B .

* 4. Describe the characteristic differences between Sikes's, Nicholson's, and the Common Hydrometer.

5. Describe the construction and use of the Stereometer.

6. What volume of cork, specific gravity .24, must be

attached to 6 lbs. of iron, specific gravity 7.6, in order that the whole may just float in water?

7. A body weighs 250 grains in a vacuum, 40 grains in water and 50 grains in spirit; find the specific gravities of the body and of the spirit.

8. A Sikes's Hydrometer floats in water with a given length (a) of its stem not immersed; it is then placed in a liquid (A), and when a weight w , volume v' , is placed on the lower end, it is found that the length of stem not immersed is the same as before; compare the specific gravity of A with that of water.

9. If a piece of metal weigh in vacuum 200 grains more than in water, and 160 grains more than in spirit, what is the specific gravity of the spirit?

10. A piece of metal whose weight in water is 15 ounces is attached to a piece of wood, which weighs 20 ounces in vacuum, and the weight of the two in water is 10 ounces; find the specific gravity of the wood.

EXAMPLES.

1. A PIECE of wood, which weighs 57 lbs. in vacuo, is attached to a bar of silver weighing 42 lbs., and the two together weigh 38 lbs. in water; find the specific gravity of the wood, that of water being 1, and that of silver 10.5.

2. The apparent weight of a sinker, weighed in water, is four times the weight in vacuum of a piece of a material, whose specific gravity is required; that of the sinker and the piece together is three times that weight. Shew that the specific gravity of the material is .5.

3. A hollow cubical metal box, the length of an edge of which is one inch and the thickness one-eighteenth of an inch, will just float in water, when a piece of cork, of which the volume is 4.34 cubic inches and the specific gravity .5, is attached to the bottom of it. Find the specific gravity of the metal.

4. A crystal of salt weighs 6.3 grains in air; when covered with wax, the specific gravity of which is .96, the whole weighs 8.22 grains in air and 3.02 in water; find the specific gravity of salt.

5. A Nicholson's Hydrometer weighs 6 oz., and it is requisite to place weights of 1 oz. and $1\frac{1}{2}$ oz. in the upper cup to

sink the instrument to the same point in two different liquids; compare the specific gravities of the liquids.

6. With the same hydrometer it is found that when a certain solid is placed in the upper cup a weight of $1\frac{1}{2}$ oz. must be placed in the upper cup to sink the instrument in a liquid to a given depth; and that, when the solid is placed in the lower cup, a weight of 3 oz. must be placed in the upper cup to sink the instrument to the same depth; compare the specific gravities of the solid and the liquid, the weight of the solid being 2 oz.

7. A ring consists of gold, a diamond, and two equal rubies, it weighs $44\frac{1}{4}$ grains, and in water $38\frac{3}{4}$ grains; when one ruby is taken out it weighs 2 grains less in water. Find the weight of the diamond, the specific gravity of gold being $16\frac{1}{2}$, of diamond $3\frac{1}{2}$, of ruby 3.

8. If the price of pure whiskey be 16s. per gallon, and its specific gravity be .75, what should be the price of a mixture of whiskey and water, which on gauging is found to be of specific gravity .8, the specific gravity of water being 1?

9. Supposing some light material, whose density is ρ , to be weighed by means of weights of density ρ' , the density of the atmosphere when the barometer stands at 30 inches being unity; shew that, if the mercury in the barometer fall one inch, the material will appear to be altered by $\frac{\rho' - \rho}{(\rho - 1)(30\rho' - 29)}$ of its former weight. Will it appear to weigh more or less?

10. A heavy bottle is filled with a fluid A and weighed in each of two other fluids B, C , the apparent weights being A_b, A_c ; it is then filled with the fluid B and weighed in C and A , the apparent weights being B_c, B_a ; lastly it is filled with fluid C and weighed in the fluids A and B , the apparent weights being C_a, C_b : shew that

$$A_b + B_c + C_a = A_c + B_a + C_b.$$

CHAPTER VIII.

Mixture of Gases, Vapours, Radiation, Conduction and Convection of Heat, Dew, Hoar Frost, Clouds and Rain, Sea and Land Breezes, Dew-Point, Hygrometers, Dilatation of Liquids, Maximum Density of Water, Congelation and Ebullition, Specific Heat.

Mixture of Gases.

133. **I**F two liquids are mixed together in a vessel, and if the vessel is left at rest, the two liquids, provided they do not act chemically on each other, will gradually separate and finally attain equilibrium with the heavier liquid lowest, and the lighter liquid superposed upon it. But if two gases are placed in communication with each other, even if the heavier gas be below the other, they will rapidly intermingle until the proportion of the two gases is the same throughout, and the greater the difference of density the more rapidly will the mixture be formed.

Take two different gases, having the same temperature and pressure, and contained in separate vessels; open a communication between the vessels, and it will be found that, unless a chemical action take place, the pressure of the mixture will be the same as before, provided the temperature be the same.

We can hence deduce the following proposition:

If two gases having the same temperature be mixed together in a vessel of volume V , and if the pressures of the gases when respectively contained in V , at the same temperature be p and p' , the pressure of the mixture will be $p+p'$.

Suppose the gases separate; change the volume of the gas, of which the pressure is p' , without change of temperature, until its pressure is p ; its volume will then be $\frac{p'}{p}V$, by Mariotte's Law.

Now mix the two gases without change of volume, so that the volume of the mixture is $V + \frac{p'}{p}V$, or $\frac{p+p'}{p}V$; by the preceding experimental fact, the pressure of the mixture will be still p .

Compress the mixture till its volume is V , and when the temperature is the same as before, the pressure, which varies inversely as the volume, will be $p+p'$.

This result is equally true of the mixture of any number of gases.

134. *Two volumes, V, V' , of different gases at the respective pressures p, p' , are mixed together in a vessel of volume U ; it is required to find the pressure.*

Change the volume of each gas to U ; their pressures will be respectively

$$\frac{V}{U}p, \quad \frac{V'}{U}p',$$

and therefore the pressure (ϖ) of the mixture will be

$$\frac{V}{U}p + \frac{V'}{U}p'.$$

Hence
$$\varpi U = pV + p'V'.$$

In Art. (133) we have assumed that Mariotte's law is true of a gas formed by the mixture of two gases; this can be shewn by direct experiment, but is in fact already proved in one case, by the original experiment with atmospheric air, which is itself composed of several different gases. Moreover, the results of the two preceding propositions are borne out by facts.

Vapours.

135. The term vapour is applied to those gaseous bodies, such as steam, which can be liquefied at ordinary

pressures and temperatures. There is no difference between the mechanical qualities, as distinguished from the chemical qualities, of vapours and gases, the laws already stated of gases being equally true of vapours within certain ranges of temperature. In fact, there is every reason to believe that all gases are the vapours of certain liquids, but those which are looked upon as permanent gases require the application of extreme cold and of very great pressure to reduce them to a liquid form.

Professor Faraday found that carbonic acid, at the temperature -11° , was liquefied by a pressure of 20 atmospheres*, but that, at the temperature 0° , a pressure of 36 atmospheres was required to produce condensation.

In 1877, M. Pictet succeeded in liquefying oxygen by subjecting it to a pressure of 300 atmospheres, and, at the end of the same year, M. Cailletet effected the liquefaction of nitrogen, atmospheric air, and hydrogen.

136. *Formation of vapour.* If water be introduced into a space containing dry air, vapour is immediately formed, and if the quantity of water be small, and the temperature high, the whole of the water will be rapidly converted into vapour, and in all cases the pressure of the air will be increased by the pressure due to the vapour thus formed.

An increase of temperature, or an enlargement of the space, increases the amount of vapour as long as the supply of water remains; but if the water be removed, an increase of temperature changes the pressure of the vapour in accordance with the general law which regulates the connection between pressure and temperature.

The formation of vapour does not in any way depend upon the presence of air or upon its density, the only effect which the air produces being a retardation of the time in which the vapour is formed. If water be introduced into a vacuum, it is instantaneously filled with vapour, but the quantity of vapour is the same as if the space had been originally filled with air.

Saturation. As long as the supply of water remains

* An atmosphere denotes the pressure due to a column of mercury 29.9 inches in height.

as a source from which vapour can be produced, any given space will be always *saturated* with vapour, that is, will contain the maximum quantity of vapour for any temperature; but if the temperature be lowered, a portion of the vapour will be immediately condensed, and become visible in the form of liquid.

The quantity of vapour by which any given space is saturated is proportional to the space for any given temperature; it follows that the pressure, or elastic force, of the vapour is independent of the space it saturates, and depends only on the temperature. No definite law has been discovered connecting the temperature and the elastic force of vapour, but tables have been formed and empirical formulæ constructed for certain ranges of temperature.

137. The laws of the mixture of gases are equally true of the mixture of vapours with each other, or of vapours with gases, provided no condensation take place; or, if any condensation should take place, provided a proper allowance be made for the loss of pressure incurred.

Thus all atmospheric air contains more or less aqueous vapour, and if p be the pressure of dry air and ϖ of the vapour in the atmosphere at any time, the actual atmospheric pressure is $p + \varpi$.

138. *Having given the pressures of a volume V of atmospheric air, and of the vapour it contains, to find the volume of the air without its vapour at the same pressure and temperature.*

Let Π be the atmospheric pressure, and ϖ that of the vapour.

Then $\Pi - \varpi$ is the pressure of the air alone when its volume is V ;

$$\text{Hence its volume at a pressure } \Pi = \frac{\Pi - \varpi}{\Pi} V.$$

139. *Having given the volume V of a dry gas at a given temperature under a pressure p , to find its volume under the same pressure, when saturated with vapour.*

Let ϖ be the pressure of the vapour.

Then the gas must be allowed to expand until its pressure is $p - \varpi$, the supply of vapour being kept up. The pressure of

the mixture is then p , and the volume will be $\frac{p}{p - \varpi} V$.

140. A gas contained in a closed vessel of volume V is in contact with water, and its pressure at the temperature t is p ; it is required to determine its pressure when V is changed to V' and t to t' .

Let ω and ω' be the pressures of the vapour at the temperatures t and t' respectively, and p' the required pressure.

Then $p - \omega$ and $p' - \omega'$ are the pressures of the gas alone, under the two sets of conditions stated.

Hence, if ρ, ρ' be the densities of the gas,

$$\begin{aligned} p - \omega &= \kappa\rho(1 + at), \\ p' - \omega' &= \kappa\rho'(1 + at'), \\ \text{also } \rho V &= \rho' V'; \\ \therefore \frac{p' - \omega'}{p - \omega} &= \frac{V}{V'} \cdot \frac{1 + at'}{1 + at}, \end{aligned}$$

whence p' is determined.

If σ, σ' be the densities of vapour under the two conditions,

$$\frac{\omega'}{\omega} = \frac{\sigma'(1 + at')}{\sigma(1 + at)},$$

and combining the two equations,

$$\begin{aligned} \frac{p' - \omega'}{p - \omega} \cdot \frac{\omega}{\omega'} &= \frac{V\sigma}{V'\sigma'}, \\ \text{or } \frac{V'\sigma'}{V\sigma} &= \frac{p\omega' - \omega\omega}{p'\omega - \omega\omega'}. \end{aligned}$$

If $p\omega' > p'\omega$, $V'\sigma'$ will exceed $V\sigma$; i. e. more vapour will have been absorbed by the gas, but if $p\omega' < p'\omega$, then $V'\sigma'$ will be less than $V\sigma$, and the gas must therefore, in changing its volume and temperature, have lost a portion of its vapour.

Radiation, Conduction, and Convection of Heat.

141. *Radiation.* All bodies give off heat from their surfaces by what is called radiation, and receive heat by radiation from other bodies. If two bodies at different temperatures are placed near each other, it is an experimental fact that the temperature of one will rise, and of the other diminish until they are both the same.

In a similar manner, if a body is placed in a confined space, the temperature of the body and of the boundary of the space will gradually approximate, the one increasing and the other decreasing till they are the same.

Difference of radiating power. Some bodies radiate heat more freely than others, and the difference appears to depend in great measure on the nature of the surfaces. Thus the leaves of trees and woollen substances radiate heat freely and rapidly, while the radiation from a polished metal surface is very slight.

Generally if the reflecting power of a surface be increased its radiating power is diminished.

142. *Conduction and convection.* There are two other modes of transference of heat from one body to another. Conduction is the term applied to the transference of heat by contact, heat being transmitted through the successive particles of a body, or from one body to another in contact with it. Convection is the actual transference of heat by the motion of fluids or other bodies from one position to another; the heat thus conveyed away from one body may be imparted by contact or radiation from the conveying body to any other.

Thus the handle of a poker, inserted in the fire, is heated by conduction, and in the process of warming rooms by hot air or hot-water pipes the heat is obtained by convection.

There are great differences in the conducting powers of different bodies; liquids generally are weak conductors, but metallic substances have large conducting powers.

The cold felt in placing the hand on a marble mantelpiece is an instance of conduction, the heat being transferred from the hand to the marble.

Woollen substances, glass, and wood, conduct heat very slowly, and this fact is practically taken advantage of in many ways. A heated body rolled up in a woollen cloth may be kept hot for a long time, and ice in a wooden pail, wrapped round with a cloth, will dissolve very slowly, even in a warm room.

Another instance of a body with very small conducting power is sand; heat is transferred through it so slowly that red-hot shot can be safely carried about in wooden barrows filled with sand.

One of the many useful applications of the non-conducting powers of certain substances is in the construc-

tion of *Fire-proof Safes*; a safe of this kind is simply an iron box enclosed within another somewhat larger, the space between being filled up with some non-conducting substance.

143. The explanations above given of the saturating density of vapour, and of the radiation of heat, will enable us to account for many of the ordinary meteorological phenomena, such as the formation of dew, and the fall of rain and snow.

Formation of Dew. Any portion of atmospheric air contains vapour in a greater or less degree, and may be saturated with it; if so, the slightest fall of temperature will produce condensation. If any solid in contact with the atmosphere be cooled down until its temperature is below that which corresponds to the saturation of the air around it, condensation will take place, and the condensed vapour will be deposited in the form of *dew* upon the surface of the body.

This accounts for the *dew* with which the ground is covered after a clear night.

Heat radiates from the ground, and from the bodies upon it, and unless there are clouds from which the heat would be radiated back, the surfaces are cooled and the vapour in the stratum of the atmosphere immediately above condenses and falls in small drops of water on the surface. Any kind of covering will more or less prevent the formation of dew beneath; very little dew, for instance, will be found under the shade of large trees. It will be seen moreover that good radiators are most abundantly covered with dew, very smooth surfaces being almost entirely free from it. This is in accordance with the facts stated above of the radiation of heat.

Hoar Frost. If after the deposition of dew the temperature fall below the freezing point, the dew is then frozen and becomes hoar frost.

The fogs seen at night on low lying or marshy lands are due to the same cause. The air is charged with moisture to saturation, and the cooling of the surface extends sometimes through three or four feet of the atmo-

sphere, producing a thick fog close to the ground, while the air above is quite clear.

144. *Clouds and Rain.* Clouds are formed by the condensation of the vapour in the upper regions of the atmosphere. The reduction of temperature requisite for condensation may occur from several different causes; a mass of air and vapour in motion may rise into a colder region or may come into contact with a larger mass of colder air, so that when the two are mingled together the temperature may not be sufficient to maintain the elasticity of the vapour.

The fact that the clouds remain suspended may be explained in various ways. It seems highly probable that in the process of condensation the vapour assumes the form of small vesicles of water containing air, and therefore not necessarily of greater specific gravity than the medium in which they are formed. Or, again, if the particles do descend, they may, as they fall into a space in which the temperature is higher, be gradually absorbed, and if new vapour be formed above, the appearance of a stationary cloud would consist with the fact of a continuous fall in the constituent particles of the cloud itself.

The cloud which is often seen about the top of a mountain is not unfrequently of this kind. A mass of warm air charged with moisture travels past a mountain, and by contact with it condensation is caused in that portion which is near to the mountain. As the condensed vapour is drifted away, it is again absorbed by the warm air around it, and thus the apparently fixed cloud merely represents a state through which the warm air passes, and from which it emerges.

If a cloud be very highly charged with moisture, and a further reduction of temperature take place, the vapour condenses still further into small drops, and descends in the form of *rain*.

145. When vapour is being condensed, if the temperature fall below the freezing point, *snow* is formed; and if rain as it falls pass through a region of the air in which the temperature is below the freezing point, the drops of rain are congealed and descend in the form of *hail*.

Fogs and *mists* are clouds formed near the earth's surface and in contact with it. The light summer rain which sometimes falls about sunrise or sunset without the appearance of a cloud is due to the same cause, the air becoming suddenly colder, and the vapour in consequence being rapidly condensed.

146. *Illustration.* The phenomena of dew and hoar frost may be obtained on a small scale by simply putting ice into a glass of water. The outside of the glass will soon be covered with a delicate dew, which after a short time freezes, and the glass is then covered with hoar frost.

The explanations of the preceding articles will enable an observer to account for most of the phenomena which depend on the existence of aqueous vapour in the atmosphere.

147. *Sea and land breezes.* Winds are partly due to changes of temperature; if, for instance, the air in the neighbourhood of any particular region become heated, it will expand and rise, its place being filled by air from other regions, and hence a wind towards the heated region.

In hot countries on the sea-coast it is noticed that during the day the wind in general blows from the sea, and during the night from the land. During the day the land becomes heated and retains heat; hence the air above it rises, and the cooler air flows in from the sea. But during the night the land cools by radiation while the temperature of the sea remains nearly the same; hence the land breeze.

Dew-Point and Hygrometers.

148. *The Dew-point* is the temperature at which the vapour in the atmosphere begins to condense.

To determine the dew-point a glass vessel must be cooled until dew begins to be deposited upon it, and its temperature must be then observed; again, observe the temperature at which the dew disappears; a mean between the two may be taken as the dew-point.

149. *Tension of vapour in the air.* If the dew-point be ascertained we can infer the pressure of the vapour in the air by means of the tables before referred to of the relation between the temperature and the saturating density.

For if t' be the dew-point, and ϖ' the corresponding pressure, t the temperature of the air, and ϖ the required pressure,

$$\varpi : \varpi' :: 1 + at : 1 + at',$$

and, the pressure being known, the quantity of vapour in the atmosphere can be determined.

150. *Hygrometers* are instruments for determining the quantity of vapour in the atmosphere, or, in other words, the degree of saturation.

This is measured by the ratio of the tension of the vapour in the air to the saturating tension.

Thus if, in the case of Art. (137), ϖ'' be the saturating tension at the temperature t , $\frac{\varpi}{\varpi''}$ is the measure required.

Hygrometers may be constructed of any substance which is affected by the amount of moisture in the air, such as a piece of cord which elongates as the quantity of vapour in the air diminishes, or a piece of seaweed, which is exceedingly sensitive to hygrometric changes in the atmosphere.

One of the hygrometers most in use is the *wet and dry bulb Thermometer*. It consists of two mercurial thermometers near each other, one of which is covered with muslin, and kept constantly wet by letting a portion of the muslin drop in a cup of water. The moisture from the muslin evaporates, and, as evaporation is always accompanied by cooling, the wet bulb thermometer falls, and, the drier the air is, the greater will be the difference between the two thermometers. Empirical formulæ and tables have been constructed by means of which the tension of the vapour can be inferred from the readings of the thermometers*.

* See Mr Glaisher's pamphlet *On the Wet and Dry Bulb Thermometer*.

The direct determination of the dew-point is a troublesome process, and the quantity of vapour in the air is most easily found by means of the hygrometer just described, and its accompanying tables.

Dilatation of Liquids.

151. In general, all solid and liquid bodies expand under the action of heat, and contract when heat is withdrawn. We have before had occasion to take account of the expansion of mercury, which is within certain limits proportional to the increase of temperature. This is also the case with solid bodies, such as glass and steel.

For water and aqueous liquids generally, the law of expansion is unknown; the rate of expansion is not constant for a constant increase of temperature, but beyond a certain limit becomes more rapid as the temperature rises.

Maximum density of water. It is a remarkable property of water that its density is a maximum at a temperature of about 4° C. or 40° F., and whether the temperature increases or decreases from this point, the water expands in volume.

152. *Freezing.* When the temperature descends to the freezing-point, a still further expansion takes place at the moment of congelation. This is sufficiently proved by the fact that ice floats in water, but it may also be rendered very distinctly evident by a direct experiment. Fill a small iron shell with water, and close the aperture with a wooden plug; if the shell be then exposed to a freezing temperature, the water within will freeze, and at the instant of congelation, the plug will be shot out with considerable violence*.

153. *Formation of ice on the surface of a lake.* It is known that ice is formed much more rapidly on the surface of shallow than on the surface of deep water; and this fact we can now account for. As the air cools, the

* The results of Playfair and Joule give 3°.945 C. as the temperature at which the density is a maximum. Prof. Miller, *Phil. Transactions*, 1856.

The temperatures at which liquids freeze are different for different liquids, but fixed for each liquid. Thus mercury freezes at a temperature -40° C.

water at the surface cools, and being contracted becomes heavier than the water beneath. The surface strata then descend, and the water from beneath rises and becomes cooled in its turn, and this process will go on until the whole of the water has attained its maximum density, after which it will remain stationary, and the upper strata being further cooled will expand and finally congeal. It is clear that the deeper the water is the longer will be the time which elapses before the whole of the water has attained its maximum density.

154. *Ebullition.* When heat is applied to water, it expands gradually until, at a certain temperature, bubbles are formed and steam is given off.

This temperature is the boiling-point, and it has been mentioned before that it depends upon the atmospheric pressure.

The bubbles are first formed by the expansion of the air which water contains. If water be heated from below, the lower strata expand and rise, the upper strata descending and becoming heated in succession, and air-bubbles ascend. As the temperature increases, small bubbles of vapour ascend, but do not always reach the surface, as they may be condensed in the less heated strata above. Finally, larger bubbles are formed, and, the whole mass being heated, ascend to the surface and give off steam, which becomes visible by a slight condensation in the air above.

These bubbles are formed when the tension of their vapour is equal to the pressure they sustain, and this explains why a diminution of atmospheric pressure permits the process of ebullition at a lower temperature; and, on the other hand, that an increase of atmospheric pressure raises the temperature of ebullition.

For instance, under a pressure of two atmospheres, the boiling-point is raised 20° C., and, if the atmospheric pressure be diminished one-half, the boiling-point is lowered about 18° .

This accounts for the fact that water boils at a low temperature on the tops of mountains, and on high table-lands.

Specific Heat.

155. It is found that a certain quantity of heat must be expended in order to raise the temperature of a mass of any substance by a given amount. The requisite quantity of heat depends on the nature of the substance and also on its mass, and for any particular substance it may be at once assumed that the quantity of heat required to raise the temperature one degree is directly proportional to the mass of the substance.

In general, the amount of heat required to change the temperature of a given mass from t^0 to $(t+1)^0$ is the same for all values of t .

Hence for the same substance the quantity of heat expended in changing the temperature from t^0 to t'^0

$\propto t' - t$ when the mass is given,

and \propto the mass when $t' - t$ is given,

and therefore generally $\propto m(t' - t)$, if m be the mass.

If this be taken equal to $cm(t' - t)$, c is called the specific heat of the substance, and it is the measure of the amount of heat which will raise by 1^0 the temperature of the unit of mass.

If two masses m, m' , of the same substance, at temperatures t, t' , be mixed together, and if τ be the temperature of the mixture, then, since the amount of heat lost by one is gained by the other,

$$mt - m\tau = m'\tau - m't,$$

$$\text{or } mt + m't = (m + m')\tau.$$

156. For different substances the quantity c has different values; thus it is found that water requires about 28 times as much heat as mercury in order to change the temperature by a given amount, and the specific heat of mercury is therefore less than that of water in the ratio of 1 : 28.

The specific heat of a gas must be considered from two different points of view, for we may suppose the volume of a gas constant, and investigate the amount of heat required to raise the temperature 1^0 , or we may suppose the pressure constant, the latter supposition permitting the expansion of the gas.

The specific heat in the second case exceeds the specific heat in the first case by the amount of heat disengaged when the gas is suddenly compressed into its original volume.

157. The specific heat of water is usually taken as the unit, and one of the methods of finding the specific heat of a substance is by immersing it in a given weight of water, and observing the temperature attained by the two substances.

Thus, if M be the mass of a body, T its temperature, and C its specific heat,

m' and m the masses of a vessel and of the water in it, and t their common temperature,

τ the temperature of the whole after immersion, and C' the specific heat of the vessel,

$$CM(T - \tau) = m(\tau - t) + C'm'(\tau - t),$$

since the quantity of heat lost by the body is equal to that gained by the water and the vessel.

If C' be known, this equation determines C ; and C' , if unknown, can be found by pouring water of a known temperature into the vessel at some other known temperature.

The following are approximate values of the specific heats of a few substances.

Water.....	1.
Thermometer-glass...	.198
Iron.....	.114
Zinc.....	.1
Mercury.....	.03
Silver.....	.06
Brass.....	.09.

EXAMINATION ON CHAPTER VIII.

1. A CUBIC foot of air having a pressure of 15 lbs. on a square inch is mixed with a cubic inch of compressed air, having a pressure of 60 lbs. on a square inch; find the pressure of the mixture, when its volume is 1729 cubic inches.

2. State the conditions under which a space is saturated with vapour.

3. A vessel of water is left in a close room for some time; what would be the effect of bringing a quantity of ice into the room?

4. Explain the radiation, conduction, and convection of heat. Why is a cloudy sky not favourable to the deposition of dew?

5. How do you account for the long trail of condensed steam which often follows a locomotive in rainy weather?

6. Define the Dew-point, and explain the use of the Wet and Dry Bulb Thermometer.

7. Explain why it is difficult to heat water from its upper surface.

8. If a piece of ice be put into a glass of water, the external surface is soon covered with a fine dew; account for this fact.

9. Explain what is meant by Specific Heat.

Three gallons of water at 45° are mixed with six gallons at 90° ; what is the temperature of the mixture?

10. At great altitudes it is sometimes found that a sensation of discomfort is felt; the lips crack and the skin of the hands is roughened; how do you account for these facts?

Can you give any reason why an east wind in England sometimes produces similar effects?

EXAMPLES.

1. Two volumes V, V' of different gases, at pressures p, p' , and temperature t are mixed together; the volume of the mixture is U , and its temperature t' , determine the pressure.

2. Two vessels contain air having the same temperature t , but different pressures p, p' ; the temperature of each being increased by the same quantity, find which has its pressure most increased.

If the vessels be of the same size, and be allowed to communicate with each other, find the pressure of the mixture at a temperature zero.

3. A glass vessel weighing 1 lb. contains 5 oz. of water, both at 20° , and 2 oz. of iron at 100° is immersed; what is the temperature of the whole, taking .2 as the specific heat of glass and .12 of iron?

4. An ounce of iron at 120° , and 2 oz. of zinc at 90° are thrown into 6 oz. of water at 10° contained in a glass vessel weighing 10 oz.; what is the final temperature, taking .1 and .12 as the specific heats of zinc and iron?

5. The pressure of a quantity of air, saturated with vapour, is observed; the mixture is then compressed into half its former volume, and, after the temperature has been lowered until it becomes the same as at first, the pressure is again observed; hence find what would be the pressure of the air (occupying its original space) if it were deprived of its vapour without having its temperature changed.

6. It is related of a place in Norway that a window of a ball-room being suddenly thrown open, a shower of snow immediately fell over the whole of the room. Account for this phenomenon.

7. A drop of water is introduced into the tube of a common barometer which just does not evaporate at the higher of the temperatures t_1^0 , t_2^0 .

Given that the elasticity of vapour increases geometrically as the temperature increases arithmetically, shew that if E_1 , E_2 be the errors of the above barometer at temperatures t_1^0 , t_2^0 , the common ratio of the geometric progression for an increase of temperature of 1^0 in the case of vapour of water is

$$\frac{\{E_1 (1 - et_1)\}^{\frac{1}{t_1 - t_2}}}{\{E_2 (1 - et_2)\}}$$

e being the coefficient of expansion for mercury.

8. A closed cylinder contains a piston moveable by means of a rod passing through an air-tight collar at the top of the cylinder. The piston is held at a distance from the bottom of the cylinder equal to one-third of its height, and vapour is introduced above and below of a known pressure, the temperature of the cylinder being such as will support vapour of twice the density without condensation. The piston on being left to itself sinks through two-ninths of the height of the cylinder. Prove that the weight of the piston is five-fourths of the pressure of the vapour upon either side at first.

9. A flask is partially filled with water which is caused to boil until the air is expelled, and then the flask is corked and allowed for a short time to cool. The flask is then placed in cold water, and it is found that the water in it recommences boiling. Explain this phenomenon.

CHAPTER IX.

Rotating Liquid.

158. **W**HEN liquid in a vessel is set rotating, it is known that the surface assumes a hollow form; by the help of a dynamical law we can determine what this form is.

If a liquid, contained in a vessel which rotates uniformly about a vertical axis, rotates uniformly, as if rigid, with the vessel, its surface is a paraboloid.


Every particle of the liquid moves uniformly in a horizontal circle, and therefore whatever the forces may be which act on any particle, their resultant must be a horizontal force tending to the centre of the circle and equal to $m\omega^2r$, where r is the distance of the particle from the axis, m its mass, and ω the angular force of the liquid*.

Since there is no relative displacement of the molecules of the liquid, we can imagine that it has no rotation, and is in equilibrium under the action of gravity and of the imaginary force $m\omega^2r$ all acting outwards.

We assume from symmetry that the surface is a surface of revolution.

Let AG be the vertical axis of revolution, and consider a point P on the surface. Round P as centre describe a very small circle, and take this small circular area as the base of a very thin circular cylinder of liquid.

If m be the mass of the element of liquid, thus imagined, it is in equilibrium under the action of gravity, of $m\omega^2r$ outwards, of the atmospheric pressure on its surface, of the liquid pressure on its



* See Garnett's *Dynamics*, or Parkinson's *Mechanics*.

inside flat end, and of the liquid pressures on its curved surface.

These latter pressures, being equal to the atmospheric pressure, are of themselves in equilibrium, and it therefore follows that the resultant of gravity, mg , and of the force $m\omega^2 r$ must be normal to the surface*.

Let the normal at P meet the axis in G , and draw PN horizontal.

Then two forces acting in directions of the lines PG , GN have their resultant, which is $m\omega^2 PN$, in the direction PN , and therefore by the triangle of forces,

$$NG : PN :: mg : m\omega^2 PN,$$

and
$$\therefore NG = \frac{g}{\omega^2}.$$

Now NG is the subnormal, and it is known that in a parabola the subnormal is constant, while it can also be shewn that the parabola is the only curve which has this property.

The vertical section in the figure is therefore a parabola the latus rectum of which is $\frac{2g}{\omega^2}$, and the surface is a paraboloid.

It will be seen that this result is independent of the form of the containing vessel. The axis of rotation, in fact, may be within or without the fluid, but in any case it will be the axis of the paraboloidal surface.

If the vessel were a surface of revolution, having the axis of rotation for its axis, it would not be necessary theoretically that the vessel should rotate. However, by making it rotate with the liquid, we get rid of the practical difficulty which would in this case arise from the friction between the fluid and the surface of the vessel.

159. *To find the pressure at any point.*

Take any point Q in the fluid, and describe a small vertical prism having Q in its base, which is to be taken horizontal.

* A particular case of the general theorem that, in fluids at rest under any forces, the resultant force at any point is normal, at that point, to the surface of equal pressure passing through it.

The prism PQ of liquid rotates uniformly under the action of the pressure around it, but its weight is entirely supported by the pressure on its base.

Hence if p be the pressure, and ρ the density,

$$p = g\rho PQ.$$

Now

$$PQ = OM - ON = \frac{\omega^2 QN^2}{2g} - ON,$$

$$\text{and } \therefore p = \rho \left(\frac{1}{2} \omega^2 QN^2 - gON \right);$$


and we thus obtain the pressure in terms of the horizontal and vertical distances from the vertex of the paraboloid.

It must be observed that ON is measured upwards and that if Q be lower than O , the equation for p is

$$p = \rho \left(\frac{1}{2} \omega^2 QN^2 + gON \right).$$

160. To find the resultant vertical pressure of a rotating liquid on any surface.


Let PQ be the surface, and draw vertical lines from its boundary to the surface; then the weight of the in-



cluded portion $PABQ$ of liquid being entirely supported by PQ , it follows that the resultant vertical pressure on PQ is equal to the weight of the liquid above it.

If the surface PQ be pressed upwards, as in the figure.

then, continuing the free surface AOP of the liquid, it can be shewn, as in Art. (52), that the resultant vertical pressure upwards on PQ is equal to the weight of the fluid which would be contained between the paraboloidal surface, the surface PQ , and vertical lines through the boundary of PQ .




161. DEF. A surface of equal pressure is the locus of points at which the pressures are the same.

If lines be drawn vertically downwards from all points of the surface equal to PQ (fig. Art. 159), it is clear that the pressures at their ends will be the same as at Q ; and, as these ends lie on the surface of a paraboloid equal to the surface paraboloid, it follows that all surfaces of equal pressure are in this case paraboloids.

162. *Floating bodies.* If a body float in a rotating mass of fluid, in a position of relative equilibrium, it is evident by the same reasoning, as in the case of a fluid at rest, that the weight of the body must be equal to the weight of the fluid displaced.

163. *Figure of the Earth.* A large portion of the earth's surface is covered with water, and, if it were not



for the earth's rotation, its surface would be a sphere having its centre at the centre of the earth.

For simplicity, imagine a solid sphere surrounded by water, and suppose the whole to be in rotation about a diameter CB of the sphere. Consider an elementary por-

tion P of the water, which describes a circle of radius PN uniformly. The attraction of the solid sphere in the direction PC , combined with the resultant fluid pressure in direction of the normal at P , must have as their resultant the force $m\omega^2PN$ in direction PN . Hence the normal at P must be inclined, as in the figure, towards the axis, and the form of the surface must be oblate.


Supposing the earth a large fluid mass, it is shown by mechanical considerations that the form would be an oblate spheroid.

It is hence seen that the normal to the surface of still water, that is, the vertical, at any point of the earth's surface is not in direction of its centre, except at the poles and the equator.

EXAMPLES.

164. (1) *A fine tube, ABCD, of which the equal branches AB, CD, are vertical, BC being horizontal, is filled with liquid, and made to rotate uniformly about the axis of AB; to find how much liquid will flow out of the end D.*

The liquid will flow out until the surface in AB is the vertex



of a parabola passing through D , and having its axis vertical and latus rectum equal to $\frac{2g}{\omega}$.

If then O be the vertex of the parabola,

$$BC^2 = \frac{2g}{\omega^2} AO.$$

This gives AO , and determines the quantity required.

If however AO be greater than AB , the surface of the liquid will be in BC , at P suppose.

In this case we have $BC^2 = \frac{2g}{\omega^2} AO'$,

$$\text{and } BP^2 = \frac{2g}{\omega^2} BO' = \frac{2g}{\omega^2} (AO' - AB),$$

which determines the position of P .

(2) *A straight tube AB , filled with liquid, is made to rotate uniformly about a vertical axis through A ; to find how much flows out at B .*

Let $OAB = \alpha$, and imagine a parabola, latus rectum $\frac{2g}{\omega^2}$, to

be drawn touching the axis of the tube, and having its axis coincident with the vertical through A .

Then if P be the point of contact, all the fluid above P will flow out.

To find P ,


$$PN^2 = \frac{2g}{\omega^2} \cdot ON$$

$$= \frac{g}{\omega^2} AN, \text{ since } OA = ON,$$

and $PN = AN \tan \alpha$;

$$\therefore AN = \frac{g}{\omega^2} \cot^2 \alpha,$$

$$\text{and } AP = \frac{g}{\omega^2} \frac{\cos \alpha}{\sin^2 \alpha}.$$



No fluid will flow out unless $AP < AB$, that is, unless

$$\omega^2 > \frac{g \cos \alpha}{AB \sin^2 \alpha}.$$

It will be seen that P is the position of relative equilibrium of a heavy particle in the rotating tube.

(3) *Let the end B be closed and the tube AB , rotating as in Ex. (2), be only partly filled with liquid; it is required to find the circumstances of relative equilibrium.*

Let AC be the portion of tube filled with liquid (fig. of previous article).

Draw the parabola touching in P as before: then, if C is below P , no change takes place, but if C is above P , the portion PC of liquid will flow to the upper end of the tube and remain there.

(4) A semicircular tube APB is filled with liquid and rotates uniformly about the vertical diameter AB ; it is required to find where a hole may be made in the tube through which all the liquid will flow out.

Draw a parabola touching the tube at P and having its vertex in BA , axis vertical, and latus rectum B equal to $\frac{2g}{\omega^2}$.

Then, if an aperture be made at P , the whole of the liquid, being above the paraboloidal surface, will flow out through P .


To find its position we have

$$PN^2 = \frac{2g}{\omega^2} \cdot ON = \frac{g}{\omega^2} NT;$$

$$\text{but } PN^2 = CN \cdot NT;$$

$$\therefore CN = \frac{g}{\omega^2}, \text{ which determines } P.$$

If a be the radius (CA) of the tube, and if $\omega^2 < \frac{g}{a}$, then $CN > CA$, and the aperture must be made at A .



(5) In a mass of liquid, rotating about a vertical axis, a very small sphere, of greater density than the liquid, is immersed, and supported by a string fastened to a point in the axis; it is required to find the position of relative equilibrium.

For one position of equilibrium it is evident that the string can be vertical, but we can shew that the sphere may rest with the string inclined at a certain angle (θ) to the vertical.

Let V be the volume of the sphere, r its distance from the axis in the position of relative equilibrium, and ρ the density of the liquid.

To find the pressure of the liquid on the sphere, imagine it removed and its place supplied by a solidified portion V of the liquid;

The resultant liquid pressure must support the weight $g\rho V$, and also supply the horizontal pressure necessary to maintain the circular motion, i. e. $\rho V\omega^2 r$.

Hence if ρ' be the density of the sphere, and t the tension of the string, we must have, for equilibrium,

$$t \sin \theta + \rho V\omega^2 r = \rho' V\omega^2 r,$$

$$t \cos \theta + \rho Vg = \rho' Vg,$$

$$\text{and } \therefore \tan \theta = \frac{\omega^2 r}{g}.$$

The position is therefore the same as if the sphere and string were in motion as a conical pendulum.


It will also be seen that the string coincides with the direction of the normal to the surface of equal pressure which passes through the centre of the sphere.

(6) *A cylindrical vessel contains liquid, which rotates uniformly about the axis of the cylinder; to find the whole pressure on its surface.*

Let AOB be a vertical section of the surface, r the radius of the cylinder.

We have shewn, Art. (159), that the pressure varies as the depth below the surface, and in this case the level of the free surface is the same for all points on the curved surface of the cylinder.

Hence the whole pressure on the curved surface



$$= g\rho 2\pi r \cdot AD \cdot \frac{1}{2} AD = \pi g\rho r \cdot AD^2.$$

Let h be the height of the liquid when at rest.

It is known that the volume of a paraboloid is half that of the cylinder on the same base and of the same height, and therefore the surface of the liquid at rest would bisect AN .

$$\text{But } ON^2 = \frac{2g}{\omega^2} AN, \text{ or } r^2 = \frac{2g}{\omega^2} AN;$$

$$\therefore AD = h + \frac{1}{2} AN = h + \frac{\omega^2 r^2}{4g}.$$

Hence the whole pressure is given in terms of h .

Also the pressure on the base is equal to the weight of the fluid, i. e. $g\rho\pi r^2 h$.

NOTE ON CHAPTER IX.

The question of the form of the surface of a rotating fluid appears to have been first discussed by Daniel Bernoulli, in his *Hydrodynamica*, which was published in 1733. He there proves that the form is that of a paraboloid, and 5 years after Clairaut, in his *Figure de la Terre*, gives a similar proof, at the same time quoting Bernoulli.

From the remarks of Art. 17, it follows that the paraboloidal form will be exactly true for viscous liquids rotating in the same manner. The point of argument lies in the phrase, *as if rigid*, for, without this condition, it would not be possible to imagine the liquid in a state of equilibrium. It must not be inferred that the paraboloidal form is that which would be assumed by a liquid set in rotation by ordinary mechanical means. The internal friction of a liquid, communicated from the surface of a rotating vessel may produce the effect, if the revolution be maintained long enough.

Theoretically, we can imagine the effect produced by enclosing ice in a strong vessel with a paraboloidal upper surface, making it rotate, and then melting the ice by pressure, or otherwise. The melted ice would retain the rotation as if rigid, and it might perhaps be possible to procure an approximation to the paraboloidal surface.

If a cup of tea be rapidly stirred, a convex surface is produced, having a hollow in the middle, but, in motion of this kind, the angular velocity decreases at increasing distances from the centre, and there is a constant displacement of the relative positions of the molecules of liquid. This is the case of Rankine's free circular vortex, and its discussion belongs to the domain of Hydrodynamics. (See *Hydromechanics*.)

EXAMPLES.

1. LIQUID contained in a closed vessel rotates uniformly about a vertical axis; prove that the difference of the pressures at any two points of the same horizontal line varies as the difference of the squares of the distances of the two points from the axis of rotation.

2. A hollow paraboloid of revolution with its axis vertical and vertex downwards is half filled with liquid. With what angular velocity must it be made to rotate about its axis in order that the liquid may just rise to the rim of the vessel?

3. If a solid cylinder float in a liquid which rotates about a vertical axis having its axis coincident with the axis of revolution, determine the portion of its surface which is submerged, the dimensions of the cylinder and the densities of the liquid and cylinder being given.

4. An open vessel, containing two liquids which do not mix, revolves uniformly round a vertical axis; find the form of the common surface.

5. A conical vessel open at the top and filled with liquid rotates about its axis; find how much runs over, 1st, when ω is less, and, 2nd, when ω is greater than $\sqrt{\frac{g}{h}} \cot \alpha$, h being the height of the cone, and α its semivertical angle.

6. A hemispherical bowl is filled with liquid, which is made to rotate uniformly about the vertical radius of the bowl; find how much runs over.

7. An elliptic tube, half full of liquid, revolves about a fixed vertical axis in its own plane, with angular velocity ω ; prove that the angle which the straight line joining the free surfaces of the liquid makes with the vertical is $\tan^{-1} \frac{g}{p\omega^2}$, where p is the perpendicular from the centre on the axis.

8. A closed cylindrical vessel, height h and radius a , is just filled with liquid, and rotates uniformly about its vertical axis; find the pressures on its upper and lower ends, and the whole pressure on its curved surface.

9. A hemispherical bowl, just filled with liquid, is inverted on a smooth horizontal table, and rotates uniformly about its vertical radius; find what its weight may be, in order that none of the liquid may escape.

10. A cylindrical vessel, containing water, rotates uniformly about its axis, which is vertical, the water rotating with it at the same rate; find the position of relative equilibrium of a small piece of cork which is kept under water by a string fastened to a point in the side of the vessel.

11. A vertical cylinder, of height h and radius a , is half full of water, which rotates uniformly about the axis; prove that the greatest angular velocity which can be imparted to the water without causing an overflow is $\sqrt{2gh} \div a$.

12. A conical vessel, of height h and vertical angle 60° , has its axis vertical and is half filled with water; prove that the greatest angular velocity which the water can have without overflowing is $\sqrt{\frac{2g}{h}}$.

CHAPTER X.

Tension of Vessels containing Fluids.

165. **I**F a cylindrical vessel contain liquid, the pressure of the liquid will produce a strain or tension in the substance of which the vessel is formed. We may imagine the vessel formed of some thin flexible substance, such as silk or paper, and it is obvious that if this substance be not strong enough, it will be torn asunder by the pressure of the liquid.

We proceed to investigate the relation between the pressure and the tension produced by it.

Measure of tension. Imagine a hollow cylindrical vessel formed of a thin flexible substance to be filled with a gas at a given pressure, so that the tension may be the same throughout.

Divide the surface along a generating line, length l , and let T be the whole force required to keep the two parts together;

then, if $T=tl$, t is the tension along any unit of length.


If the cylinder be vertical and filled with water, so that the pressure and therefore the tension vary at different depths, then the tension t at any point is the tension that would be exerted along an unit of length, if it were the same throughout the unit as it is at the point in question.

166. *A vertical cylindrical vessel contains fluid; to find the relation between the pressure and tension.*

The pressure being the same at all points of the same

horizontal plane, it follows that the tension will be the same at all points of the same horizontal section.

Let PQ , $P'Q'$ be small portions of two horizontal sections very near each other. PP' and QQ' being vertical. The dimensions of PQ are taken so small that the pressure and tension at all points of it are sensibly the same.



Let p , t be the pressure and tension; then $t.PP'$, $t.QQ'$ are the horizontal forces acting on the portion PQ of the surface at the middle points A , B of its ends, and these forces must counterbalance the pressure of the liquid, which is $p.PP'.PQ$.


This resultant pressure acts in the direction CE bisecting the angle ACB , and the two tensions in the directions of the tangents at P and Q .

Hence, resolving the forces in the direction CE ,

$$\begin{aligned} p.PP'.PQ &= 2t.PP' \sin \frac{1}{2}ACB \\ &= 2t.PP' \cdot \frac{1}{2} \cdot \frac{AB}{r} = \frac{t}{r}.PP'.PQ, \end{aligned}$$

if r be the radius of the cylinder,

and $\therefore t = pr$.



If the cylinder contain a gaseous fluid of which the pressure is sensibly the same throughout its mass, the relation $t = pr$ is true at every point, whether the axis be vertical or not.

This result can also be obtained by considering the equilibrium of a semi-circular portion of thickness PP' , for the resultant pressure will be parallel to the tensions at the two ends, and will be equal to the pressure on the projected area $2r \cdot PP'$, so that

$$2t \cdot PP' = p \cdot 2r \cdot PP', \text{ or } t = pr.$$

167. If the pressure along the arc AB is a variable quantity the cylindrical form is not a form of equilibrium for a vessel formed of thin flexible material, but, taking r as the radius of curvature at E , the relation, $t = pr$, is true of every point.

For example consider the *Lintearia*, which is the form assumed by a rectangular piece of a thin membrane, two opposite sides of which are fastened to the sides of a box, while the other sides fit the box closely, so that liquid can be poured in without escaping.

The figure is a section of the cylindrical surface so formed, by a plane perpendicular to its generating lines, BC being the surface of the liquid.



The tension (t) along BAC is constant, because the liquid pressure is normal, and, if r be the radius of curvature at P ,

$$t = pr = qprPN, \therefore \frac{1}{r} \propto PN,$$

i. e. the curvature at P is proportional to the depth below the surface.

This curve is the same as the *Elastica*, the curve formed by a bent rod, and is also, as will be seen subsequently, the same as the *Capillary curve*.


168. *A spherical surface contains gas at a given pressure, it is required to find the tension at any point.*

From symmetry we may take the tension to be the same at every point.

Moreover, if any line be drawn on the surface we may assume that the tension between the two portions parted by that line acts in a direction perpendicular to it.

Consider a very small square portion of the surface $ABCD$, and let t be the tension, O the centre of the sphere, and r its radius.

Then a being the length of any side of the square, there are four forces, each equal to ta , acting at the middle points of the lines AB , BC , CD , DA , perpendicular to these lines and tangential to the surface, and the resultant of these forces must counterbalance the pressure of the gas.



The resultant of ta on AB and CD , in the direction EO ,

$$= 2ta \cdot \sin FOE = 2ta \frac{EF}{r} = \frac{t}{r} a^2;$$

and similarly the resultant of the tensions on AC and BD

$$= \frac{t}{r} a^2;$$

therefore, if p be the pressure of the gas,

$$2 \frac{t}{r} a^2 = pa^2, \text{ and } 2t = pr.$$

Instead of taking a small element we may consider the equilibrium of a hemisphere, under the action of the tension $2\pi rt$, and of the resultant pressure which is equal to the pressure on a circular area of radius r , so that

$$2\pi rt = \pi r^2 p, \text{ or } 2t = pr.$$

Hence it appears that a spherical vessel is twice as strong as a cylindrical vessel of the same material and the same radius.

169 We have not compared with each other the tensions of vessels formed of substances of different thickness. To do this it will be seen that for a given value of the tension t , as we have measured it, the intrinsic strain of any substance will be diminished by increasing the thickness.

Now if e be the thickness of any flexible lamina, and if $t = e\tau$, then τ will be the tension of an unit of area of the section, and for the comparison of different thicknesses, this latter measure of tension must be employed.

Ex. *A bar of metal one square inch in section can sustain a weight of 1000 lbs., and of this metal a cylinder is made one-twentieth of an inch in thickness, and one foot in diameter; find the greatest fluid pressure which the cylinder can sustain.*

In this case the greatest possible value of τ is 1000, and the greatest value of $t = \frac{1}{20} \cdot \tau = 50$;

$$\therefore p = \frac{t}{r} = \frac{50}{6} = 8\frac{1}{3} \text{ lbs.}$$


Hence $8\frac{1}{3}$ lbs. per sq. inch is the greatest pressure which can be applied without bursting the cylinder.

170. *A conical vessel, formed of a flexible substance, is held by the rim with its vertex downwards, and is filled with liquid; it is required to find the tension at any point in the direction of the generating line passing through the point.*

Let PP' be a horizontal section of the cone. It is obvious that along the section PP' the tension is the same at any point and is in direction of the generating line through that point.

Let then t be the tension, which is at all points of the circle PP' in a direction inclined at an angle α to the vertical, if 2α be the vertical angle.

The vertical resultant of the tension on the whole circle PP' , that is, $2\pi \cdot PN \cdot t \cos \alpha$, is equal to the resultant vertical pressure on the surface POP' .



Now this pressure

$$\begin{aligned} &= \text{weight of fluid } POP' + \text{weight of fluid } P'Q \\ &= g\rho \left(\frac{1}{3} \pi PN^2 \cdot ON + \pi PN^2 \cdot PQ \right), \end{aligned}$$

and therefore if $ON=x$, and $OE=h$,

$$2\pi x \tan \alpha \cdot t \cos \alpha = g\rho \pi x^2 \tan^2 \alpha \left(\frac{x}{3} + h - x \right),$$

or
$$t = \frac{1}{2} g\rho \frac{\sin \alpha}{\cos^2 \alpha} \left(hx - \frac{2x^2}{3} \right).$$

Since
$$hx - \frac{2x^2}{3} = \frac{2}{3} \left\{ \frac{9h^2}{16} - \left(x - \frac{3h}{4} \right)^2 \right\},$$

it follows that t has a maximum value when $x = \frac{3h}{4}$.

A little consideration will shew that there is a horizontal tension at all points along a generating line, in a direction perpendicular to that line, but the investigation of this other tension would be beyond the limits which must be assigned to an elementary course, and must therefore be deferred to treatises taking a higher range.

EXAMPLES.

1. Two vertical cylinders of the same thickness and the same material, contain equal quantities of water; compare their greatest tensions.

2. Two cylindrical boilers are constructed of the same material, the diameter of one being three times that of the other, and the thickness of the larger one twice that of the other; compare the strengths of the boilers.

3. A bar of metal, one-fourth of a square inch in section, can support a weight of 1000 lbs.; find the greatest fluid pressure which a cylindrical pipe made of this metal can sustain, the diameter being 10 inches and the thickness one-tenth of an inch.

4. Equal quantities of the same material are formed into two thin spherical vessels of given radii; compare the greatest fluid pressures they will sustain.

5. The natural radius of an elastic spherical envelope containing air at atmospheric pressure is a , and, when a certain quantity of air is forced into it, its radius is b . It is then placed under an exhausted receiver and its radius becomes c . Find the quantity of air forced in, supposing that the increase of tension of the envelope varies directly as the increase of its surface.

6. The top of a rectangular box is closed by an uniform elastic band, fastened at two opposite sides, and fitting closely to the other sides; the air being gradually removed from the box, find the successive forms assumed by the elastic band, and when it just touches the bottom of the box, find the difference between the external and internal atmospheric pressures.

7. A vertical cylinder formed of a flexible and inextensible material contains water; find the tension at any point.

If this flexible cylinder be put into a square box, the width of which is less than the diameter of the cylinder, and water be then poured in to the same height as before, find the change in the tension at any depth.

8. An elastic and flexible cylindrical tube contains ordinary atmospheric air; if the ends be kept closed, and the pressure of the air inside be increased by a given amount, find the increase in the radius of the cylinder.

If the radius be doubled by a given increase of pressure, prove that the modulus of elasticity is in that case twice the tension that would have been produced in the cylinder, if inelastic, by the same increase of pressure.

9. An inelastic flexible cylindrical vessel, closed rigidly at the top, is filled with water, and the whole rotates uniformly about the axis of the cylinder, which is vertical; find the tension at any point.

CHAPTER XI.

Capillarity.

171. **W**HEN a glass tube, of very small bore, with its two ends open, is dipped in water it is observed that the water rises in the tube, and that it is in equilibrium with the surface of the water inside at a higher level than the surface outside. If the tube is dipped in mercury, it is found that the mercury inside is in equilibrium at a lower level than the mercury outside.

In either case, the ascent, or depression, is greater if the experiment be made with tubes of smaller bore.


If the surface of water be examined close to the vertical side of a vessel containing it, the surface will be found to be curved upwards, the water appearing to cling to, and hang from the wall, at a definite angle.

Phænomena of this kind, with others, such as those presented by drops of liquid, or by liquid films, are grouped together as being instances of *Capillary Action*.

172. Consider the equilibrium of a thin column of liquid PQ , as in the figure.

If π be the atmospheric pressure, pressure at $Q = \pi - g\rho \cdot QN$, and therefore, taking a as the cross section, the column PQ is acted upon by gravity and also by the force $gpaQN$ downwards, that is by

the force $gpaPN$ downwards, which is in some way counterbalanced.



This suggests the theory of the existence of a surface tension, the vertical resultant of which, acting on the upper boundary, at P , of the column, will exactly counterbalance the weight of the column PN .

Various facts support the idea of the existence of a surface tension. The familiar experiment of gently placing a needle on the surface of water, on which it will sometimes float, is a case in point. The needle appears to be supported on a thin membrane, which bends beneath its weight.

In summer weather insects may be seen on the surface of water, apparently indenting, without breaking through, the superficial membrane.

As the results of observation and experiment we can state two laws relating to surface tension.

(1) *At the bounding surface separating air from any liquid, or between two liquids, there is a surface tension which is the same at every point and in every direction.*

(2) *At the line of junction of the bounding surface of a gas and a liquid with a solid body, or of the bounding surface of two liquids with a solid body, the surface is inclined to the surface of the solid body at a definite angle, depending upon the nature of the solid and the liquids.*

In the case of water in a glass vessel the angle is acute; in the case of mercury it is obtuse. It will be seen that in a tube containing water, the top of the water is concave; in a mercurial barometer tube the top of the mercury is convex.

173. *Rise of a liquid between two plates.*

Taking the figure of Art. 172 as a vertical section perpendicular to the plates,

T the surface tension, α the constant angle at which the surface meets either plate, h the mean rise, and d the distance between the plates, we have, for the equilibrium of one unit of breadth of the liquid,

$$2T \cos \alpha = gphd,$$

so that h varies inversely as d .

Rise of a liquid in a circular tube.

Taking the same figure as a section through the axis, and r as the radius of the tube, we have

$$2\pi r T \cos a = g\rho\pi r^2 h,$$

and therefore h varies inversely as r .


It will be seen that the rise in a circular tube of radius r is the same as the rise between two plates at a distance r .

In each case the pressure at any point of the suspended column is less than the atmospheric pressure, and, if the column were high enough, this pressure would merge into a state of tension, which would still follow the law of fluid pressure, of being the same, at any point, in every direction.

The rise of sap in trees may perhaps afford an instance of this state of things.

174. *The Capillary Curve* is the form assumed by the liquid near a vertical wall.

Let PN be the height above the level of the water of a point P of this curve, and consider the equilibrium of the column $PQLN$, taking one unit of breadth perpendicular to the plane of the paper.



The resultant of the tensions at P and Q is in direction of the normal at the middle point of PQ , and, if r be the radius of curvature, it is equal to $T \cdot \frac{PQ}{r}$.

The vertical component of this resultant being equal to the weight of the column,

$$T \cdot \frac{PQ}{r} \cos \theta = g\rho PN \cdot NL,$$

where θ is the inclination to the vertical of the normal.
Now $NL = PQ \cos \theta$,

$$\therefore T = g\rho r \cdot PN,$$

i.e. the curvature at P is proportional to PN .

This is the property which we found to be true of the *Lintearia*, and which can be shewn to be also the characteristic property of the *Elastica*.

175. Liquid Films possess the characteristic property that the tension is the same at every point, and in every direction.

If t be the tension of a soap bubble of radius r , and p the difference between the internal and external air-pressure, then $2t = pr$.

Liquid films may be formed, and examined, by shaking a clear glass bottle containing some viscous liquid, or by dipping a wire frame into a solution of soap and water, and slowly drawing it out.

In this way films, apparently plane, can be obtained, shewing that the action of gravity is unimportant in comparison with the tension of the film.

These films give way and break under the least tangential action, and we therefore infer that the tension across any line is normal to that line.

We can hence deduce the property above stated. For, considering a small triangular portion, the actual tensions on the sides must be proportional to the lengths of the sides, and therefore the measures of the three tensions are the same.

If one part of the boundary of a plane film be a light thread, we can prove that it will take the form of an arc of a circle.

Since the tension of the film is at all points normal to the thread, it follows that the tension, t , of the thread is constant.

Let τ be the intrinsic tension of the film, and consider

the element PQ of the thread; for equilibrium, if r be the radius of curvature,

$$t \frac{PQ}{r} = \tau \cdot PQ,$$

and therefore r is constant.

176. *Energy of a plane film.*

In drawing out a film a certain amount of work is expended, and this represents the energy of the film.

Consider for instance a plane rectangular film $ABCD$, bounded by wires, and imagine the wire CD moveable on AC and BD ;

then releasing CD the film will draw CD towards AB , and the work done, if t be the tension, will be $t \cdot CD \cdot AC$. But, if S be the superficial energy per unit of area, the actual energy is $S \cdot CD \cdot AC$;

$$\therefore S = t,$$

i. e. the superficial energy per unit of area is equal to the tension per unit of length. (Maxwell's *Heat*, Chapter xx.)

177. *Energy of a soap bubble.*

If p be the difference between the internal and external pressures, and t the tension, the work done in expanding a soap-bubble from radius to a radius r' slightly greater is

$$p \cdot 4\pi r^2 (r' - r), \text{ or } 8\pi tr (r' - r).$$

If we assume t constant, the total work done in the formation of a bubble of radius $c = \Sigma 8\pi tr (r' - r)$,

and, taking $r' - r = \frac{c}{n}$ and $r = \frac{mc}{n}$,

$$\begin{aligned} \text{this} &= 8\pi t \Sigma_1^n \frac{mc^2}{n^2}, \text{ when } n \text{ is indefinitely increased,} \\ &= 4\pi tc^2, \end{aligned}$$

and therefore the superficial energy = t .

CHAPTER XII.

The Motion of Fluids.

178. **I**F an aperture be made in the base or the side of a vessel containing liquid, it immediately flows out with a velocity which is greater the greater the distance of the aperture below the surface. The relation between the velocity and the depth, taking the aperture to be small, was discovered experimentally by Torricelli.


The following is Torricelli's Theorem :

If a small aperture be made in a vessel containing liquid, the velocity with which the particles of fluid issue from the vessel, into vacuum, is the same as if they had fallen from the level of the surface to the level of the aperture;

that is, if x be the depth of the aperture below the surface, and v the velocity of the issuing particles,

$$v^2 = 2gx.$$

The experimental proof of this is that if the aperture be turned upwards, as in the figure, the particles of liquid will rise to the same level as the surface of the liquid in the vessel. Practically the resistance of the air and friction in the conducting-tube destroy a portion of this velocity, but experiments tend to prove the truth of the law, which moreover can be established as an approximate result of mathematical reasoning.



Assuming the principle of energy*, we can give a theoretical proof of the theorem.

Let K be the area of the upper surface of the liquid, and suppose that, during a short time, the height of the upper surface is diminished by a small quantity y , so that Ky is the volume which has flowed out through the orifice. Taking v as the velocity of efflux, the kinetic energy which has issued through the orifice is $\frac{1}{2} \rho Kyv^2$, and if we neglect the kinetic energy of the liquid in the vessel, this must be equal to the loss of potential energy, which is $g\rho Kyx$,
and $\therefore v^2 = 2gx$.

Form of a jet of liquid. If the aperture be opened in any direction not vertical, each particle of liquid having the same velocity, will follow the same path, which by the laws of Dynamics, is a parabola. Hence the form of the jet is a parabola.

Contracted vein. If the aperture be made in the base of a vessel, and if the base be of thin material, it is observed that the issuing jet is not cylindrical, but that it contracts for a short distance (a fraction of an inch) and then expands afterwards contracting gradually as it descends, and finally breaking into separate drops. The amount of contraction depends on the thickness of the vessel, and the size and form of the aperture.

The rate of Efflux is the rate at which the liquid flows out, and this clearly depends both on the velocity of the issuing particles, and the size of the aperture.

If k be the area of the aperture and v the velocity, then in an unit of time a portion of liquid will have passed through equal to a length v of a cylinder of which k is the base, and therefore vk is the quantity which flows out in an unit of time, that is, vk is the rate of efflux.

This is however not true unless the liquid issue from a pipe of some length, in which case there is no contracted vein. In general k must be taken as the section of the contracted vein, it being found that the velocity at the contracted vein is that which is given by Torricelli's theorem.

* See Maxwell's *Matter and Motion*.

179. *Steady motion.* When a fluid moves in such a manner that, at any given point, the velocities of the successive particles which pass the point are always the same, the motion is said to be steady. Thus if a vessel having a small aperture in its base be kept constantly full, the motion is steady.

180. *Motion through tubes of different size.* The continuity of a fluid leads to a simple relation between the velocities of transit through successive tubes. Thus if a liquid, after passing through a tube AB , pass through CD , the tubes being full, it is clear that during any given time the quantity which passes through a given plane AB in one tube must be equal to the quantity which passes any given plane CD in the other. Let k, k' be the areas of these planes, and v, v' the respective velocities at AB and CD . Then $kv, k'v'$ are the quantities which pass through in an unit of time, and therefore



$$kv = k'v'.$$

Hence, as the section of a mass of fluid decreases, its velocity increases in the same proportion. For instance, the stream of a river is more rapid at places where the width of the river is diminished. This also accounts for the gradual contraction of the descending jet of liquid, Art. (172), for the velocity increases, and therefore the section diminishes.

181. *A cylindrical vessel containing liquid has a small orifice in its base; to find the velocity at the surface.*

If the orifice be small and the surface large, the surface will descend very slowly.

Let h be the height of the surface, then $\sqrt{2gh}$ is approximately the velocity at the orifice. Take K for the area of the base of the vessel, and k of the orifice.

Then, neglecting the change of velocity at the orifice in the unit of time, $\sqrt{2gh}$ is the quantity of liquid which passes through the orifice, and therefore if V be the velocity at the surface,

$$VK = k\sqrt{2gh}.$$

If the vessel be kept constantly full, the motion is steady and the velocities are constant: hence the time in which a quantity of liquid, equal in volume to the cylinder, would, under these circumstances, flow through the

$$\text{orifice} = \frac{h}{V} = \frac{K}{k} \sqrt{\frac{h}{2g}}.$$

It will be seen that this is only a rough approximation to the actual facts of the case, but its insertion will serve to illustrate the laws above mentioned.

182. *Pressure of air in motion.* Early in the 18th century Hawksbee observed that if a current of air be transmitted through a small box the air becomes rarefied. This fact is illustrated by the following experiment.

Take a small straight tube, and at one end of it fix three smooth wires parallel to the tube and projecting from its edge, and let a flat disc be moveable on these wires, with its plane perpendicular to the axis of the tube. Blow steadily into the other end, and it will be found that the disc will not be blown off, but will oscillate about a point at a short distance from the end of the tube.

The reason of this apparent paradox is that the diminution of the density of the air in motion diminishes the pressure on the disc which would otherwise result from the continued action of the air impinging upon it, and the result is that it is balanced by the atmospheric pressure on the other side.

A full account of this experiment, and of other facts connected with it, was given by Professor Willis in the third volume of the Cambridge Philosophical Transactions.

A similar experiment was performed, in 1826, by M. Hachette, with a stream of water, and it was found that the pressure of the water was diminished by an increase of velocity.

It is worthy of remark that large ships of the Devastation class are observed to sink more deeply in the water when their speed is increased.

In a series of papers in *Nature*, for November and December 1875, Mr W. Froude has given a number of experimental illustrations, with explanations in general terms, of the connection between the pressure and the kinetic energy of a liquid.

Impulsive Action.

183. Imagine a closed vessel filled with liquid and having an aperture in its surface fitted with a piston. Let an impulse be applied to this piston; then assuming the incompressibility of the liquid, it can be shewn by the same reasoning as for finite pressures, that the impulse is transmitted throughout the mass, and is, at any point, the same in every direction.

The impulse at any point is measured in the same manner as a finite pressure; that is, if ϖ be the impulsive pressure at a point, ϖk is the impulse on a small area k containing the point.

A cylindrical vessel, containing liquid, is descending with a given velocity and is suddenly stopped; to find the impulsive action at any point.

The impulsive pressure at all points of the same horizontal plane will be the same, and if ϖ be the pressure at a depth x , and k the area of the base of the cylinder, ϖk is the impulse between the portion of the liquid above and below the plane at a depth x , and this impulse evidently destroys, and is therefore equal to, the momentum of the liquid mass above, which is $\rho k x v$.

Hence $\varpi k = \rho k x v$,
and $\therefore \varpi = \rho v x$.

If a vessel of any shape, containing liquid, descend vertically and be suddenly stopped, we can prove, by considering a small vertical prism of liquid, that the impulse at any point varies as the depth below the surface of the liquid.

This being the case, it follows that the propositions relating to whole pressure, and to resultant vertical and horizontal pressures in Chapters III. and IV., are equally true of impulsive pressures for the particular case in which the motion destroyed is vertical. The question is really the same if the vessel be made to ascend suddenly from rest, or have its velocity suddenly changed.

Ex. In a closed vessel of liquid a ball of metal is suspended by a vertical string fastened to the upper part of the vessel. Find the impulsive tension of the string when the vessel is suddenly raised with a given velocity.

The resultant impulse of the liquid on the ball will be the same as if its place were occupied by the liquid, and therefore will be equal to the momentum of the ball of liquid.

If U be the volume, and v the velocity, this is ρvU . But if ρ' be the density of the metal, the momentum of the ball is $\rho'vU$, and this is produced by the impulse of the liquid, and the tension T of the string.

Hence $\rho'vU = \rho vU + T,$
and $T = (\rho' - \rho) vU.$

CHAPTER XIII.

On Sound.

184. **T**HE sensation which we call sound is produced by a vibratory movement of the atmosphere; however it is first caused, it finally affects the organs of hearing by means of the air. A blow struck on any elastic body will produce sound, and the more highly elastic the body is the more easily will the sound be produced; a piece of metal when struck will ring sharply while the same blow on a piece of wood produces a dull sound of less intensity. A sound may traverse intervening bodies and be finally imparted through air which has no direct communication with the air in which it originated.

The fact that *air is necessary for the transmission of sound* may be shewn experimentally. Suspend a bell within the receiver of an air-pump, and provide a means of striking the bell from without, for instance, by a rod sliding in an air-tight collar. Then proceed to exhaust the receiver, and it will be found that as the exhaustion progresses, the sound of the bell becomes fainter, and is finally lost altogether.

That there is an actual motion in the particles of air is shewn by the transmission of sound through solid bodies, and also by the fact that a musical note sounded on any instrument will sometimes produce a sound, in unison with it, from some other body not in contact with the instrument.

Velocity of Sound.

The rate at which sound travels depends on the temperature of the atmosphere; it has been found experi-

mentally that at the freezing temperature the velocity is about 1089 feet per second, and that at a temperature of 61° F., when the height of the barometer is 29.8 inches, the velocity is nearly 1118 feet per second. We may therefore take 1100 feet per second as the velocity of sound under average atmospheric conditions.

Distance of a sounding body. Knowing the velocity of sound, we can estimate the distance of a sounding body whenever the production of sound is accompanied by the production of light. The velocity of light is so great that its transmission through all ordinary distances on the earth may be considered instantaneous, and thus if a cannon be fired from a ship at sea, the interval between seeing the flash and hearing the report will determine the distance of the ship. In the same manner the interval between a flash of lightning and the thunder which follows it will determine the distance of the cloud from which the flash is evolved.

The rolling of thunder may be accounted for in two ways. A single explosion may accompany the lightning, in which case a peal of thunder will be due to the reflection of the sound by clouds in different directions, and will be in fact a succession of echoes. Or the electric flash may pass rapidly from cloud to cloud, and thus the sounds of a series of explosions taking place almost at the same instant, but at different distances from the spectator, will arrive in succession and produce a continuous peal. In this latter case the peal is probably intensified and lengthened by echoes.

Velocity of sound through water. Sound is transmitted with much greater velocity through water, and through highly elastic solids, than through air. By experiments made in the lake of Geneva, the velocity was found to be 4708 feet per second, when the temperature of the water was 8° C. The rate of transmission through metallic substances is very much greater.

Velocity through gases. We have stated that the velocity in air depends on the temperature, and not on the density. In fact it depends on the value of k ,

which is different for different gases, and therefore the velocities in gases differ from each other. For instance, the velocity in hydrogen is nearly four times that in air at the same temperature, the elasticity of hydrogen being much greater than that of air.

Transmission through the atmosphere. The various portions of the atmosphere through which a sound passes, may have different temperatures, and consequently the sound will travel with a variable velocity. Moreover, the passage through varying strata tends to disturb the vibrations and to diminish the intensity. This accounts for the fact that distant sounds are heard more distinctly at night than during the day, the atmosphere being in general more quiescent, and having a more equable temperature.

Sound Waves.


185. A *wave* is the term applied to any state of vibratory motion which is transmitted progressively through the particles of a body. The effect of dropping a stone in still water is a familiar illustration; the rise and fall of the water produced by the plunge of the stone travels outwards in an expanding circle, while the particles of water merely rise and fall in succession as the wave passes over them.

Thus a portion of the atmosphere being in any way set in motion, the vibrations are communicated to the surrounding air, and the expanding spherical wave impinging on the ear produces the sensation of sound.

The intensity of a sound diminishes as the distance of the sounding body is increased. As a spherical wave expands, its thickness remaining constant, the vibrations are communicated to larger masses of air, and, in accordance with a general law of mechanics, the intensities of the vibrations are diminished. The intensity in fact is diminished in the inverse ratio of the square of the distance. This law however does not hold, if the sound be transmitted through tubes or pipes. In such cases the intensity is very slowly diminished.

Propagation of a Wave along a Straight Tube.

186. Consider a straight tube filled with air, and let a disc AB at one end oscillate rapidly over the space aa' .



When the disc oscillates from A to a , it compresses the air before it, and when the disc is at a , the compression has traversed and extends over the space AC . This compression travels along the tube with a constant velocity, and is called the condensing wave.

As the disc returns from a to A , it rarefies the air behind it; and this rarefaction extends over AC , while the previous compression has been transferred to the space CD , and thus a rarefying wave follows the condensing wave.

As the disc moves from A to a' , another rarefying wave is produced, and when the disc returns to A , a condensing wave is produced, while during these two processes the first condensing and rarefying waves have been transferred to EF and DE respectively.

The disc having its greatest velocity at A , and coming to rest at a and a' , it is obvious that the condensation is greatest at F , and diminishes gradually to E , where there is no condensation, or where the density is the same as if the air were at rest; from E to D the air is rarefied, and at D the rarefaction is greatest; from D to C the rarefaction decreases, and at C condensation commences and increases to A .

Thus a complete wave or undulation is formed, and if the disc oscillate once only, a single wave will travel along

the tube taking successive positions as in the figure; if the disc continue to vibrate, a succession of these waves will be produced and will follow each other continuously along the tube. If these waves, on emergence from the tube, impinge on the ear, the sensation produced will be that of a continuous and uniform sound.


The vibrations can be produced without the aid of the disc, as, for instance, by blowing across the end of the tube.

It will be observed that the velocities of the vibrating particles of air are zero at F and D , and greatest at E and C .

The length of a wave is the distance between any two points at which the *phases* of vibration are the same, that is, at which the velocities of the vibrating particles are the same in direction and magnitude.

Motion of a Wave along a Stretched String.

187. In a similar manner, if a portion of a stretched cord PQ be set in motion, a wave, or succession of waves,



will travel along the cord, and on arriving at Q will be reflected and travel back again.

The string may vibrate somewhat in the form of the curve $ABCDE$, AE being the length of a wave, B and C the points at which the displacement is greatest, and the velocity zero, and A , C , and E the points at which the displacement is zero and the velocity greatest.

In this case the vibration is perpendicular to the line in which the wave travels, but its analogy with the case of the tube is sufficiently evident.

The vibrations of the string are communicated to the air and thereby conveyed to the ear.

Musical Sounds.

188. Any series of waves, following in close succession, may produce a continued sound; if they are irregular in magnitude, the result is a *noise*, but a *musical note* is produced by a constant succession of equal waves.

Pitch, intensity, and quality. Notes may differ from each other in three characteristics; thus, a note may be grave or acute, that is, its *pitch* may be high or low; and the pitch of a note depends on the length of the constituent wave, and is higher as the length of the wave is less. The *intensity* of a sound depends on the extent of vibration of the particles of air, and its *quality* is a characteristic by which notes of the same pitch and intensity are distinguished from each other. The quality of a note, or, as it is sometimes called, its *timbre*, depends on the nature of the instrument from which it is produced.

A further distinction of sounds is sometimes marked by the word *tone*. Thus the tone of a flute differs from the tone of other instruments, while two flutes may, and will in general, produce sounds which differ in quality.

189. *Sounds of different pitch travel with the same velocity.* This appears to be the case from the fact that if a musical band be heard at a distance there is no loss of harmony, and therefore there can be no sensible difference in the velocities of the different sounds.

190. *Reflection of waves in a tube of finite length.* It is found both by experiment and theory that a wave on arriving at the end of a tube is reflected, whether the end be open or closed, and travels back again, changed only in intensity, to be again reflected at the other end.

This accounts for the *resonance* in a tube when the air within it has been set in vibration.

191. *Coexistence and interference of undulations.*

Different sound waves travelling through the air traverse each other without alteration either of pitch or in-


tensity. In other words, different undulations coexist without affecting each other, and the actual vibration of any particle of air is the sum or difference of the coexistent vibrations which are at the same instant traversing the particle of air.

A simple illustration of this coexistence may be seen by dropping two stones in water. The expanding circular waves intersect, and at the points of intersection it will be seen that a depression and an elevation neutralize each other, and that two depressions or two elevations at the same point increase the amount of one or the other. If there be a sufficient number of circular waves the points of greatest elevation will be seen to lie in regular curves, as also those of depression, and of neutralization*. The vibrations in this case being transverse to the direction of transmission of the wave are different from those of sound waves, which are longitudinal or in the direction of transmission, but the effect of coexistence is the same in all cases.

The effect of coexistence in producing neutralization, or increase of intensity, is called the interference of undulations, and it must be observed that, while two sets of undulations are physically independent of each other, their geometrical resultant may be a form of undulation different from that of either component, as in the case just referred to of the undulations in the surface of water.

* These curves are hyperbolas, for, if A and B be the centres of disturbance, and P, P' the points of intersection of two particular waves, AP and BP increase uniformly with the time, and the rate of increase of each is the same.

Hence, their difference is constant, and the locus of P is an hyperbola of which A and B are the foci. As other waves follow in succession the series of such points will lie in confocal hyperbolas.



The Notes which can be produced from a Tube closed at one end.

192. When a definite note is being sounded from a tube, the air within the tube vibrates regularly, every particle maintaining the same vibration, and there are certain points of the tube at which the air remains at rest. These points, or planes of division of the tube, are called *nodes*, and the planes of maximum vibration are called *loops*.

The motion in fact is the same as if there were fixed waves in the tube, and the nodes and loops are the points of zero velocity and zero condensation.

The motion thus described is called *steady* motion, and its existence is requisite to the continuance of a definite note.

In the case of a tube closed at one end B , it is clear that the end B must be a node, and since the end A is open its density is sensibly that of the air outside, and we may take it to be a loop.

It is therefore evident that the longest possible wave for which the motion can be steady is four times the length of AB ; and the corresponding sound is the fundamental note of the tube.

Further, AB may be any odd multiple of the distance from a node to a loop, and if $AB=l$, and λ be the length of a wave, we must have

$$l = (2n + 1) \frac{\lambda}{4},$$

$$\text{or } \lambda = \frac{4l}{2n + 1}.$$

Hence the notes which can be produced from AB have for their wave lengths,

$$4l, \frac{4l}{3}, \frac{4l}{5}, \text{ \&c.,}$$

and, if v be the velocity with which a wave traverses the tube, the times of vibration are

$$\frac{4l}{v}, \frac{4l}{3v}, \frac{4l}{5v}, \dots$$

and are therefore in the ratio of the fractions

$$1, \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \text{ \&c.}$$

The Notes of a Tube open at both ends.

193. In this case each end is a loop, and there is therefore a node between; hence the greatest possible wave length is twice the length of the tube, and further the length of the tube must be some multiple of half the length of a wave.

Hence
$$l = m \frac{\lambda}{2}, \text{ and } \lambda = \frac{2l}{m}.$$

The successive waves are therefore

$$2l, l, \frac{2l}{3}, \frac{l}{2}, \frac{2l}{5}, \text{ \&c.},$$

and the vibrations in the ratio of the fractions

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$$


It will be observed that the fundamental note of the open tube is an octave higher than that of the closed tube, the wave length for the former being half that for the latter.

The Formation of Nodes and Loops.

194. Taking the case of a tube HK closed at the end K , the aerial particles at the end K are permanently at rest, while those at A are in a state of permanent vibration. We have stated, as an experimental fact, that a series of waves travelling along HK in regular succession are reflected at K and travel in


the opposite direction; and this fact enables us to account for the existence of nodes and loops.

In order to give the required explanation we must first explain a method of representing geometrically the state of motion of the aerial particles in a wave.



Take AE as a wave length, and let the ordinates of the curve $ABCDE$ represent the velocities of the several particles parallel to the line AE ; thus NP represents the velocity of the particle at N , NP being drawn upwards when the velocity is in the direction AE , and downwards when the velocity is in the opposite direction.

Hence, if two distinct sets of vibrations coexist along a line of aerial particles, we can determine the resultant motion by drawing the two curves for the two waves, and the algebraic sum of the ordinates at any point will represent the resultant velocity at that point.



Imagine now a wave travelling along AB , and impinging on the fixed end K ; this wave will be reflected and will travel along BA with reversed velocities.

In the figure the dotted line will represent the reflected wave, and the effect of the reflected wave is the same as that of a wave $B'C'$ travelling in the direction KA .

It will be seen from the figure that the velocities at K from the two waves are always equal and opposite, and that the resultant velocity at K is always zero, in accordance with the given conditions. In other words, the effect of the fixed end K is replaced by the effect of a reversed wave travelling in the opposite direction.

It will also be seen that there is a succession of points, K, L, M, \dots at which the velocity is always zero, and a succession K', L', \dots at which the velocity varies between its greatest values in both directions, the former set of points being nodes, and the latter loops.

Let a dotted line $KPLQMR$ be drawn such that its ordinate at any point is the algebraic sum of the ordinates corresponding to the incident and reflected wave; this dotted line will represent the state of vibration of the air in the tube at the instant considered, and it will be observed that while the points K, L, M, \dots are points of permanent rest, all the intermediate points represent the positions of aerial particles which vibrate steadily, their velocities being zero at regular intervals.

Thus, the opposing waves may be so placed that their extremities C, C' may coincide at K' ; in the figure this will occur when the incident waves have traversed the space CK' , and the opposing waves the space $C'K'$, and at this instant the velocity at K' will be zero. Subsequently the two waves travelling in opposite directions will produce at K' a velocity double that of either, so that the velocity at K' will then be a maximum, the interval of time being that during which the vibration has traversed a space equal to one-fourth of a wave length.

It will be now clear that, if a permanent vibration be maintained at the open end H , a succession of nodes and loops will necessarily be formed in the tube, provided that the wave length emitted from the end H is such as to satisfy the condition of Art. (192). This condition is that the wave length should be an odd submultiple of 4 times the length of the tube.

In a similar manner, if KH be a tube open at both ends, it is found that a wave or a set of waves travelling along HK are reflected at K , and traverse the tube in the direction HK . An important difference however exists between the two cases; in the former case the end K is a node, in the present case it is a loop, the particles of air vibrating freely, and the density being the same (very nearly) as that of the external air.



An analogous explanation will account in this case also for the formation of nodes and loops.

In the case of the tube closed at K , the reflected wave on arriving at H is partly emitted into the open air and partly reflected, thereby reinforcing the new vibration which is at the instant being excited at H , and aiding to produce another series of waves which are again reflected at K .

The effect which is thus produced on the ear is that of a sustained note, the character of which depends on the material of which the tube is formed, while its pitch and intensity depend solely on the lengths of the constituent waves and the extent of the vibrations of the aerial particles.

The Notes produced by a Vibrating Cord.

195. A stretched cord in a state of vibration may either oscillate as a whole, fig. (1), or in parts, as in figures



(2) and (3), the curved lines representing the actual positions at certain instants of the cord itself.

In any case the two ends are points of zero velocity or nodes, and the wave corresponding to the fundamental note has twice the length of the cord for its length on the cord.

In general, if the wave length on the cord be λ' , the length l of the cord must be some multiple of $\frac{1}{2}\lambda'$,

$$\text{i. e. } l = m \frac{\lambda'}{2}.$$

The velocity of propagation along the cord will depend on its tension, thickness and density; and if v' be this velocity the time of vibration is $\frac{\lambda'}{v'}$.

The pitch of the note produced is determined by the time of vibration, and therefore, if λ be the wave length produced in air by the vibrations of the cord and thereby

conveyed to the ear as a sound, and v the velocity of propagation in air, we shall obtain the note by the relation

$$\frac{\lambda}{v} = \frac{\lambda'}{v'}$$

since the aerial vibrations are performed in the same time as those of the cord.

Hence
$$\frac{\lambda}{v} = \frac{2l}{mv'}$$

and the wave lengths are

$$2l \frac{v}{v'}, \quad l \frac{v}{v'}, \quad \frac{2l}{3} \frac{v}{v'}, \quad \frac{l}{2} \frac{v}{v'}, \quad \&c.$$

These wave lengths give the series of harmonics producible from the cord, and it should be observed that any one may be produced alone, or any number of them may exist simultaneously.

196. *Vibration of rods.* We know that sounds are produced by vibrating rods, and we can determine the series of notes producible in any simple case by the considerations of the preceding articles. A rod fixed at one end and free at the other, will have for its fundamental note a wave length four times its own length, the fixed end corresponding to a node and the free end to a loop.

The analogy between a vibrating rod and a vibrating column of air will be now seen, but attention must be paid to the fact that the vibrations of air which produce sound are longitudinal, while the vibrations of a string are transversal, and those of a rod may be either transversal or longitudinal.

A common instance occurs in the humming of a telegraph-post, which is probably due to a series of longitudinal vibrations traversing the post in a vertical direction.

The transmission of sound through water is analogous to the transmission of sound by means of longitudinal vibrations along a rod, and is treated theoretically in exactly the same manner.

197. *The pitch of a note produced by a vibrating cord*

depends on the tension and substance of the cord, and is heightened by an increase of tension; and in a similar manner the pitch of a note produced from a rod is found to depend on its size and substance.

This is due to the fact that the rates of propagation of vibrations depend on the characteristics above mentioned, and thus a long wave length, traversing a cord or a rod very rapidly, may give rise to a short wave length in the aerial vibrations which result from those of the cord or the rod, and a high-pitched note be produced.

Unison and Harmony of Musical Notes.

198. Two notes are said to be in *unison* when the times of vibration, or the wave lengths, are the same for both.

The *harmony* of two notes consists in the recurrent coincidence, at short intervals, of their constituent vibrations; thus, if a note and its octave be sounded, the vibration belonging to the fundamental note coincides exactly with two vibrations of the octave, and the two sounds are said to be in harmony with each other.

More generally two notes are in harmony when a small number of vibrations belonging to one of them coincides exactly, in time, with a small number of the vibrations belonging to the other. An instance of this is the harmony of a note with its fifth in the diatonic scale, three vibrations of the upper note being coincident with two of the lower note.

199. *Communication of vibrations.* If two different bodies can vibrate in unison or in harmony with each other, that is, if their fundamental notes are either in unison or in harmony, it is a known fact that when one is set vibrating, the other, if not too far off, will vibrate also. The reason is that the sound waves diverging from one body impinge on the other, and when the vibrations of the latter can be in harmony with those of the former, the slight vibration at first established is maintained and intensified by the continued impulses of the same aerial vibration.

Thus a person singing or whistling in a room may sometimes hear notes sounding from thin glass jars or metallic tubes, and these notes will always be in harmony with the note originally sounded.

200. *Beats.* When a note of a pianoforte is sounded a series of alternations is generally to be noticed in the intensity of the sound, these alternations, which are called *beats*, occurring at regular intervals.

This phenomenon depends on the fact that there are in general two strings to each note, which are intended to be exactly in unison with each other. Practically the unison is seldom perfect, and hence the two sets of waves do not exactly coincide with each other.

The intensities are however very nearly the same, and hence, when the vibrations of the two waves oppose each other, a diminution of the intensity results, but when they are in the same direction the intensity is increased.

Suppose that τ and τ' are the times of vibration of the two notes; then if x vibrations of one coincide with $x+1$ of the other, we have

$$\tau x = \tau'(x+1),$$

or

$$x = \frac{\tau'}{\tau - \tau'};$$

and $\therefore \frac{\tau\tau'}{\tau - \tau'}$ is the interval between the instants of time at which the vibrations oppose each other, and is therefore the period of the beats.

It is evident that the more nearly τ and τ' are equal to each other, the longer is the period of the beats, and the less the number of beats heard while the sound is perceptible.

Beats are also produced when two notes are very nearly in harmony with each other; the explanation is the same as for the simple beats above mentioned.

Tartini's Beats. Again, when two notes are actually in concord, a note is sometimes heard in addition to the two notes, and of lower pitch than either. The vibrations of the two notes coincide at regular intervals; these coin-

cidences are Tartini's beats, and the effect of a series of such beats, at regular and rapidly recurring intervals, is that of a note which is grave in comparison with the original notes. This lower note is called a *subharmonic* of the two notes by which it is produced.

Notes.

Velocity of sound. A calculation from theoretical principles of this velocity was made by Newton, and again by Lagrange; the result obtained was about 916 feet per second.

This notable discrepancy between fact and theory remained unexplained until Laplace remarked that the heat developed by the sudden compression of the air would increase the elasticity, and therefore increase the calculated velocity.

New calculations were made, and the result is in complete accordance with fact.

The theoretical expression is $\sqrt{k\beta(1+at)}$, where β is the coefficient introduced by the consideration of the heat developed.

Intensity of sound after traversing pipes. Experiments were made by Biot with some water-pipes in Paris, and it was found that a whispered conversation could be carried on through a pipe 3000 feet in length.

The use of speaking-tubes in large houses is another illustration of the fact mentioned in Art. (185).

Vibrating Cords. It is found that the velocity with which a wave traverses a stretched cord is the same as the velocity which would be acquired by a heavy body falling through a vertical space equal to half a length of the cord of which the weight is equal to its tension. In other words, if the weight of a length l of the cord be equal to its tension, the velocity with which a wave travels along it is \sqrt{gl} .

The existence of nodes and loops in the case of a cord may be practically manifested by placing on the vibrating cord small pieces of paper, cut so as to rest on the cord; those which are placed at the nodes will remain on the cord, while those which are placed near the loops will be thrown off.


The *Monochord* is a simple instrument for trying the experiments just mentioned, and for testing other results of theory.

A cord fastened at one end is stretched over a sounding board, and passing over a bridge is tightened by a weight at the other end; the tension may be varied by changing the weight, and by means of another bridge, moveable along the board, the length of the vibrating portion may be diminished. The notes obtained for different lengths and different tensions can be thus compared, and the wave lengths for different notes can be directly measured.

Practical illustration of the interference of aerial vibrations. From Art. (191) we can see that if two waves, exactly similar to each other, travel in the same direction, and one be half a wave length behind or before the other, the result will be a permanent quiescence of the aerial particles along the direction in which the waves travel.

This has been shewn visibly by an experiment, which is due to Mr Hopkins.

A straight tube AB branches off at the end B into two portions BC, BD ; the end A is closed by a tight membrane and fine sand is scattered over the membrane. A vibrating plate of glass is placed beneath C and D so that the two portions immediately beneath C and D shall be in opposite phases of vibration. The waves thus produced in CB and DB traverse these branches of the tube, and arrive at B in opposite phases, that is, one is the half of a wave length before the other, and therefore there is theoretically no resultant vibration in BA . Practically it is found that the sand on A is undisturbed, but, if the plate be turned round, the sand is immediately thrown into a state of violent commotion.



Beats. The theory of beats is given in Smith's *Harmonics*, published in 1749. Tartini's treatise, in which the sounds called by his name were first discussed, appeared in 1754.

The diatonic scale. The ordinary or diatonic scale consists of a series of notes, for which the times of vibration are in the ratio of the numbers in the following table:

C	D	E	F	G	A	B	C
1,	$\frac{8}{9}$,	$\frac{4}{5}$,	$\frac{3}{4}$,	$\frac{2}{3}$,	$\frac{3}{5}$,	$\frac{8}{15}$,	$\frac{1}{2}$;

or, in other words, the numbers of vibrations per second are in the ratio of

$$1, \frac{9}{8}, \frac{5}{4}, \frac{4}{3}, \frac{3}{2}, \frac{5}{3}, \frac{15}{8}, 2,$$

that is, of the numbers

$$24, 27, 30, 32, 36, 40, 45, 48.$$

As a matter of fact the actual number of vibrations corresponding to the particular *C* employed as a central note varies in different places, and from time to time. As an ordinary standard for the concert pitch of this note *C*, about 128 vibrations in a second is taken as making the note*, and the numbers of vibrations for the several notes of the scale are then respectively

$$128, 144, 160, 170, 192, 214, 240.$$

The range of sounds appreciable by the human ear varies for different persons, but in general extends over above nine octaves. A series of aerial impulses will produce the impression of a continuous note when they recur with such rapidity that the ear cannot appreciate the succession of impulses, and it is found that this is the case for a wave length of about 68 feet. On the other hand it has been found that the highest note which is in general appreciable has about eight-fifths of an inch for its wave length.

* See Spencer's *Treatise on Music*, in Weale's Series.

APPENDIX I.

The equilibrium of fluids under the action of any given forces.

201. In any field of force the measure of the force at any point is the force which would be exerted upon the unit of mass supposed to be concentrated at that point.

As in Art. (10), it can be shown that the pressure at any point is the same in all directions; for if we consider the equilibrium of a very small prism, the forces at all points of the prism will be ultimately equal and parallel, and the case then becomes the same as that of a prism under the action of gravity.

202. *The measure of the force at a point, in a given direction, multiplied by the density, is equal to the rate of change, per unit of length, of the pressure in that direction.*

If P be the point, take any length PQ in the direction considered and describe a very thin cylinder about PQ .

The equilibrium of this cylinder is maintained by the pressures on its ends and on its curved surface and by the external forces in action.

Therefore the difference of the pressures on the ends P and Q is equal to the force on the cylinder in the direction PQ , and, if a be the cross section, and PQ be very small, so that its density may be considered uniform, and the measure of force, f , the same at all points of PQ , we have, taking p and p' the pressures,

$$(p' - p)a = \rho a f PQ,$$

so that
$$\rho f = \frac{p' - p}{PQ},$$

which is the rate of change of pressure.

203. DEF. Surfaces of equal pressure are surfaces in the fluid over which the pressure is constant.

Surfaces of equal pressure are at every point perpendicular to the resulting force.

To prove this, consider two consecutive surfaces of equal pressure containing between them a stratum of fluid, and let a small circle be described about a point P in one surface, and a portion of the fluid cut out by normals through its circumference.

This small cylinder of fluid is kept at rest by the external force and by the pressures on its ends and on its circumference.

The pressures at all points of the circumference being equal, the pressures on the two forces must be counter-balanced by the external force, which must therefore act in the direction of these pressures, i.e. perpendicular to the surface of equal pressure.

Again, if d be the distance at P between the consecutive surfaces, we have, as before,

$$\rho f d = p' - p,$$

so that


$$\rho d \propto \frac{1}{f},$$

and, in the case of a homogeneous liquid,

$$d \propto \frac{1}{f}.$$

204. If in any field of force a particle be in contact with a smooth surface, it will be in equilibrium if the normal to the surface coincide with the direction of the resultant force.

Surfaces of equilibrium are therefore at all points perpendicular to the resultant force.



If a particle be moved over a surface of equilibrium no work is done against the force, and these surfaces are therefore surfaces of equal energy, or equipotential surfaces.

If a particle of mass unit be carried along the normal from one surface to another the work done is $f \cdot PQ$, which is the change of energy and is constant ;


$$\therefore f \cdot PQ \text{ is constant.}$$

Surfaces of equal pressure are also surfaces of equal density ;

For ρfd is constant and we have just shewn that fd is constant, $\therefore \rho$ is constant.

205. Examples. (1) *A mass of liquid at rest under the action of a force to a fixed point varying as the distance from that point.*

The surfaces of equilibrium, and therefore of equal pressure, are clearly concentric spheres, and the free surface is a sphere.



To find the pressure at any point P , take a thin cylindrical column from P to the surface and observe that its equilibrium is maintained by the pressure at the end P counterbalancing the attractive force.

If a be the cross section, $OP = r$, $OA = a$, and if μr be the force at the distance r ,

$$pa = \text{force on the column } AP$$

$$= \rho a (a - r) \mu \frac{1}{2} (a + r), \text{ by Leibnitz's theorem ;}$$

$$\therefore p = \frac{1}{2} \mu \rho (a^2 - r^2).$$

The Pressure on a diametral plane

$$= \text{Force on a hemisphere}$$

$$= \frac{2}{3} \rho \pi a^3 \cdot \mu \cdot \frac{3a}{8} = \frac{1}{4} \mu \rho \pi a^4.$$

(2) *Liquid at rest under the action of forces to any number of centres varying as the distance.*

The resulting force is directed to a fixed point and varies as the distance from that point; this case is therefore the same as the preceding.

(3) *Liquid at rest under the attraction of a straight rod, the molecules of which attract with force varying inversely as the square of the distance.*

If AB be the rod, it can be shewn by elementary geometry that the direction of the resulting attraction at any point P bisects the angle APB ; from this it follows that the surfaces of equal pressure are confocal spheroids, having their foci at A and B .

APPENDIX II.

WE shall now conclude with the solution of some problems of a more extended character than those which have been hitherto discussed; to these the student will find that the principles of the preceding pages are directly applicable, but that a larger demand than before will be made upon his skill in algebraic operations.

(1) *Centre of Pressure.* A general expression can be obtained for the depth of the centre of pressure of any plane area.

Let the area be divided by horizontal lines into a number of very small portions, and let a be the area of one of these portions and z its depth below the surface.

Then the pressure upon it = $gpza$, and if \bar{z} be the depth of the centre of pressure, we have by the usual formula for the centre of a system of parallel forces,

$$\bar{z} = \frac{\sum gpza \cdot z}{\sum gpza} = \frac{\sum (z^2 a)}{\sum (za)},$$

$gp \sum (za)$ being the pressure on the whole area.

Ex. An isosceles triangle is immersed vertically, its base being horizontal and its vertex A at a depth c below the surface.

Let $AD = h$,

$$AN = r \frac{h}{n}, \text{ and } NM = \frac{h}{n},$$

the line AD being divided into n equal portions.


Then $PP' = 2 \frac{rh}{n} \tan \frac{A}{2}$, and $z = c + \frac{rh}{n}$,

$$\sum (z^2 a) = \sum \left(c + \frac{rh}{n} \right)^2 2r \frac{h^2}{n^2} \tan \frac{A}{2}.$$

$$\tan \theta = \frac{\pi}{2} \tan \alpha,$$

where θ is the inclination of RS to the horizon.

S is therefore the centre of pressure.



To find its position, we have

$$\tan \theta = \frac{RM}{SM} = \frac{RV - MV}{MV \cdot \tan \alpha};$$

$$\therefore \frac{RV}{MV} - 1 = \frac{\pi}{2} \tan^2 \alpha;$$

$$\therefore MV = \frac{RV}{1 + \frac{\pi}{2} \tan^2 \alpha}.$$


But
$$RV = \frac{1}{2} KV = \frac{1}{2} EV \sec \alpha;$$

$$\therefore SV = MV \sec \alpha = \frac{\frac{1}{2} EV \sec^2 \alpha}{1 + \frac{\pi}{2} \tan^2 \alpha}.$$

(3) One asymptote of an hyperbola lies in the surface of a fluid; it is required to find the depth of the centre of pressure of the area included between the immersed asymptote, the curve, and two given horizontal lines in the plane of the hyperbola.

Taking OA, OB as the axes, let $PN, P'N'$ be two lines near each other and parallel to OA .

The pressure on the small area PN'
 $= g\rho ON \sin \omega . PN . NN'$.



But $ON . PN \sin \omega$ is the area of the parallelogram $OMP N$, the constancy of which is a known property of the hyperbola.

Hence the pressure on PN' varies as its vertical thickness, and therefore the depth of the centre of pressure of any finite area contained between two horizontal lines, the curve and the asymptote, is half the sum of the depths of the horizontal lines.

(4) *Having given the position of the centre of pressure of a plane area increased vertically at a given depth, it is required to find its position when the area is immersed in the same position to any other given depth.*

Let K be the position of the centre of pressure when G the centre of gravity is at the depth h .

If the depth be increased to h' , the increase of pressure on the area A is $wA(h' - h)$ acting at G .


Take the point K' in GK such that

$$wAh' . GK' = wAh . GK, \text{ or } GK' = GK \frac{h}{h'};$$

then K' is the new centre of pressure.

(5) *A triangular area is immersed with one angular point in the surface; it is required to find its centre of pressure.*

Dividing the base BC into a large number of equal parts, the centre of pressure of an elementary triangle AP will be at a point R such that $AR = \frac{3}{4} AP$, P being the middle point of the base of the elementary triangle.



If $AE = \frac{3}{4} AB$, the centre of pressure, K , of ABC will be on the line EF parallel to BC .

Further, all the elementary triangles being equal, the pressure on AP will be proportional to the depth of its centre of gravity, and therefore will vary as RG .

Hence it follows that K is the same as the centre of gravity of the frustum EF of a triangle, vertex G , and

$$\therefore GK (GE^2 - GF^2) = \frac{2}{3} (GE^3 - GF^3)$$

$$\text{or } GK = \frac{2}{3} \cdot \frac{GE^2 + GE \cdot EF + GF^2}{GE + GF} = \frac{1}{2} \cdot \frac{BD^2 + BD \cdot CD + CD^2}{BD + CD}.$$

If β, γ be the depths of B and C ,

$$\text{the depth of } K = \frac{1}{2} \frac{\beta^2 + \beta\gamma + \gamma^2}{\beta + \gamma}.$$

(6) We can now by the aid of (4) find the depth, z , of the centre of pressure of a triangle ABC in terms of the depths α, β, γ of its angular points.

Draw a horizontal plane through A and remove the liquid above; then, if z' be the depth of the centre of pressure below A ,

$$z' = \frac{1}{2} \frac{(\beta - \alpha)^2 + (\beta - \alpha)(\gamma - \alpha) + (\gamma - \alpha)^2}{\beta + \gamma - 2\alpha}.$$

Replacing the liquid, and taking S for the area, we have a new pressure wSa at the centre of gravity, and therefore

$$wS \frac{a + \beta + \gamma}{3} z = wS \frac{\beta + \gamma - 2a}{3} (z' + a) + wSa \frac{a + \beta + \gamma}{3},$$

or $2z(a + \beta + \gamma) = a^2 + \beta^2 + \gamma^2 + \beta\gamma + \gamma\alpha + \alpha\beta.$

If h, k, l be the depths of the middle points of the sides of the triangle,

$$z(h + k + l) = h^2 + k^2 + l^2.$$

(7) A similar method may be employed to find the centre of pressure of a sector of a circle with its centre in the surface.

Taking the case of a sector with one bounding radius (c) in the surface, divide the sector into a large number of small triangles; the centres of pressure of these triangles will be on the arc of a circle of radius $\frac{3}{4}c$, and it can be shewn, by the summation of a trigonometrical series, that the depth of the centre of pressure is

$$\frac{3c}{16} \frac{2a - \sin 2a}{1 - \cos a},$$

$2a$ being the angle of the sector.


(8) *A cylindrical vessel, open at the top, is inverted and pushed down vertically in water; the substance of the vessel being of greater density than water, it is required to prove that, at a certain depth, it will be in a position of equilibrium which for vertical displacements is unstable.*

As the vessel is forced downwards the pressure of the water compresses the air within, and there must be some depth at which the air will be so compressed that the weight of the water displaced by the vessel and the air is exactly equal to the weight of the vessel and air together. At this point there will be equilibrium; but, if the vessel be slightly lifted, the air within will expand, and the weight of water displaced will be too great for equilibrium; hence the vessel will ascend. If on the other hand it be slightly depressed, a further compression of the air will take place, and the vessel will then descend.

(9) *A square lamina floats with its plane vertical and one angular point below the surface; it is required to find its positions of equilibrium.*

Let PQ be the surface of the liquid, G the centre of gravity

of the square, and H of the liquid displaced, E being the middle point of PQ .



Then, if $OP = x$, and $OQ = y$, and if ρ , σ be the densities of the liquid and the lamina, and $2a$ the side of the square,

$$\frac{1}{2} \rho xy = 4\sigma a^2, \quad \text{or } xy = 8 \frac{\sigma}{\rho} a^2 = c^2 \text{ suppose.}$$

We have now to express the condition that GH is vertical.

Draw HN perpendicular to OP ;

$$\text{Then} \quad ON = \frac{1}{3} x, \quad \text{and } HN = \frac{1}{3} y.$$

Hence, if GM , HL be perpendicular and parallel to OP , the tangent of the angle which HG makes with OP

$$= \frac{GL}{HL} = \frac{GM - HN}{OM - ON} = \frac{a - \frac{1}{3} y}{a - \frac{1}{3} x};$$

but this angle is the complement of OPQ , of which the co-tangent is $\frac{x}{y}$;

$$\therefore \frac{3a - y}{3a - x} = \frac{x}{y};$$

$$\text{or } x^2 - y^2 = 3a(x - y).$$

This equation gives $x=y$,
and $x+y=3a$.

The first result gives the symmetrical position of equilibrium, for which $x=y=c$.

From the second,

$$x + \frac{c^2}{x} = 3a,$$

$$\therefore x = \frac{3a}{2} \pm \sqrt{\frac{9a^2}{4} - c^2}.$$


Hence, if $\frac{9a^2}{4} > c^2$, i. e. if $\frac{\rho}{\sigma} > \frac{32}{9}$, there are two other positions of equilibrium.

If $\frac{\rho}{\sigma} = \frac{32}{9}$, it will be seen that these three positions coincide.

(10) *A vessel in the form of a paraboloid is immersed with its open end downwards, in a trough of mercury. Supposing the length of the axis of the vessel to be to the height of the barometer as 45 is to 64, it is required to find the depth of the surface of the mercury within the vessel when the whole vessel is just immersed.*

Let AM be the height of the vessel, and h the height of the barometer; then

$$AM = \frac{45}{64} h.$$



If PN be the surface of the mercury within the vessel, and Π' the pressure of the air within,

$$\frac{\Pi'}{\Pi} = \frac{\text{volume } AQM}{\text{volume } APN} = \frac{AM^2}{AN^2};$$

but $\Pi' = \Pi + g\sigma AN$, and $\Pi = g\sigma h$;

\therefore , if $AN = x$,

$$\frac{h+x}{h} = \left(\frac{45}{64}\right)^2 \frac{h^2}{x^2},$$

$$\text{or } x^3 + hx^2 = \left(\frac{45}{64}\right)^2 h^3.$$

Writing $\frac{z}{16}$ for $\frac{x}{h}$, this becomes

$$z^3 + 16z^2 = 45^2,$$

from which we find easily by trial $z=9$,

$$\text{and } \therefore AN = \frac{9}{16} h.$$

(11) A cylindrical vessel contains a given quantity of fluid. In this fluid is placed another cylindrical vessel of half the diameter of the first and containing half the quantity of fluid which is of half the specific gravity of that in the first vessel. In this second vessel is placed a third related to the second as the second is to the first; and so on indefinitely. Find the distance between the surfaces of the first and n^{th} fluids, neglecting the weights of the vessels.

Let $\rho, \frac{1}{2}\rho, \frac{1}{2^2}\rho$, &c. be the densities,

$r, \frac{1}{2}r, \frac{1}{2^2}r$, the radii, and

h_1, h_2, h_3 , the heights of fluid in the respective cylinders.

$$\text{Then } r^2 h_1 = 2 \left(\frac{r}{2}\right)^2 h_2 = 2^2 \left(\frac{r}{2^2}\right)^2 h_3 \dots = 2^{n-1} \left(\frac{r}{2^{n-1}}\right)^2 h_n;$$

$$\therefore h_n = 2^{n-1} h.$$

If $\pi r^2 h = V$, the whole weight of fluid in all the cylinders beginning with the second

$$= g\rho \left(\frac{1}{2} \frac{V}{2} + \frac{1}{2^2} \frac{V}{2^2} + \dots \text{ to infinity} \right)$$

$$= \frac{1}{3} g\rho V.$$

This whole weight is floating in the fluid of the first cylinder, and therefore if z be the depth immersed of the second cylinder,

$$g\rho \frac{\pi r^2 z}{4} = \frac{1}{3} g\rho V = \frac{1}{3} g\rho \pi r^2 h,$$

$$\text{whence } z = \frac{4}{3} h.$$

But the effect of this immersion is to raise the surface in the first cylinder to a certain height x such that

$$\pi r^2 x - \pi \frac{r^2 z}{4} = \pi r^2 h,$$

$$\text{or } x = \frac{4}{3} h.$$

The base of the second cylinder therefore just descends to the base of the first, and the same is the case with all the successive cylinders.


Hence the successive heights of the surfaces above the base are

$$\frac{4}{3} h, \frac{4}{3} 2h, \frac{4}{3} 2^2 h, \text{ \&c.}$$

and the required distance is

$$\frac{4}{3} h (2^{n-1} - 1).$$

(12) A straight tube $ABCD$ of small bore is bent at B and C so as to make ABC and BCD right angles, AB being equal to CD . The tube thus formed is moveable in a vertical plane about its centre of gravity, and being placed with BC horizontal



and downwards, water is poured in (at A or D) so that c is the length of BA or CD occupied by the fluid. It is required to determine the condition of stability.

Let $BC=2a$, and take b as the distance of G , the centre of gravity of the tube, from BC , and P, Q , as the surfaces of the water.

Turn the tube through a small angle θ so that P', Q' are the new surfaces, and therefore

$$PP' = QQ' = a \tan \theta.$$

If the moment of the weight of the water about G be in the direction opposite to the displacement, the equilibrium will be stable.

Taking κ as the area of a section of the tube, this moment

$$= g\rho\kappa \{2ab \sin \theta + (c - a \tan \theta) EN - (c + a \tan \theta) E'N'\},$$

E, E' being the middle points of $P'B, Q'C$; $EN, E'N'$ perpendiculars on the new vertical through G , and FL perpendicular to EN .

$$\text{But } EN = LN + BF \cos \theta - EB \sin \theta$$

$$= b \sin \theta + a \cos \theta - \frac{1}{2}(c - a \tan \theta) \sin \theta,$$

$$\text{and } E'N' = a \cos \theta + \frac{1}{2}(c + a \tan \theta) \sin \theta - b \sin \theta.$$

Hence, supposing θ very small, $\sin \theta = \theta$, $\cos \theta = 1$, and the moment

$$= g\rho\kappa \left\{ 2ab\theta + (c - a\theta) \left(b\theta + a - \frac{1}{2}c\theta \right) - (c + a\theta) \left(a - b\theta + \frac{1}{2}c\theta \right) \right\}$$

$$= g\rho\kappa (2ab\theta + 2bc\theta - c^2\theta - 2a^2\theta),$$

and this is positive if

$$2ab + 2bc > 2a^2 + c^2,$$

$$\text{or } c^2 - 2bc + b^2 < b^2 + 2ab - 2a^2.$$

If $c > b$, this leads to

$$c < b + \sqrt{b^2 + 2ab - 2a^2},$$

$c < b$, to

$$c > b - \sqrt{b^2 + 2ab - 2a^2}.$$


If we suppose the ends A, D , joined by a continuance of the tube and the figure $ABCD$ to be a square, $b = a$, and the condition is simply

$$c < 2a,$$

so that in this case the equilibrium is always stable.

(13) A rectangular lamina floats with two of its sides vertical in a liquid; it is required to determine when the equilibrium is stable for a small angular displacement such that the volume of liquid displaced remains unchanged.

In the figure let PQ be the line in the surface, and $P'Q'$ the line in the surface in the displaced position; H the centre of gravity of the liquid displaced in the position of equilibrium, and K, L the centres of gravity of the triangles EQQ', EPP' .



Draw HN, KM, LM' perpendicular to the horizontal line through G .

Then, if $GA = b, EA = c, BC = 2a$, and $\theta =$ the small angle QEQ' , the moment about G , tending to turn the rectangle back to its original position,

$$= g\rho \left(\frac{1}{2} a^2 \theta \cdot GM + \frac{1}{2} a^2 \theta \cdot GM' - 2ac \cdot GN \right),$$

but $GM = \frac{2}{3} a - EG \cdot \theta, GM' = \frac{2}{3} a + EG \cdot \theta$, and $GN = HG \cdot \theta$;

$$\therefore \text{the moment} = g\rho \left(\frac{2}{3} a^3 \theta - 2ac \theta \cdot HG \right),$$

$$\text{which is positive if } HG < \frac{a^2}{3c},$$

or

$$b - \frac{c}{2} < \frac{a^2}{3c}.$$

Let m be the point in which the line of action of the fluid

pressure after displacement meets HG ; then the moment above considered is equal to

$$gp2ac\theta \cdot Gm,$$

and, equating the two expressions, we obtain

$$\frac{2}{3}a^3 - 2acHG = 2ac(Hm - HG);$$

$$\therefore Hm = \frac{a^3}{3c}.$$

This point m is the *metacentre*, and we thus see that the stability depends on the position of this point with regard to G , as in Art. 63.

(14) *A cylindrical vessel, containing liquid, is raised upwards from rest with a given acceleration; it is required to determine the pressure at any point of the liquid.*

The acceleration here supposed may be obtained by attaching the vessel to a string passing over a fixed pulley, and having a weight at its other end; but, however the acceleration be obtained, the fact to be considered is that every element of the liquid ascends with a constant acceleration.

Taking P a point in the surface, imagine a thin prism PQ of the liquid to become rigid, and observe that its vertical acceleration is caused by the pressure of the liquid on the end Q , the atmospheric pressure on the end P , and the weight of the prism.

If $PQ = z$, $p =$ the pressure at Q , $\kappa =$ the area of a section of the prism, and $f =$ the given acceleration, we obtain, by aid of the second law of motion,

$$\rho z \kappa f = p \kappa - \Pi \kappa - g r z \kappa;$$

$$\therefore p = \Pi + \rho z (g + f).$$

Hence the whole pressure and the resultant pressure on the surface may be obtained as in the case of a liquid at rest, writing $g + f$ for g .

(15) *A closed vessel, just filled with liquid, slides down a smooth inclined plane; when the liquid is in a state of relative equilibrium it is required to find the pressure at any point and the surfaces of equal pressure.*

Every element of the liquid moves in a straight line with a constant acceleration $g \sin \alpha$, and since the forces on any element are the resultant fluid pressure upon it and its weight, it follows that the resultant of these forces is $mg \sin \alpha$, parallel to the plane, m being the mass of the element.

It is hence easy to see that the resultant pressure is perpendicular to the plane and is equal to $mg \cos \alpha$.

Whatever be the shape of the element, the resultant fluid pressure upon it in the direction parallel to the plane is zero, and therefore it follows that the surfaces of equal pressure are planes parallel to the inclined plane, and that the *surface* of the liquid is the plane through its highest point parallel to the inclined plane.

If there be no air within the vessel the pressure at the surface is zero, it being given that the vessel is only just filled, or, which is the same thing, just not filled.

Taking z as the depth of a point in the liquid below the surface thus defined, and drawing a thin cylinder or prism from this point to the surface, the pressure on the base will be the resolved part of the weight of the prism perpendicular to the plane, and, as before,

$$p\kappa = gpz\kappa \cos \alpha,$$

$$\text{or } p = gpz \cos \alpha.$$

As in the previous article the whole pressure and resultant pressure may be obtained, employing $g \cos \alpha$ for g .

The reasoning employed in this and the preceding example is applicable to any analogous case, that is, to any case in which the fluid, while bodily in motion, is within its own mass in a state of relative equilibrium.

MISCELLANEOUS PROBLEMS.

1. A triangle ABC is immersed in a fluid, its plane being vertical, and the side AB in the surface. If O be the centre of the circumscribing circle, prove that pressure on triangle OCA : pressure on triangle $OCB :: \sin 2B : \sin 2A$.

2. Water is gently poured into a vessel of any form; prove that when so much water has been poured in that the centre of gravity of the vessel and water is in the lowest possible position, it will be in the surface of the water.

3. A closed hollow cone is just filled with liquid, and is placed with its vertex upwards; divide its curved surface by a horizontal plane into two parts on which the whole pressures are equal.

Also do the same when the vertex is downwards.

4. If the cone be placed on its side on a horizontal table, compare the whole pressures on the curved surface and the base.

5. A triangle ABC has its plane vertical and the side AB in the surface of a liquid; divide it by straight lines drawn from A into n triangles on each of which the pressure shall be the same.

6. A solid displaces $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$ of its volume respectively when it floats in 3 different fluids; find the volume it displaces when it floats in a mixture formed, 1st, of equal volumes of the fluids, 2nd, of equal weights of the fluids.

7. A float is made by attaching to a hemisphere (radius r) a cone of the same base, and axis of length $2r$. If this will float in a fluid A with the cone just immersed, and in a fluid B with the hemisphere just immersed, compare the densities of A and B .

8. Compare the whole pressures on the curved surface and plane base of a solid hemisphere, radius r , immersed in water with its base horizontal and at a depth (r).

Note. The centre of gravity of the portion of the surface of a sphere contained between two parallel planes which intersect or touch the surface is equidistant from the planes.

9. A parabolic lamina floats in a liquid with its axis vertical and vertex downwards; having given the densities, σ , ρ , and the height (h) of the parabola, find the depth to which its vertex is immersed.

10. A heavy sphere, weight W , is placed in a vertical cylinder, filled with atmospheric air, which it exactly fits. Find the density of the air in the cylinder when the sphere is in a position of permanent rest, r being the radius and h the height of the cylinder.

11. If half a second be the unit of time, and the acceleration of a falling body that of acceleration, determine the ratio of the unit of density to the density of distilled water, in order that the formula, $p = g\rho z$, may give the pressure in pounds.

12. A cone, of given weight and volume, floats in a given fluid with its vertex downwards; shew that the surface of the cone in contact with the fluid is least, when the vertical angle of the cone is $2 \tan^{-1} \frac{1}{\sqrt{2}}$.

13. A hollow sphere is filled with fluid and a plane drawn through the centre divides the surface into two parts, the total normal pressures upon which are as $m : 1$; find the position of the plane and the greatest and least values of m .

14. A uniform tube is bent into the form of a parabola, and placed with its vertex downwards and axis vertical: supposing any quantities of two fluids of densities ρ , ρ' to be poured into it, and r , r' to be the distances of the two free surfaces respectively from the focus, then the distance of the common surface from the focus will be $\frac{r\rho - r'\rho'}{\rho - \rho'}$.

15. If water be the standard substance, 4 feet the unit of length, and 2 seconds the unit of time, find the unit of weight in the equation $W = g\rho V$, assuming 32 as the value of g when a foot and a second are units.

16. If there be n fluids arranged in strata of equal thickness, and the density of the uppermost be ρ , of the next 2ρ , and so on, that of the last being $n\rho$; find the pressure at the lowest point of the n^{th} stratum, and thence prove that the pressure at any point within a fluid whose density varies as the depth is proportional to the square of the depth.

17. A fine tube, bent into the form of an equilateral triangle with its vertex upwards and base horizontal, contains equal quantities of two liquids, each liquid filling a length of the tube equal to a side of the triangle. Prove that the height of the surface of the lighter fluid above that of the heavier : the altitude of the triangle :: $\rho' - \rho : \rho' + \rho$, ρ and ρ' being the densities.

18. A cylinder is filled with equal volumes of n different fluids which do not mix; the density of the uppermost is ρ , of the next 2ρ , and so on, that of the lowest being $n\rho$: shew that the whole pressures on the different portions of the curved surface of the cylinder are in the ratio

$$1^2 : 2^2 : 3^2 : \dots : n^2.$$

19. Equal volumes of n fluids are disposed in layers in a vertical cylinder, the densities of the layers, commencing with the highest, being as $1 : 2 : \dots : n$; find the whole pressure on the cylinder, and deduce the corresponding expression for the case of a fluid in which the *increase* of density varies as the depth.

Also, if the n fluids be all mixed together, shew that the pressure on the curved surface of the cylinder will be increased in the ratio $3n : 2n + 1$.

20. A hollow cone floats with its vertex downwards in a cylindrical vessel containing water. In the position of equilibrium the area of the circle in which the cone is intersected by the surface of the fluid bears to the base of the cylinder the ratio of 6 : 19. Prove that, if a volume of water equal to $\frac{19}{8}$ ths of the volume originally displaced by the cone be poured into the cone, and an equal volume into the cylinder, the position in space of the cone will remain unaltered.

21. A body is wholly immersed in a liquid and is capable of motion about a horizontal axis. It is found that the total pressure of the fluid on the surface is increased by A when the body

is turned through one right angle, and further increased by Z when it is turned through another right angle. Prove that the difference between the greatest and least pressures on the surface is $\sqrt{2(A^2 + B^2)}$.

22. A frustum of a right cone, formed by a plane parallel to the base and bisecting the axis, is closed and filled with fluid by means of a thin vertical pipe, which is also filled. If the top of this pipe be on a level with the vertex of the cone, find the whole pressure on the curved surface, and if this bear to the pressure on the base the ratio of 7 to 6, find the vertical angle of the cone.

23. If in the last example the base be removed, and the vessel then placed on a horizontal plane, and filled to the top of the pipe, find the least weight of the vessel which will prevent its being lifted.

24. An open cylindrical vessel, axis vertical, contains water, and a cone the radius of which is equal to that of the cylinder is placed in the water vertex downwards. Prove that, in the position of equilibrium, if the density of the cone be one-eighth of the density of water, the surface of the water will be raised above its original level through a height equal to one-twenty-fourth the height of the cone.

25. A solid cone of wood (density σ) rests with its base on the plane base of a large vessel, and water (density ρ) is then poured in to a given height; B a piece of the same wood is then attached by a string to the vertex of the cone so as to be wholly immersed; find what the size of the piece must be in order that it may just raise the cone.

26. An elliptic lamina floats with its plane vertical in a liquid of twice the density of the lamina, 1st, with its axis vertical, 2ndly, with its axis horizontal; determine in each case whether the equilibrium is stable or unstable, the lamina being displaced in its own plane.

27. A regular tetrahedron has one of its faces removed and is filled with fluid; the other faces, which are capable of moving round the lowest point, are kept together by means of strings which join the middle points of the horizontal edges of the vessel; shew that the tension of the strings is to the weight of the fluid as $\sqrt{3}$ to $4\sqrt{2}$.

28. A number of weights of different densities are attached to points of a thin weightless rod. Find the density of the fluid in which it is possible for them to rest, when all are totally immersed.

If there be three weights W_1, W_2, W_3 , of densities ρ_1, ρ_2, ρ_3 , respectively, and x, y be the distances of W_1, W_3 from W_2 , the middle weight, shew that, in order that the system may rest in equilibrium in any position when totally immersed in the corresponding fluid, the following condition must hold true,

$$\frac{x}{W_3} \left(\frac{1}{\rho_2} - \frac{1}{\rho_1} \right) + \frac{y}{W_1} \left(\frac{1}{\rho_3} - \frac{1}{\rho_2} \right) = \frac{x+y}{W_2} \left(\frac{1}{\rho_1} - \frac{1}{\rho} \right).$$

29. Two heavy liquids rest in equilibrium, one on the top of the other; one extremity of a heavy rod of length (a) is fixed at a given depth (c) in the lower liquid, and the other end reaches into the upper liquid. Find the positions of equilibrium, and determine whether they are stable or unstable.

30. A glass cylindrical vessel is inverted and plunged into water; by inclining the vessel half the air is allowed to escape, and the cylinder is then held vertically with the open end immersed and raised until one-fourth only of its length is below the surface; find the height of the water within.

31. A parallelogram is immersed in a fluid with a diagonal vertical, one extremity of which is in the surface of the fluid. Through this point lines are drawn dividing the parallelogram into three equal parts. Compare the pressures on these three parts; and, if P_2 be the pressure on the middle part, and P_1, P_3 those on the other two, prove that

$$16P_2 = 11(P_1 + P_3).$$

32. If a solid right cone whose angle is 2α be immersed in a liquid with its vertex in the surface and axis vertical, prove that if P be the whole pressure on the curved surface and base, and P' the resultant pressure,

$$\frac{P}{P'} = \frac{2 + 3 \sin \alpha}{\sin \alpha}.$$

Also, determine this ratio when the axis is inclined at an angle θ to the vertical, θ being less than the complement of α .

33. Three faces of a regular tetrahedron, which rests with the remaining face on a horizontal table, are heavy plates capa-

ble of moving about their horizontal edges. If they fit accurately and the tetrahedron be filled with fluid through a small hole at the vertex, shew that it will hold together if the ratio of the weight of each plate to the weight of the contained fluid be not less than 9 to 2.

34. A vertical cylinder is closed by an airtight piston, and when the piston is at the top of the cylinder it is filled with vapour at a given pressure: if temperature be such as would maintain vapour of three times the density, find the least weight of the piston which will not condense the whole of the vapour.

35. If a Differential Thermometer be constructed with unequal bulbs, will it shew any indication of a change of temperature to which both bulbs are subject?

36. A thin conical surface (weight W) just sinks to the surface of a fluid when immersed with its open end downwards; but when immersed with its vertex downwards a weight equal to mW must be placed within it to make it sink to the same depth as before. Shew that if a be the length of the axis, and h the height of a column of the fluid, the weight of which equals the atmospheric pressure,

$$\frac{a}{h} = \frac{1}{m} \sqrt[3]{1 + \frac{1}{m}}.$$

37. If A be the area of the section of each pump of a fire engine, l the length of the stroke, n the number of strokes per minute, B the area of the hose, find the mean velocity with which the water rushes out.

38. A piston without weight fits into a vertical cylinder, closed at its base and filled with air, and is initially at the top of the cylinder; water being poured slowly on the top of the piston, find how much can be poured in before it will run over. Explain the case in which the height of the cylinder is less than the height of the water barometer.

39. Within a cylinder of height a , open at the top, is placed another cylinder of the same height, and half the content, closed at the top, and a quantity of mercury sufficient to fill the interior cylinder is poured into the exterior. If x and y be the dis-

tances of the surfaces in the two cylinders from the top, prove that

$$\frac{y}{x}(y-x)=h,$$

and find x and y ; h being the height of the mercury barometer.

40. The sides of a rectangle are in the ratio $\pi : 4$, and semicircles are described on the longer sides as diameters. Prove that, if the rectangle be immersed in water, with one of the shorter sides in the surface, the pressure on the two parts external to both semicircles will together be equal to that on the part common to them.

41. A plane rectangular lamina is bent into the form of a cylindrical surface of which the transverse section is a rectangular hyperbola. If it be now immersed in water so that first the transverse, secondly the conjugate, axes of the hyperbolic sections be in the surface, prove that the horizontal pressure on any the same immersed surface will be in the two cases the same.

42. A double funnel formed by joining two equal hollow cones at their vertices stands upon a horizontal plane with the common axis vertical, and fluid is poured in until its surface bisects the axis of the upper cone. If the fluid be now on the point of escaping between the lower cone and the plane, prove that the weight of either cone is to that of the fluid it can hold as 27 : 16.

43. A square lamina $ABCD$, which is immersed in water, has the side AB in the surface; draw a line BE to a point E in CD such that the pressures on the two portions may be equal. Prove that, if this be the case, the distance between the centres of pressure : the side of the square $:: \sqrt{505} : 48$.

44. A cubical vessel, having one of its vertical sides moveable about a hinge in the base, is filled with water, the moveable side inclining inwards; prove that the tangent of its inclination to the horizon is to unity as the weight of the side is to the weight of the water contained by the vessel when the side is vertical.

45. A semicircular area is immersed in a liquid with its bounding diameter in the surface; find the pressure on any por-

tion of the area contained between two radii, and find the area contained between the surface and a radius such that the pressure upon it may be one-fourth of the pressure upon the whole.

46. A vertical cylinder is filled with liquid; find the centre of pressure of the portion of its curved surface contained between two vertical planes through the axis.

47. Find the centre of pressure of the surface contained between two planes drawn through a radius of the top of the cylinder, and through the extremities of that diameter of the base which is perpendicular to the radius.

Also, find the centre of pressure of the same surface when the cylinder is inverted.

48. A solid, in the form of a right pyramid, the base of which is a regular polygon of n sides, is completely immersed in a liquid, with its base vertical; find the direction and magnitude of the resultant pressure on its inclined surfaces.

Solve the same question when the base is inclined to the vertical at a given angle.

49. An oblique cone on a circular base is completely immersed in water with its base vertical; find the resultant pressure on the curved surface.

50. A vessel in the form of an oblique cone on a circular base is held with its base horizontal and vertex downwards and is filled with liquid; find the resultant pressure on the surface and its point of action.

51. If a parabolic area be just immersed in water, and be turned about in a vertical plane so that the surface is always a tangent, prove that the centre of pressure of the part above a fixed horizontal plane lies in the diameter through the point of contact and at a given distance from that point.

52. A portion of a right circular cone cut off by a plane through the axis and a plane perpendicular to the axis is immersed in fluid with the vertex in the surface, and axis vertical; shew that the resultant horizontal pressure on any part of the curved surface intercepted between two horizontal planes will pass through the centre of gravity of the intercepted portion of the cone.

53. In exhausting a receiver by an air-pump a cloud is sometimes seen in the receiver; explain the cause of this.

54. A hollow sphere is just filled with liquid; find the line of action and magnitude of the resultant pressures on either of the portions into which it is divided by a vertical plane through its centre.

55. A hollow cone, vertex downwards, is filled with liquid; find the direction and magnitude of the resultant pressure on the portion of its surface contained between two vertical planes through its axis.

56. Solve the same question when the vertex of the cone is upwards.

57. A hollow cylinder is closed at one end and open at the other, and a fixed stop perpendicular to the axis divides the cylinder into two equal parts cutting off the communication between the parts; the weight of the whole cylinder is half the weight of the water which it would contain. Prove that if the cylinder be placed mouth downwards in water the depth of the stop in the position of rest will be only half as great as if a hole had been made in the stop.

58. If a thermometer plunged incompletely in a liquid whose temperature is required indicate a temperature t , and τ be that of the air, the column not immersed being m degrees, prove that the correction to be applied is $\frac{m(t-\tau)}{6480+\tau-m}$, $\frac{1}{6480}$ being the expansion of mercury in glass for 1° of temperature, assuming that the temperature of the mercury in each part is that of the medium which surrounds it.

59. A weightless cone is very nearly filled with liquid and inverted on a horizontal table; the liquid is made to rotate with an angular velocity ω , and the pressure required to keep the cone in contact with the table is equal to three times the weight of the liquid; prove that

$$\omega = 2 \cot \alpha \sqrt{\frac{g}{3h}},$$

where h is the height of the cone, and α the semivertical angle.

60. A right circular cone is constrained to rest in a fluid with its axis horizontal and the highest point C of its base in the surface of the fluid. Find the magnitude and direction of the resultant fluid pressure on the curved surface of the cone, and determine the vertical angle of the cone when the direction of

this pressure (1) passes through C , (2) is parallel to a generating line. Shew also that its direction can never be perpendicular to a generating line.

61. A conical vessel, having its vertex downwards, is filled with two liquids which do not mix, their common surface bisecting the axis; compare the whole pressures on the two portions of the surface.

62. A tube, in the form of an equilateral triangle, is filled with equal volumes of three liquids, the densities of which are as 1 : 2 : 3; if the tube be held with one side horizontal, and the opposite angle upwards, prove that the common surfaces of the liquids divide the sides in the ratio 1 : 2.

63. An open hemispherical cup, filled with water, is placed on a horizontal table, and the whole is made to rotate uniformly about its vertical radius; prove that the pressure on the table : the original weight of liquid :: $8g - 3\omega^2 r$: $8g$.

64. A hollow vessel in the shape of a wedge of a cylinder, formed by two planes through its axis, is filled with water and closed at the top; it is then made to rotate uniformly about the axis, which is vertical; find the pressure on the top.

65. In the previous problem find the whole pressure on the curved surface of the cylinder.

66. An isosceles triangular prism, the vertical angle of which is a right angle, floats in water with its edge horizontal, and its base above the surface, find its positions of equilibrium.

67. A cone is totally immersed in a fluid, the depth of the centre of its base being given. Prove that P , P' , P'' , being the resultant pressures on its convex surface, when the sines of the inclination of its axis to the horizon are s , s' , s'' , respectively,

$$P^2 (s' - s'') + P'^2 (s'' - s) + P''^2 (s - s') = 0.$$

68. A hollow cone filled with liquid is suspended freely from a point in the rim of its base; prove that the total pressures on the curved surface and the base are in the ratio

$$1 + 11 \sin^2 \alpha : 12 \sin^3 \alpha.$$

69. A hollow cone without weight, closed and filled with water, is suspended from a point in the rim of its base; if ϕ be

the angle which the direction of the resultant pressure on the curved surface makes with the vertical, and α the semi-vertical angle of the cone, prove that

$$\cot \phi = \frac{28 \cot \alpha + \cot^3 \alpha}{48}.$$

70. A heavy uniform chain is suspended from its two ends under water; prove that its form will be the same as if suspended in air.

71. An open conical shell, the weight of which may be neglected, is filled with water, and is then suspended from a point in the rim, and allowed gradually to take its position of equilibrium; prove that, if the vertical angle be $\cos^{-1} \frac{2}{3}$, the surface of the water will divide the generating line through the point of suspension in the ratio of 2 : 1.

72. A tube of small bore in the form of an ellipse is half filled with equal volumes of two fluids which do not mix; find in what manner the tube must be placed in order that the free surfaces of the two fluids may be the extremities of the minor axis.

73. If any curved surface, having for its base a plane area A and enclosing a volume V , be totally immersed in a fluid, find the resultant pressure on the curved surface, when the depth of the centre of gravity, and the inclination to the horizon, of the plane of the base are given.

If P_1, P_2, P_3 , be these resultant pressures when the depths of the centre of gravity of the base, in a fluid of density ρ , are x, y, z respectively, and the inclinations of the base to the horizon are the same, shew that

$$P_1^2(z-y) + P_2^2(x-z) + P_3^2(y-x) = g^2 \rho^2 A^2 (z-y)(x-z)(y-x).$$

74. A heavy chain is suspended from two points and hangs partly immersed in a fluid; shew that the curvatures of the portions just inside and just outside the surface of fluid are as $\rho - \sigma : \rho$, ρ and σ being the densities of the chain and fluid.

75. Close to the base of a vertical cylinder there is a small aperture turned upwards as in the figure, Art. 170, but, instead

of the surface in the cylinder being free, a heavy piston rests upon it; find the height to which the jet rises.

76. A ball of lead is let fall in water; assuming that the pressure of the water is the same as if the ball were not in motion, find its velocity at any given depth.

77. A weightless inextensible envelope full of air floats in equilibrium in the receiver of an air-pump; find the velocity of its descent after n strokes of the piston, supposed instantaneous, and made at equal intervals.

78. If the volume of the receiver be n times that of the piston, and if v be the limit of the above velocity when n is infinite, and v' the velocity which would have been obtained in vacuo in the same time, shew that $v' = \epsilon v$.

79. A spherical bubble of air ascends in water; having its size at depth a , find its size when its depth is $\frac{1}{2}a$.

80. A vertical cylinder containing water is made to rotate with a uniform angular velocity about its axis: if $\frac{1}{n}$ th of the axis of the cylinder was above the surface before the rotation commenced, shew that the greatest angular velocity that can be given to the cylinder, without causing any of the water to leave the cylinder, is

$$\frac{2}{a} \left(\frac{gh}{n} \right)^{\frac{1}{2}},$$

where h is the height of the cylinder, and a the radius of its base.

81. A bent tube ABC contains fluid, and the tube rotates uniformly with an angular velocity ω about the leg AB , which is vertical: find the position of equilibrium of the fluid.

If l be the whole length of tube occupied by the fluid, and the angle $ABC = \alpha$, examine the case in which $\omega^2 > \frac{g}{2l} \cot^2 \frac{\alpha}{2}$.

82. Two equal uniform rods AB, AC are rigidly connected at A , and the system floats symmetrically with the point A downwards.

If a be the length of each rod, and c the length of each immersed, prove that the equilibrium will be stable for a small angular displacement in the vertical plane of the rods if

$$c(3 - \cos \omega) > a(1 + \cos \omega),$$

where ω is the angle BAC .

83. A hollow vertical polygonal prism, open at both ends, rests upon a horizontal plane; every two contiguous faces are moveable about their common edge. Supposing the prism to be in equilibrium, when filled with fluid, prove that

$$\frac{c_1}{\sin a_1} = \frac{c_2}{\sin a_2} = \frac{c_3}{\sin a_3} = \dots$$

a_1, a_2, a_3, \dots being the angles of a transverse section $A_1A_2A_3 \dots A_nA_1$, and c_1, c_2, c_3, \dots denoting the lines $A_nA_3, A_1A_3, A_2A_4, \dots$

84. Two cylindrical vessels containing water are suspended with their axes vertical to the ends of a string passing over a fixed smooth pulley in a vertical plane; neglecting the weights of the vessels, compare the whole pressures, during the motion, on the curved surfaces of the cylinders.

85. Through the plane vertical side of a vessel containing fluid, small holes are bored in the circumference of a circle, which has its highest point in the surface of the fluid; shew that the trace of the issuing fluid on a horizontal plane through the lowest point of the circle is two straight lines.

86. A tuning fork held over a glass jar of a certain depth has its sound greatly augmented; but a jar an inch deeper, or an inch shallower, produces but a slight augmentation. Why is this the case?

87. On clapping your hands near a long railing, a sound is heard resembling that produced by the swift passage of a switch through the air; state the cause of this sound.

88. A hollow cone, vertex downwards, and containing liquid, is attached to a string passing over a pulley and supporting at its other end a given weight: determine the motion and find the whole pressure of the fluid on the cone and also the resultant pressure.

89. Two vessels contain air having the same pressure H but different temperatures t, t' ; the temperature of each being increased by the same quantity, find which has its pressure most increased.

If the vessels be of the same size, and the air in one be forced into the other, find the pressure of the mixture at a temperature zero.

90. The temperature of the air in an extensible spherical envelope is gradually raised from 0° to t° , and the envelope is allowed to expand till its radius is n times its original length; compare the pressure of the air in the two cases.

91. A cylindrical vessel, closed at both ends, and placed so that its axis is vertical, is half filled with mercury at a temperature $0^\circ C$, the remaining space being occupied by air at the same temperature. The expansion of mercury between the temperatures 0° and $100^\circ C$ being $\cdot 018$ of its original volume, and that of air $\cdot 3665$ of its original volume for the same pressure, shew that if the temperature be raised to $20^\circ C$ the pressure of the air will be increased in the ratio $1\cdot 0772 : 1$.

92. If a given body lose in air, when the height of the barometric column is h , the m^{th} part of its weight, find what part of its weight it will lose when the height of the barometric column is h' .

93. The specific gravity of mercury compared with that of water at 68° is $13\cdot 568$ and at 212° is $13\cdot 704$. If the expansion of mercury between these points be $\frac{1}{69}$ th of its volume at the lower temperature, find that of water between the same points.

94. A hemispherical bowl is filled with water; if the internal surface be divided by horizontal planes into n portions, on each of which the whole pressure is the same, and h_r be the depth of the r^{th} of these planes, prove that

$$\frac{h_r}{a} = \sqrt{\frac{r}{n}},$$

a being the radius.

95. If a lamina in the form of a regular hexagon be immersed in liquid with one side in the surface, the depth of its centre of pressure is to the depth of its centre of gravity as 23 to 18.

96. Find the centre of pressure upon a portion of a vertical cylinder containing liquid, the portion being such as when unwrapped to form an isosceles triangle, the base of which when forming part of the cylinder is horizontal, and the vertex at the surface of the fluid.

97. Two very small spheres, of the same size but different densities, are connected by a fine string and immersed in a liquid, which rotates uniformly about a fixed axis, and is not acted upon by any forces; find their position of relative equilibrium.

98. A hollow cone open at the top is filled with water; find the resultant pressure on the portion of its surface cut off, on one side, by two planes through its axis inclined at a given angle to each other; also determine the line of action of the resultant pressure, and shew that, if the vertical angle be a right angle, it will pass through the centre of the top of the cone.

99. Two equal light spheres of the same substance are attached by strings of lengths r, r' to a point in the bottom of a vessel of water—they are mutually repulsive and rest at a distance x from each other: shew that the line joining them is inclined to

the horizon at
$$\sin^{-1} \frac{r^2 \sim r'^2}{x \sqrt{2(r^2 + r'^2) - x^2}};$$

also if $\phi(x)$ be the repulsion

$$\phi(x) = \frac{Px}{\sqrt{2(r^2 + r'^2) - x^2}},$$

P being the fluid pressure on either sphere.

100. A cylindrical tube, containing air, is closed at one extremity by a fixed plate, the other extremity being open; a piston just fitting the tube slides within it, and the centres of the plate and piston are connected by an elastic string, the modulus of elasticity of which is equal to the atmospheric pressure on the piston; prove that, if l be the natural length of the string, and a its length when the air between the piston and the fixed plate is in its natural state, l being less than a , the length of the string in the position of equilibrium will be $(la)^{\frac{1}{2}}$.

101. The readings of a faulty barometer containing some air are 29.4 and 29.9 inches, the corresponding readings of a correct instrument being 29.8 and 30.4 inches respectively; prove that the length of the tube occupied by the air is 2.9 inches, when the reading of the barometer is 29 inches; and find the corresponding correct reading.

102. A cylinder of density 2ρ floats with its axis vertical between two liquids of densities ρ and 3ρ , its height being equal to the depth of the upper liquid; prove that the pressures on its ends are in the ratio of 1 to 5.

103. A heavy rope, the density of which is double the density of water, is held by one end, which is above the surface, the other end being under water; find the tension at the middle section of the immersed portion.

104. If the depths of the angular points of a triangle below the surface of a fluid be a, b, c , shew that the depth of the centre of pressure below the centre of gravity is

$$\frac{(b-c)^2 + (c-a)^2 + (a-b)^2}{12(a+b+c)}.$$

105. Given that the centre of pressure of a disc of radius r , with one point in the surface, is at a distance p from the centre, prove that for a disc of radius R wholly immersed with its centre at a distance h from the surface, the distance between the centre of the circle and the centre of pressure is $pR^2 \div hr$.

106. If an air-pump be fitted with a barometer gauge of small section κ , and length l , prove that at the end of the first stroke the mercury will have risen a height

$$\frac{Bh}{A+B} \left\{ 1 - \kappa \frac{Ah + (A+B)l}{(A+B)^2} \right\} \text{ nearly;}$$

h being the height of the barometer.

107. A hemispherical shell is floating on the surface of a liquid, and it is found that the greatest weight which can be attached to the rim is one-fourth of the weight of the hemisphere; prove that the weight of the liquid which would fill the hemisphere bears to the weight of the hemisphere the ratio of

$$25\sqrt{5} : 20\sqrt{5} - 28.$$

108. A cylindrical diving-bell fully immersed is in equilibrium without a chain. Shew that if the exterior atmospheric pressure increase slightly, the ratio of the distance moved through by the bell if free to that moved through by the surface of the water in the bell when held fixed is $Hh + x^2 : x^2$ approximately; where H is the height of the water barometer, h the height of the bell, and x the length of that part of it which is filled with air.

96. Find the centre of pressure upon a portion of a vertical cylinder containing liquid, the portion being such as when unwrapped to form an isosceles triangle, the base of which when forming part of the cylinder is horizontal, and the vertex at the surface of the fluid.

97. Two very small spheres, of the same size but different densities, are connected by a fine string and immersed in a liquid, which rotates uniformly about a fixed axis, and is not acted upon by any forces; find their position of relative equilibrium.

98. A hollow cone open at the top is filled with water; find the resultant pressure on the portion of its surface cut off, on one side, by two planes through its axis inclined at a given angle to each other; also determine the line of action of the resultant pressure, and shew that, if the vertical angle be a right angle, it will pass through the centre of the top of the cone.

99. Two equal light spheres of the same substance are attached by strings of lengths r, r' to a point in the bottom of a vessel of water—they are mutually repulsive and rest at a distance x from each other: shew that the line joining them is inclined to

the horizon at $\sin^{-1} \frac{r^2 \sim r'^2}{x \sqrt{2(r^2 + r'^2) - x^2}}$;

also if $\phi(x)$ be the repulsion

$$\phi(x) = \frac{Px}{\sqrt{2(r^2 + r'^2) - x^2}},$$

P being the fluid pressure on either sphere.

100. A cylindrical tube, containing air, is closed at one extremity by a fixed plate, the other extremity being open; a piston just fitting the tube slides within it, and the centres of the plate and piston are connected by an elastic string, the modulus of elasticity of which is equal to the atmospheric pressure on the piston; prove that, if l be the natural length of the string, and a its length when the air between the piston and the fixed plate is in its natural state, l being less than a , the length of the string in the position of equilibrium will be $(la)^{\frac{1}{2}}$.

101. The readings of a faulty barometer containing some air are 29.4 and 29.9 inches, the corresponding readings of a correct instrument being 29.8 and 30.4 inches respectively; prove that the length of the tube occupied by the air is 2.9 inches, when the reading of the barometer is 29 inches; and find the corresponding correct reading.

102. A cylinder of density 2ρ floats with its axis vertical between two liquids of densities ρ and 3ρ , its height being equal to the depth of the upper liquid; prove that the pressures on its ends are in the ratio of 1 to 5.

103. A heavy rope, the density of which is double the density of water, is held by one end, which is above the surface, the other end being under water; find the tension at the middle section of the immersed portion.

104. If the depths of the angular points of a triangle below the surface of a fluid be a, b, c , shew that the depth of the centre of pressure below the centre of gravity is

$$\frac{(b-c)^2 + (c-a)^2 + (a-b)^2}{12(a+b+c)}.$$

105. Given that the centre of pressure of a disc of radius r , with one point in the surface, is at a distance p from the centre, prove that for a disc of radius R wholly immersed with its centre at a distance h from the surface, the distance between the centre of the circle and the centre of pressure is $pR^2 \div hr$.

106. If an air-pump be fitted with a barometer gauge of small section κ , and length l , prove that at the end of the first stroke the mercury will have risen a height

$$\frac{Bh}{A+B} \left\{ 1 - \kappa \frac{Ah + (A+B)l}{(A+B)^2} \right\} \text{ nearly;}$$

h being the height of the barometer.

107. A hemispherical shell is floating on the surface of a liquid, and it is found that the greatest weight which can be attached to the rim is one-fourth of the weight of the hemisphere; prove that the weight of the liquid which would fill the hemisphere bears to the weight of the hemisphere the ratio of

$$25\sqrt{5} : 20\sqrt{5} - 28.$$

108. A cylindrical diving-bell fully immersed is in equilibrium without a chain. Shew that if the exterior atmospheric pressure increase slightly, the ratio of the distance moved through by the bell if free to that moved through by the surface of the water in the bell when held fixed is $Hh + x^2 : x^2$ approximately; where H is the height of the water barometer, h the height of the bell, and x the length of that part of it which is filled with air.

109. Two hollow cones, filled with water, are connected together by a string attached to their vertices which passes over a fixed pulley; prove that, during the motion, if the weights of the cones be neglected, the total pressures on their bases will be always equal, whatever be the forms and dimensions of the cones. If the heights of the cones be h, h' , and heights mh, nh' be unoccupied by water, the total normal pressures on the bases during the motion will always be in the ratio

$$n^2 + n + 1 : m^2 + m + 1.$$

110. A hollow cone, whose vertical angle is given, is filled with water and placed with its base on a horizontal plane; determine a point in its surface at which, an orifice being made, the issuing fluid will just fall outside the base of the cone.

111. The times of the aerial vibrations constituting a note (C) and its fifth (G) are in the ratio 3 : 2; compare the times of the vibrations corresponding to (C) and the fifth of (G).

112. A pyramid on a square base floats with its vertex downwards and base horizontal in a liquid. The pyramid is bisected by a vertical plane perpendicular to two sides of the base, and the two parts are connected at the vertex by a hinge. Prove that the parts will remain in contact if the ratio of the density of the pyramid to that of the liquid exceed

$$\left(\frac{3a^2}{2h^2 + 3a^2} \right)^3,$$

where h is the height and $2a$ the side of the base.

113. A circular tube of fine bore, whose plane is vertical, contains a quantity of heavy uniform fluid, which subtends an angle 2α at the centre; a heavy spherical particle, just fitting the tube, is let fall from the extremity of a horizontal radius; find the impulsive pressure at any point of the fluid.

114. A cylindrical vessel containing inelastic fluid is descending with a given velocity (v) and is suddenly stopped; its axis being vertical, find the whole impulse on the curved surface.

115. A closed hollow cone, filled with inelastic fluid, and having its vertex upwards, is suddenly raised with a given velocity; find the whole impulse on the curved surface, and the resultant impulse on the base.

116. A hollow sphere formed of a rigid inelastic substance, and filled with inelastic fluid, is let fall on a horizontal plane; find the whole impulse on its curved surface, and on each half of its surface above and below the horizontal plane through its centre.

Also determine the resultant impulses on each of these surfaces.

117. A flexible and elastic cylindrical tube is placed within a rigid hollow prism, in the form of an equilateral triangle, which it just fits when unstretched; if there be no air between the tube and the prism, and if air at a given pressure be forced into the tube, find the extension and the portion in contact with the sides of the prism.

118. A conical bag, which is filled with liquid, has its rim fastened to a horizontal plate, and is then inverted; prove that the tension at any point, in the direction of a generating line, varies as the square of the distance from the vertex.

119. A bag, in the form of a paraboloid, formed of thin flexible substance, is supported by its rim, and is filled with water; find the tension at any point in direction of the tangent to the generating parabola at the point.

Hence prove that the tension in every direction at the vertex $=g\rho ah$, if h be the depth of the bag, and $4a$ the latus rectum.

Also obtain this last result independently by aid of Art. (167).

120. If the same bag, when filled, be closed and inverted, prove that the tension at any point P , in direction of the tangent to the generating parabola, varies as $AN \cdot \sqrt{SP}$, A being the vertex of the bag, S the focus, and AN the depth of P below the vertex.

SPECIFIC GRAVITIES.

Ratios of the Specific Gravities of different substances to that of water at 60°.

Diamond	3.52		Nickel	8.38
Sulphur	2.		Iron	7.844
Iodine	4.94		Flint-glass	3.33
Arsenic	5.959		Plate-glass	2.5
Gold	19.4		Marble	2.716
Platina	21.53		Rock-salt	1.92
Silver	10.5		Ivory	1.917
Mercury	13.568		Ice (at 0°)	0.926
Copper	8.85		Sea-water	1.027
Tin	7.285		Olive-oil	0.915
Lead	11.445		Alcohol	0.794
Zinc	6.862		Æther	0.724

Ratios of the densities of gases and vapours of different substances to that of atmospheric air at the same temperature and under the same pressure.

Oxygen	1.103		Water	0.62
Hydrogen	0.069		Alcohol	1.613
Nitrogen	0.976		Carbonic Acid	1.524
Chlorine	2.44		Ammonia	0.591
Bromine	5.395		Sulphurous Acid	2.212
Iodine	8.701		Sulphuric Acid	2.763
Arsenic	10.365		Æther	2.586
Mercury	6.978			

ANSWERS TO THE EXAMPLES.

CHAPTER I. EXAMINATION.

4. $10\frac{1}{2}$ lbs. and 42 lbs. 5. *wa.*
 6. 180 lbs. 8. 82944 lbs.

CHAPTER II. EXAMINATION.

2. 848 lbs. 3. 104976 lbs.
 4. $\frac{m+1}{mn+1}\sigma$ and $\frac{mn+n}{mn+1}\sigma$, σ being the specific gravity of the mixture.

5. 2ρ . 6. $202\frac{1}{2}$ lbs. 7. 13. 8. $3\sigma - s' - s''$.
 9. $\frac{V(\sigma - s) + V'(\sigma' - s)}{s}$. 10. $2s$.

CHAPTER II. EXAMPLES.

1. $\frac{n+1}{2}$, $\frac{2n+1}{3}$, and $\frac{n+2}{3}$.
 2. $3\sigma' - 2\sigma$ and $4\sigma - 3\sigma'$. 3. 4 : 405.
 4. 84 : 125. 5. 1 : 32.
 6. $\frac{1}{\sqrt{32}}$ th of a second.
 7. The densities are as 3 : 8. 8. 4 : 1.
 9. 9 : 512.

CHAPTER III. EXAMINATION.

2. 1st. $43\frac{2}{3}$ lbs. on a square inch. 2nd. about 58 lbs.
 4. $73\frac{1}{4}$ lbs. on a square inch, neglecting atmospheric pressure.
 6. 125 oz. 8. $\frac{2}{3}g\rho\pi rh\sqrt{r^2+h^2}$.

11. If h be the vertical side, the depth of the horizontal line = $\frac{h}{\sqrt{2}}$.

12. The depths of the horizontal lines are

$$\frac{1}{\sqrt{n}} h, \quad \sqrt{\frac{2}{n}} h, \quad \sqrt{\frac{3}{n}} h, \quad \&c.$$

13. The depth = $\frac{1}{\sqrt{2}} h$.

CHAPTER III. EXAMPLES.

2. 3125 lbs.

3. The line divides the opposite side in the ratio of 3 : 1.

4. $\frac{1}{8\sqrt{2}}$ (whole length of liquid).

5. $16 + \frac{125\pi}{4}$ lbs.

6. $20 + \frac{125\pi}{4}$ lbs.

8. 1 : 1.

10. The point lies in the line from the vertex bisecting the base and at a depth $\frac{1}{\sqrt{3}}$ (the depth of the vertex).

11. 1 : $\sqrt{2} - 1$.

12. $\frac{4}{3}(1 + \sqrt{10})$ inches.

13. 1 : 4 : 9.

14. If α , $\pi - \alpha$, be the angular spaces occupied by the liquids ρ , ρ' , the inclination to the vertical of the bounding diameter = $\tan^{-1} \left(\cot \alpha + \frac{1}{\sin \alpha} \frac{\rho' + \rho}{\rho' - \rho} \right)$.

16. The increase = 14 (the weight of the fluid).

18. $\left\{ 1 - \frac{\rho'}{\rho} \left(1 - \frac{1}{\sqrt{2}} \right) \right\}$. radius.

20. Produce the rectangle to the surface ; then, knowing the centres of pressure of the whole and of the upper part, and the pressures on these parts, the position of the centre of pressure of the lower part can be inferred.

23. The densities are equal.

24. 2 feet.

25. Unit of time = $\frac{2}{5\sqrt{5}}$ seconds ;

Unit of space = $\frac{2}{5\sqrt{5}}$ feet. 26. $\frac{5\sqrt{15}}{2}$ secs.

27. 3π units of weight.

29. $4(5\sqrt{2}-7) : 3$.

30. $3 : 6 : 4$.

CHAPTER IV. EXAMINATION.

7. 12 feet.

8. $15 : 16$.

9. The forces are equal.

13. $\frac{4}{5}$ ths of a cubic foot.

15. Weight of wood + weight of water it displaces.

16. If σ , σ' be the specific gravities, V , V' the volumes, and ρ the specific gravity of water, the condition is

$$\frac{V}{V'} = \frac{\sigma' - \rho}{\sigma - \rho}.$$

CHAPTER IV. EXAMPLES.

2. $\frac{1}{866}$ of a cubic yard.

3. Surface divides altitude in ratio $1 : \sqrt{2} - 1$.

4. $\sqrt[3]{2} - 1 : 1$.

7. $\frac{1}{4}$ th of the cylinder is in the upper liquid.

8. $\frac{9}{4}$. density of wood.

9. Half that of water.

10. $\left(\frac{375\pi}{64} - 8\right)$ oz.

14. $\sigma : \rho = r^3 : 2(r^3 - r)$. \times

15. If w , w' be the weight of the cone and of the fluid displaced, the force = $w - w'$, and its line of action must be at a distance from the centre of gravity of the solid cone equal to

$$\frac{w - w'}{12} \cdot \frac{h}{2}.$$

16. One-third of the axis is immersed.

18. $h\sqrt{\frac{\sigma}{\rho}}$, h being the height of the paraboloid.

19. $2186\frac{2}{3}$ tons.

20. Height = $\frac{r^2}{r'^2} \frac{h}{2}$.

21. $gpr^2h\left(\frac{\pi+4}{2}\right)$, and $gprh^2$ at depth $\frac{2}{3}h$.

22. $\frac{r}{\sqrt{2}}$, assuming distance of centre of gravity of hemisphere from centre to be $\frac{3}{8}r$.

24. 19 : 56.

25. $\sqrt{3} \cdot r$.

33. The resultant vertical pressure = $g\rho hr^2 \left(1 - \frac{\pi}{6}\right)$.

34. Horizontal pressure = $W \sin a \cos a$, and vertical pressure = $W \sin^2 a$, where W is the weight of fluid displaced. Hence direction of resultant is inclined to the vertical at an angle $\frac{\pi}{2} - a$, as is obvious, *a priori*.

CHAPTER V. EXAMINATION.

2. $4\frac{4}{9}$, $3\frac{5}{9}$.

4. 72.7 inches nearly.

5. 1 : 12.

6. 1 : 8 ($1 + at$), and 1 : 2 ($1 + at$).

10. $11\frac{3}{7}$ and $-11\frac{3}{7}$.

17. 1909 lbs. nearly.

CHAPTER V. EXAMPLES.

1. $1 + at : n^3$.

3. If W , W' be the weights of the fluid and the piston, Π the pressure of the fluid at a density ρ , the length of cylinder occupied = $\frac{W}{W'} \cdot \frac{\Pi}{g\rho}$.

4. $\frac{P'}{P} = n \frac{1 + at'}{1 + at}$.

5. If m be the ratio of the air-pressure on the piston to its weight when the piston is in the middle, its height above the base is given by the equation

$$x^2 - 2ax + 2ma(a - x) = 0,$$

$2a$ being the height of the cylinder.

6. (The area of surface) $\times \kappa pat$.

8. $h : h'$.

9. If Π be external air-pressure, W the weight of the piston, A its area, λ the modulus of elasticity, a the natural length of the string, and t the increase of temperature, the increase x of the length of the string is given by the equation

$$\lambda(ax + x^2) = a(aat - x)(\Pi A + W).$$

10. Length above surface is changed in ratio 1 : .9987.

14. $7h$, where h is the height of the water barometer.

16. If a be the depth to which the original open surface of the mercury is lowered, ρ , σ the densities of the water and mercury, and k , K the sectional areas of the tube, the height of the mercurial column is increased by

$$\frac{\rho \left(1 + \frac{K'}{k}\right)}{\sigma \left(1 + \frac{K}{k}\right) - \rho} \cdot a.$$

CHAPTER VI. EXAMINATION.

1. To $\frac{1}{3}$ rd of its original volume.
5. 512 lbs.
6. Early in the 4th stroke.
9. About 45 inches above mercury level.
11. $\frac{5000\pi}{16}$ lbs. or about 983 lbs.
12. $\frac{3000\pi}{16}$ lbs.

CHAPTER VI. EXAMPLES.

1. $1 - \left(\frac{4}{5}\right)^5$.
4. 1.8 nearly; $\frac{2}{11}$ ths of the volume of the bell.
6. If h be the height of the barometer, the ascent x is given by the equation

$$\frac{15}{15-x} + \frac{x}{h} = \frac{5}{3} + \frac{20}{3h}.$$

If $h=30$, $x=6.1$ nearly.

11. If a be the length originally occupied by air, h the height of the barometer, and ρ the density of atmospheric air, the difference x is given by the equation

$$x^2 + 2a(x-h) + \frac{\rho^n}{\rho} h(2a+x) = 0$$

CHAPTER VII. EXAMINATION.

1. 5 : 7.
2. 9 : 7.
3. $1280 - \pi$: $1280 - 2\pi$.
6. $\frac{198}{1805}$ of a cubic foot.
7. $\frac{25}{21}$ and $\frac{20}{21}$.

8. $W+w : W+sv'$, W being the weight of the instrument, and s the density of water.

9. $\frac{4}{5}$.

10. $\frac{4}{5}$.

CHAPTER VII. EXAMPLES.

1. The same as that of water.

3. $\frac{729}{217}(3.17) = 10.8$ nearly.

4. 1.9 nearly.

5. $4 : 5$.

6. $4 : 3$.7. $5\frac{1}{4}$ grains.8. $12\frac{1}{8}$ shillings.

CHAPTER VIII. EXAMINATION.

1. $15\frac{45}{1729}$ lbs.

9. 75° .

CHAPTER VIII. EXAMPLES.

1. $\frac{pV+p'V'}{U} \cdot \frac{1+at'}{1+at}$.

2. The air at greatest pressure.

3. $22^{\circ}\frac{58}{211}$.

4. $13^{\circ}\frac{53}{104}$.

5. The difference of the observed pressures.

CHAPTER IX.

2. If $l =$ Latus Rectum, $\omega^2 = \frac{g}{l}$.

3. Length submerged $= \frac{\sigma}{\rho}h + \frac{r^2\omega^2}{4g}$.

4. A paraboloid.

6. $\frac{1}{4} \frac{\pi a^4 \omega^2}{g}$.

3. $g\rho\pi a^2 \frac{a^4 \omega^2}{4g}$, $g\rho\pi a^2 \left(\frac{a^2 \omega^2}{4g} + h \right)$, and $g\rho\pi a h \left(h + \frac{\omega^2 a^2}{g} \right)$.

9. $g\rho\pi a^2 \left(\frac{a^2 \omega^2}{4g} + \frac{a}{3} \right)$.

CHAPTER X.

1. Inversely as the radii.
2. 3 : 2.
3. 80 lbs. on a square inch.
4. $r^3 : r^3$.
5. If $\frac{4}{3}\pi x^3$ be the volume of atmospheric air forced in,

$$(a^3 + x^3) \left\{ 1 - \frac{b^3}{c^3} \cdot \frac{b^3 - a^3}{c^3 - a^3} \right\} = b^3.$$

ANSWERS TO MISCELLANEOUS PROBLEMS.

3. First. The plane divides the axis in the ratio $1 : \sqrt[3]{2} - 1$,
Second. The plane bisects the axis.

Measuring x from the vertex, the equation is

$$x^3 - \frac{3h}{2}x^2 + \frac{h^3}{4} = 0,$$

one root of which is $\frac{h}{2}$.

4. The ratio is $4 : 3 \sin \alpha$.
5. The distances from B of the points of division are in the ratio $1 : \sqrt{2} : \sqrt{3} : \&c$.
6. 1st, $\frac{1}{3}$, 2nd, $\frac{13}{36}$ of its volume.
7. The densities are equal.
8. The pressures are either equal or in the ratio $1 : 3$.
9. $h \left(\frac{\sigma}{\rho} \right)^{\frac{2}{3}}$.
10. If Π = atmospheric pressure, the height above the base of the centre of the sphere is

$$\frac{\frac{2}{3}rW + h\pi r^2\Pi}{W + \pi r^2\Pi}.$$

11. 1 : 32000.

13. The inclination to the horizon = $\cos^{-1} 2 \frac{1-m}{1+m}$.

15. 2000 oz.

20. 60° .

23. The weight of the vessel must be at least $\frac{17}{7}$ (the weight of the fluid).

25. The volume must be $\frac{\rho}{\rho-\sigma} \pi r^2 \left(a - \frac{h}{3} \right)$, a being the depth of water, and h, r the height and radius of the cone.

29. If ρ, ρ' be the densities of the lower and upper liquids respectively, σ the density of the rod, and θ its inclination to the vertical,

$$\cos^2 \theta = \frac{c^2}{a^2} \frac{\rho - \rho'}{\sigma - \rho'}.$$

30. If a be the height of the cylinder, and h the height of the water-barometer, the length (x) of the cylinder occupied by air is given by the equation

$$x^2 + \left(h - \frac{3a}{4} \right) x = \frac{ah}{2}.$$

37. $\frac{2A \ln}{B}$.

45. The inclination of the radius vector to the surface is 60° .

46. The point lies in the central generating line, dividing it in the ratio 2 : 1.

47. In the first case the point divides the central generating line in the ratio 3 : 1; in the second it bisects the generating line.

48, and 49. See Example 8, page 55.

54. Resultant pressure : weight of liquid :: $\sqrt{13} : 2$, and its inclination to the horizon = $\tan^{-1} \frac{2}{3}$.

55, and 56. See Articles 51, 52, 53, and Example 2, page 47.

61. If ρ be the density of the upper and ρ' of the lower liquid, the pressures are in the ratio

$$4\rho : 3\rho + \rho'.$$

64. See Art. 160.

65. See Example 6, page 154.

66. See Appendix (6).

75. Remove the piston and replace it by an equivalent weight of water.

84. The pressures are in the inverse ratio of the radii.

88. If f be the acceleration with which the cone ascends, and $2a$ the vertical angle, the whole pressure : the weight of the liquid :: $f + g : g \sin a$.

90. $n^3 : 1 + at.$ 92. $\frac{mh}{h'}$.

97. ρ and ρ' being the densities and σ the density of the liquid, the axis of rotation divides the string in the ratio
 $\rho - \sigma : \rho' - \sigma$.

103. The tension is zero.

114. The whole impulse = $\rho\pi r h^2 v$, r being the radius and h the height of the fluid.

116. If $U = \pi r^3 \rho v$, the whole impulses are $4U$, U and $3U$.
 The resultant impulses are $\frac{4}{3}U$, $\frac{1}{3}U$, and $\frac{5}{3}U$.

NOTE ON ART. 84.

The value of α is very nearly the same for all gases, and moreover remains nearly the same for different pressures. M. Regnault has investigated the values of α for different substances; for instance, between 0° and 100° he finds the value of α for carbonic acid gas to be .003689. It has also been observed that the coefficients for two gases separate more from each other when the pressure is very much increased.

Regnault's results: values of α for

Air003665.
Hydrogen003667.
Azote003668.
Sulphuric Acid003669.
Hydrochloric Acid	.003681.
Cyanogen003682.
Carbonic Acid003689.

INDEX.

	PAGE		PAGE
ABSOLUTE Temperature.....	78	Heights by the Barometer...	80
Air a ponderable body	73	Homogeneous Atmosphere..	76
—in motion	176	Hopkins.....	195
Air-Pumps	103—105	Hydraulic Ram	112
Aneroid Barometer.....	88	Hydrometers	126—128
Archimedes.....	63, 118	Hydrostatic Balance	124
Balloon	59, 91	— Bellows	11
Barker's Mill	110	— Paradox	12
Barometer	74	Hygrometers.....	143
— gauge	106	Impulsive Action	177
Beats	193	Lintearia	163
Bourdon's Metallic Baro- meter	89	Liquid Films.....	171
Boyle's Law	76	Magdeburgh Hemispheres...90	
Bramah's Press.....	102	Manometers	108—110
Camel	52	Metacentre	57
Capillarity.....	166	Mixture of gases	134
Capillary Curve	163, 170	Monochord	194
Cecil's Lamp	56	Musical sounds	184
Centre of Pressure.....	38, 201	Nodes and loops.....	187—189
Coexistence and Interfer- ence	184	Pascal	90
Common surface of two liquids	31	Piezometer	111, 119
Condenser	107	Pumps	96—99
Contracted Vein	174	Rotating liquid.....	150
Density	18	Safety Valves	13
Des Cartes	91	Siphon	81
Dew	140, 142	Siphon gauge	107
Diatonic Scale	195	Sounds, velocity of,...	180, 184
Differential Thermometer...83		Specific gravity	20
Dilatation of Liquids	144	Stability	57
Diving-bell	94	Steady motion	175
Ebullition	145	Steam Engines	113—116
Elastica	163	Stereometer	129
Elasticity of air	69	Stevinus	64
Energy of Film	172	Tartini	193
Equilibrium of a floating body.....	50	Temperature of air.....	78
Equipotential Surfaces ...198		Tension	161
Freezing.....	144	Thermometer	71
Galileo	64	Torricelli.....	72, 173
Heat	138, 146	Unison and Harmony	192
		Vapours.....	136
		Virtual Velocities	14
		Water Barometer	75
		Waves	181

May 1883.

A CLASSIFIED LIST

OF

EDUCATIONAL WORKS

PUBLISHED BY

GEORGE BELL & SONS.

Full Catalogues will be sent post free on application.

BIBLIOTHECA CLASSICA.

A Series of Greek and Latin Authors, with English Notes, edited by eminent Scholars. 8vo.

Æschylus. By F. A. Paley, M.A. 18s.

Cicero's Orations. By G. Long, M.A. 4 vols. 16s., 14s., 16s., 18s.

Demosthenes. By R. Whiston, M.A. 2 vols. 16s. each.

Euripides. By F. A. Paley, M.A. 3 vols. 16s. each.

Homer. By F. A. Paley, M.A. Vol. I. 12s.; Vol. II. 14s.

Herodotus. By Rev. J. W. Blakesley, B.D. 2 vols. 32s.

Hesiod. By F. A. Paley, M.A. 10s. 6d.

Horace. By Rev. A. J. Maclean, M.A. 18s.

Juvenal and Persius. By Rev. A. J. Maclean, M.A. 12s.

Plato. By W. H. Thompson, D.D. 2 vols. 7s. 6d. each.

Sophocles. Vol. I. By Rev. F. H. Blaydes, M.A. 18s.

— Vol. II. Philoctetes. Electra. Ajax and Trachiniæ. By F. A. Paley, M.A. 12s.

Tacitus: The Annals. By the Rev. P. Frost. 15s.

Terence. By E. St. J. Parry, M.A. 18s.

Virgil. By J. Conington, M.A. 3 vols. 14s. each.

An Atlas of Classical Geography; Twenty-four Maps. By W. Hughes and George Long, M.A. New edition, with coloured Outlines. Imperial 8vo. 12s. 6d.

Uniform with above.

A Complete Latin Grammar. By J. W. Donaldson, D.D. 3rd Edition. 14s.

GRAMMAR-SCHOOL CLASSICS.

A Series of Greek and Latin Authors, with English Notes. Fcap. 8vo.

Cæsar: De Bello Gallico. By George Long, M.A. 5s. 6d.

— Books I.—III. For Junior Classes. By G. Long, M.A. 2s. 6d.

— Books IV. and V. in 1 vol. 1s. 6d.

Catullus, Tibullus, and Propertius. Selected Poems. With Life. By Rev. A. H. Wratlaw. 3s. 6d.

- Cicero:** De Senectute, De Amicitia, and Select Epistles. By George Long, M.A. 4s. 6d.
- Cornelius Nepos.** By Rev. J. F. Macmichael. 2s. 6d.
- Homer:** Iliad. Books I.—XII. By F. A. Paley, M.A. 6s. 6d.
- Horace.** With Life. By A. J. Maclean, M.A. 6s. 6d. [In 2 parts, 3s. 6d. each.]
- Juvenal:** Sixteen Satires. By H. Prior, M.A. 4s. 6d.
- Martial:** Select Epigrams. With Life. By F. A. Paley, M.A. 6s. 6d.
- Ovid:** the Fasti. By F. A. Paley, M.A. 5s.
- Sallust:** Catilina and Jugurtha. With Life. By G. Long, M.A. 5s.
- Tacitus:** Germania and Agricola. By Rev. P. Frost. 3s. 6d.
- Virgil:** Bucolics, Georgics, and Æneid, Books I.—IV. Abridged from Professor Conington's Edition. 5s. 6d.—Æneid, Books V.—XII. 5s. 6d. Also in 9 separate Volumes, 1s. 6d. each.
- Xenophon:** The Anabasis. With Life. By Rev. J. F. Macmichael. 5s. Also in 4 separate volumes, 1s. 6d. each.
- The Cyropædia. By G. M. Gorham, M.A. 6s.
- Memorabilia. By Percival Frost, M.A. 4s. 6d.
- A Grammar-School Atlas of Classical Geography,** containing Ten selected Maps. Imperial 8vo. 5s.

Uniform with the Series.

- The New Testament, in Greek.** With English Notes, &c. By Rev. J. F. Macmichael. 7s. 6d.

CAMBRIDGE GREEK AND LATIN TEXTS.

- Æschylus.** By F. A. Paley, M.A. 3s.
- Cæsar:** De Bello Gallico. By G. Long, M.A. 2s.
- Cicero:** De Senectute et de Amicitia, et Epistolæ Selectæ. By G. Long, M.A. 1s. 6d.
- Ciceronis Orationes.** Vol. I. (in Verrem.) By G. Long, M.A. 3s. 6d.
- Euripides.** By F. A. Paley, M.A. 3 vols. 3s. 6d. each.
- Herodotus.** By J. G. Blakesley, B.D. 2 vols. 7s.
- Homeri Ilias.** I.—XII. By F. A. Paley, M.A. 2s. 6d.
- Horatius.** By A. J. Maclean, M.A. 2s. 6d.
- Juvenal et Persius.** By A. J. Maclean, M.A. 1s. 6d.
- Lucretius.** By H. A. J. Munro, M.A. 2s. 6d.
- Sallusti Crispi Catilina et Jugurtha.** By G. Long, M.A. 1s. 6d.
- Sophocles.** By F. A. Paley, M.A. 3s. 6d.
- Terenti Comædiæ.** By W. Wagner, Ph.D. 3s.
- Thucydides.** By J. G. Donaldson, D.D. 2 vols. 7s.
- Virgilius.** By J. Conington, M.A. 3s. 6d.
- Xenophontis Expositio Cyri.** By J. F. Macmichael, B.A. 2s. 6d.
- Novum Testamentum Græcum.** By F. H. Scrivener, M.A. 4s. 6d. An edition with wide margin for notes, half bound, 12s.

CAMBRIDGE TEXTS WITH NOTES.

A Selection of the most usually read of the Greek and Latin Authors, Annotated for Schools. Fcap. 8vo. 1s. 6d. each, with exceptions.

Euripides. Alcestis.—Medea.—Hippolytus.—Hecuba.—Bacchæ.
Ion. 2s.—Orestes.—Phoenissæ.—Troades. By F. A. Paley, M.A.

Æschylus. Prometheus Vincetus.—Septem contra Thebas.—Agamemnon.—Persæ.—Eumenides. By F. A. Paley, M.A.

Sophocles. Œdipus Tyrannus.—Œdipus Coloneus.—Antigone.
By F. A. Paley, M.A.

Homer. Iliad. Book I. By F. A. Paley, M.A. 1s.

Terence. Andria.—Hauton Timorumenos.—Phormio.—Adelphoe.
By Professor Wagner, Ph.D.

Cicero's De Senectute, De Amicitia, and Epistolæ Selectæ. By
G. Long, M.A.

Ovid. Selections. By A. J. Maclean, M.A.
Others in preparation.

PUBLIC SCHOOL SERIES.

A Series of Classical Texts, annotated by well-known Scholars. Cr. 8vo.

Aristophanes. The Peace. By F. A. Paley, M.A. 4s. 6d.

—— The Acharnians. By F. A. Paley, M.A. 4s. 6d.

—— The Frogs. By F. A. Paley, M.A. 4s. 6d.

Cicero. The Letters to Atticus. Bk. I. By A. Pretor, M.A. 4s. 6d.

Demosthenes de Falsa Legatione. By R. Shilleto, M.A. 6s.

—— The Law of Leptines. By B. W. Beatson, M.A. 3s. 6d.

Plato. The Apology of Socrates and Crito. By W. Wagner, Ph.D.
7th Edition. 4s. 6d.

—— The Phædo. 6th Edition. By W. Wagner, Ph.D. 5s. 6d.

—— The Protagoras. 3rd Edition. By W. Wayte, M.A. 4s. 6d.

—— The Euthyphro. 2nd edition. By G. H. Wells, M.A. 3s.

—— The Euthydemus. By G. H. Wells, M.A. 4s.

—— The Republic. Books I. & II. By G. H. Wells, M.A. 5s. 6d.

Plautus. The Aulularia. By W. Wagner, Ph.D. 2nd edition. 4s. 6d.

—— Trinummus. By W. Wagner, Ph.D. 2nd edition. 4s. 6d.

—— The Menæchmei. By W. Wagner, Ph.D. 4s. 6d.

Sophocles Trachiniæ. By A. Pretor, M.A. 4s. 6d.

Terence. By W. Wagner, Ph.D. 10s. 6d.

Theocritus. By F. A. Paley, M.A. 4s. 6d.

Others in preparation.

CRITICAL AND ANNOTATED EDITIONS.

Ætina. By H. A. J. Munro, M.A. 3s. 6d.

Aristophanis Comœdiæ. By H. A. Holden, LL.D. 8vo. 2 vols.
23s. 6d. Plays sold separately.

—— Pax. By F. A. Paley, M.A. Fcap. 8vo. 4s. 6d.

Catullus. By H. A. J. Munro, M.A. 7s. 6d.

Corpus Poetarum Latinorum. Edited by Walker. 1 vol. 8vo. 18s.

Horace. Quinti Horatii Flacci Opera. By H. A. J. Munro, M.A.
Large 8vo. 1l. 1s.

Livy. The first five Books. By J. Prendeville. 12mo. roan, 5s.
Or Books I.-III. 3s. 6d. IV. and V. 3s. 6d.

- Lucretius.** Titi Lucretii Cari de Rerum Natura Libri Sex. With a Translation and Notes. By H. A. J. Munro, M.A. 2 vols. 8vo. Vol. I. Text. (New Edition, Preparing.) Vol. II. Translation. (Sold separately.)
- Ovid.** P. Ovidii Nasonis Heroides XIV. By A. Palmer, M.A. 8vo. 6s.
- Propertius.** Sex Aurelii Propertii Carmina. By F. A. Paley, M.A. 8vo. Cloth, 9s.
- Sex. Propertii Elegiarum.** Lib. IV. By A. Palmer. Fcap. 8vo. 5s.
- Sophocles.** The Ajax. By C. E. Palmer, M.A. 4s. 6d.
- Thucydides.** The History of the Peloponnesian War. By Richard Shilleto, M.A. Book I. 8vo. 6s. 6d. Book II. 8vo. 5s. 6d.

LATIN AND GREEK CLASS-BOOKS.

- Auxilia Latina.** A Series of Progressive Latin Exercises. By M. J. B. Baddeley, M.A. Fcap. 8vo. Part I. Accidence. 2nd Edition, revised. 2s. Part II. 4th Edition, revised. 2s. Key to Part II. 2s. 6d.
- Latin Prose Lessons.** By Prof. Church, M.A. 6th Edit. Fcap. 8vo. 2s. 6d.
- Latin Exercises and Grammar Papers.** By T. Collins, M.A. 3rd Edition. Fcap. 8vo. 2s. 6d.
- Unseen Papers in Latin Prose and Verse.** With Examination Questions. By T. Collins, M.A. 2nd Edition. Fcap. 8vo. 2s. 6d.
- in Greek Prose and Verse. With Examination Questions. By T. Collins, M.A. Fcap. 8vo. 3s.
- Analytical Latin Exercises.** By C. P. Mason, B.A. 3rd Edit. 3s. 6d.
- Latin Mood Construction, Outlines of.** With Exercises. By the Rev. G. E. C. Casey, M.A., F.L.S., F.G.S. Small post 8vo. 1s. 6d. Latin of the Exercises. 1s. 6d.
- Scala Græca: a Series of Elementary Greek Exercises.** By Rev. J. W. Davis, M.A., and R. W. Baddeley, M.A. 3rd Edition. Fcap. 8vo. 2s. 6d.
- Greek Verse Composition.** By G. Preston, M.A. Crown 8vo. 4s. 6d.
- Greek Particles and their Combinations according to Attic Usage.** A Short Treatise. By F. A. Paley, M.A. 2s. 6d.
- BY THE REV. P. FROST, M.A., ST. JOHN'S COLLEGE, CAMBRIDGE.
- Ecologæ Latinæ; or, First Latin Reading-Book, with English Notes and a Dictionary.** New Edition. Fcap. 8vo. 2s. 6d.
- Materials for Latin Prose Composition.** New Edition. Fcap. 8vo. 2s. 6d. Key, 4s.
- A Latin Verse-Book.** An Introductory Work on Hexameters and Pentameters. New Edition. Fcap. 8vo. 3s. Key, 5s.
- Analecta Græca Minora, with Introductory Sentences, English Notes, and a Dictionary.** New Edition. Fcap. 8vo. 3s. 6d.
- Materials for Greek Prose Composition.** New Edit. Fcap. 8vo. 3s. 6d. Key, 5s.
- Florilegium Poeticum.** Elegiac Extracts from Ovid and Tibullus. New Edition. With Notes. Fcap. 8vo. 3s.
- BY THE REV. F. E. GRETTON.
- A First Cheque-book for Latin Verse-makers.** 1s. 6d.
- A Latin Version for Masters.** 2s. 6d.
- Reddenda; or Passages with Parallel Hints for Translation into Latin Prose and Verse.** Crown 8vo. 4s. 6d.
- Reddenda Reddita** (*see next page*).

Anthologia Græca. A Selection of Choice Greek Poetry, with Notes. By F. St. John Thackeray. 4th and Cheaper Edition. 16mo. 4s. 6d.

Anthologia Latina. A Selection of Choice Latin Poetry, from Nævius to Boëthius, with Notes. By Rev. F. St. John Thackeray. Revised and Cheaper Edition. 16mo. 4s. 6d.

By H. A. HOLDEN, LL.D.

Follorum Silvula. Part I. Passages for Translation into Latin Elegiac and Heroic Verse. 9th Edition. Post 8vo. 7s. 6d.

—— Part II. Select Passages for Translation into Latin Lyric and Comic Iambic Verse. 3rd Edition. Post 8vo. 5s.

—— Part III. Select Passages for Translation into Greek Verse. 3rd Edition. Post 8vo. 8s.

Folia Silvulæ, sive Eclogæ Poetarum Anglicorum in Latinum et Græcūm conversæ. 8vo. Vol. I. 10s. 6d. Vol. II. 12s.

Follorum Centuriæ. Select Passages for Translation into Latin and Greek Prose. 8th Edition. Post 8vo. 8s.

TRANSLATIONS, SELECTIONS, &c.

* * Many of the following books are well adapted for School Prizes.

Æschylus. Translated into English Prose by F. A. Paley, M.A. 2nd Edition. 8vo. 7s. 6d.

—— Translated into English Verse by Anna Swanwick. Post 8vo. 5s.

Horace. The Odes and Carmen Sæculare. In English Verse by J. Conington, M.A. 8th edition. Fcap. 8vo. 5s. 6d.

—— The Satires and Epistles. In English Verse by J. Conington, M.A. 5th edition. 6s. 6d.

—— Illustrated from Antique Gems by C. W. King, M.A. The text revised with Introduction by H. A. J. Munro, M.A. Large 8vo. 1l. 1s.

Horace's Odes. Englished and Imitated by various hands. Edited by C. W. F. Cooper. Crown 8vo. 6s. 6d.

Lusus Intercisi. Verses, Translated and Original, by H. J. Hodgson, M.A., formerly Fellow of Trinity College, Cambridge.

[Immediately.]

Propertius. Verse Translations from Book V., with revised Latin Text. By F. A. Paley, M.A. Fcap. 8vo. 3s.

Plato. Gorgias. Translated by E. M. Cope, M.A. 8vo. 7s.

—— Philebus. Translated by F. A. Paley, M.A. Small 8vo. 4s.

—— Theætetus. Translated by F. A. Paley, M.A. Small 8vo. 4s.

—— Analysis and Index of the Dialogues. By Dr. Day. Post 8vo. 5s.

Reddenda Reddita : Passages from English Poetry, with a Latin Verse Translation. By F. E. Gretton. Crown 8vo. 6s.

Sabrinæ Corolla in hortulis Regiæ Scholæ Salopiensis contexuerunt tres viri floribus legendis. Editio tertia. 8vo. 8s. 6d.

Theocritus. In English Verse, by C. S. Calverley, M.A. New Edition, revised. Crown 8vo. 7s. 6d.

Translations into English and Latin. By C. S. Calverley, M.A. Post 8vo. 7s. 6d.

—— into Greek and Latin Verse. By R. C. Jebb. 4to. cloth gilt. 10s. 6d.

Between Whiles. Translations by B. H. Kennedy. 2nd Edition. revised. Crown 8vo. 6s.

REFERENCE VOLUMES.

- A Latin Grammar.** By Albert Harkness. Post 8vo. 6s.
 ——— By T. H. Key, M.A. 6th Thousand. Post 8vo. 8s.
- A Short Latin Grammar for Schools.** By T. H. Key, M.A.,
 F.R.S. 14th Edition. Post 8vo. 3s. 6d.
- A Guide to the Choice of Classical Books.** By J. B. Mayor, M.A.
 Revised Edition. Crown 8vo. 3s.
- The Theatre of the Greeks.** By J. W. Donaldson, D.D. 8th
 Edition. Post 8vo. 5s.
- Keightley's Mythology of Greece and Italy.** 4th Edition. 5s.
- A Dictionary of Latin and Greek Quotations.** By H. T. Riley.
 Post 8vo. 5s. With Index Verborum, 6s.
- A History of Roman Literature.** By W. S. Teuffel, Professor at
 the University of Tübingen. By W. Wagner, Ph.D. 2 vols. Demy 8vo. 21s.
- Student's Guide to the University of Cambridge.** 4th Edition
 revised. Fcap. 8vo. Part 1, 2s. 6d.; Parts 2 to 6, 1s. each.
-

CLASSICAL TABLES.

- Latin Accidence.** By the Rev. P. Frost, M.A. 1s.
- Latin Versification.** 1s.
- Notabilia Quædam; or the Principal Tenses of most of the**
 Irregular Greek Verbs and Elementary Greek, Latin, and French Con-
 struction. New Edition. 1s.
- Richmond Rules for the Ovidian Distich, &c.** By J. Tate,
 M.A. 1s.
- The Principles of Latin Syntax.** 1s.
- Greek Verbs.** A Catalogue of Verbs, Irregular and Defective; their
 leading formations, tenses, and inflexions, with Paradigms for conjugation,
 Rules for formation of tenses, &c. &c. By J. S. Baird, T.C.D. 2s. 6d.
- Greek Accents (Notes on).** By A. Barry, D.D. New Edition. 1s.
- Homeric Dialect.** Its Leading Forms and Peculiarities. By J. S.
 Baird, T.C.D. New Edition, by W. G. Rutherford. 1s.
- Greek Accidence.** By the Rev. P. Frost, M.A. New Edition. 1s.
-

CAMBRIDGE MATHEMATICAL SERIES.

- Algebra.** Choice and Chance. By W. A. Whitworth, M.A. 3rd
 Edition. 6s.
- Euclid.** Exercises on Euclid and in Modern Geometry. By
 J. McDowell, M.A. 3rd Edition. 6s.
- Trigonometry.** Plane. By Rev. T. Vyvyan, M.A. 3s. 6d.
- Conics.** The Geometry of. 3rd Edition. By C. Taylor, D.D. 4s. 6d.
- Solid Geometry.** By W. S. Aldis, M.A. 3rd Edition. 6s.
- Rigid Dynamics.** By W. S. Aldis, M.A. 4s.
- Elementary Dynamics.** By W. Garnett, M.A. 3rd Edition. 6s.

Heat. An Elementary Treatise. By W. Garnett, M.A. 2nd Edit.
3s. 6d.

Hydromechanics. By W. H. Besant, M.A., F.R.S. 4th Edition.
Part I. Hydrostatics. 5s.

Mechanics. Problems in Elementary. By W. Walton, M.A. 6s.

CAMBRIDGE SCHOOL AND COLLEGE TEXT-BOOKS.

*A Series of Elementary Treatises for the use of Students in the
Universities, Schools, and Candidates for the Public
Examinations. Fcap. 8vo.*

Arithmetic. By Rev. C. Elsee, M.A. Fcap. 8vo. 10th Edit. 3s. 6d.

Algebra. By the Rev. C. Elsee, M.A. 6th Edit. 4s.

Arithmetic. By A. Wrigley, M.A. 3s. 6d.

——— A Progressive Course of Examples. With Answers. By
J. Watson, M.A. 5th Edition. 2s. 6d.

Algebra. Progressive Course of Examples. By Rev. W. F.
M'Michael, M.A., and R. Prowde Smith, M.A. 2nd Edition. 3s. 6d. With
Answers. 4s. 6d.

Plane Astronomy, An Introduction to. By P. T. Main, M.A.
4th Edition. 4s.

Conic Sections treated Geometrically. By W. H. Besant, M.A.
4th Edition. 4s. 6d. Solution to the Examples. 4s.

Elementary Conic Sections treated Geometrically. By W. H.
Besant, M.A. [*In the press.*]

Statics, Elementary. By Rev. H. Goodwin, D.D. 2nd Edit. 3s.

Hydrostatics, Elementary. By W. H. Besant, M.A. 10th Edit. 4s.

Mensuration, An Elementary Treatise on. By B. T. Moore, M.A. 6s.

Newton's Principia, The First Three Sections of, with an Appen-
dix; and the Ninth and Eleventh Sections. By J. H. Evans, M.A. 5th
Edition, by P. T. Main, M.A. 4s.

Trigonometry, Elementary. By T. P. Hudson, M.A. 3s. 6d.

Optics, Geometrical. With Answers. By W. S. Aldis, M.A. 3s. 6d.

Analytical Geometry for Schools. By T. G. Vyvyan. 3rd Edit. 4s. 6d.

Greek Testament, Companion to the. By A. C. Barrett, A.M.
4th Edition, revised. Fcap. 8vo. 5s.

Book of Common Prayer, An Historical and Explanatory Treatise
on the. By W. G. Humphry, B.D. 6th Edition. Fcap. 8vo. 4s. 6d.

Music, Text-book of. By H. C. Banister. 10th Edit. revised. 5s.

——— Concise History of. By Rev. H. G. Bonavia Hunt, B. Mus.
Oxon. 6th Edition revised. 3s. 6d.

ARITHMETIC AND ALGEBRA.

See foregoing Series.

GEOMETRY AND EUCLID.

- Euclid. The First Two Books explained to Beginners. By C. P. Mason, B.A. 2nd Edition. Fcap. 8vo. 2s. 6d.
- The Enunciations and Figures to Euclid's Elements. By Rev. J. Brasse, D.D. New Edition. Fcap. 8vo. 1s. On Cards, in case, 5s. 6d. Without the Figures, 6d.
- Exercises on Euclid and in Modern Geometry. By J. McDowell, B.A. Crown 8vo. 3rd Edition revised. 6s.
- Geometrical Conic Sections. By W. H. Besant, M.A. 4th Edit. 4s. 6d. Solution to the Examples. 4s.
- Elementary Geometrical Conic Sections. By W. H. Besant, M.A. *[In the press.]*
- Elementary Geometry of Conics. By C. Taylor, D.D. 3rd Edit. 8vo. 4s. 6d.
- An Introduction to Ancient and Modern Geometry of Conics. By C. Taylor, M.A. 8vo. 15s.
- Solutions of Geometrical Problems, proposed at St. John's College from 1830 to 1846. By T. Gaskin, M.A. 8vo. 12s.

TRIGONOMETRY.

- Trigonometry, Introduction to Plane. By Rev. T. G. Vyvyan, Charterhouse. Cr. 8vo. 3s. 6d.
- Elementary Trigonometry. By T. P. Hudson, M.A. 3s. 6d.
- An Elementary Treatise on Mensuration. By B. T. Moore, M.A. 5s.

ANALYTICAL GEOMETRY AND DIFFERENTIAL CALCULUS.

- An Introduction to Analytical Plane Geometry. By W. P. Turnbull, M.A. 8vo. 12s.
- Problems on the Principles of Plane Co-ordinate Geometry. By W. Walton, M.A. 8vo. 16s.
- Trilinear Co-ordinates, and Modern Analytical Geometry of Two Dimensions. By W. A. Whitworth, M.A. 8vo. 16s.
- An Elementary Treatise on Solid Geometry. By W. S. Aldis, M.A. 2nd Edition revised. 8vo. 8s.
- Elementary Treatise on the Differential Calculus. By M. O'Brien, M.A. 8vo. 10s. 6d.
- Elliptic Functions, Elementary Treatise on. By A. Cayley, M.A. Demy 8vo. 15s.

MECHANICS & NATURAL PHILOSOPHY.

- Statics, Elementary. By H. Goodwin, D.D. Fcap. 8vo. 2nd Edition. 3s.
- Dynamics, A Treatise on Elementary. By W. Garnett, M.A. 3rd Edition. Crown 8vo. 6s.
- Elementary Mechanics, Problems in. By W. Walton, M.A. New Edition. Crown 8vo. 6s.

- Theoretical Mechanics**, Problems in. By W. Walton. 2nd Edit. revised and enlarged. Demy 8vo. 16s.
- Hydrostatics**. By W. H. Besant, M.A. Fcap. 8vo. 10th Edition. 4s.
- Hydromechanics**, A Treatise on. By W. H. Besant, M.A., F.R.S. 8vo. 4th Edition, revised. Part I. Hydrostatics. 5s.
- Dynamics of a Particle**, A Treatise on the. By W. H. Besant, M.A. [Preparing.]
- Optics**, Geometrical. By W. S. Aldis, M.A. Fcap. 8vo. 3s. 6d.
- Double Refraction**, A Chapter on Fresnel's Theory of. By W. S. Aldis, M.A. 8vo. 2s.
- Heat**, An Elementary Treatise on. By W. Garnett, M.A. Crown 8vo. 2nd Edition revised. 3s. 6d.
- Newton's Principia**, The First Three Sections of, with an Appendix; and the Ninth and Eleventh Sections. By J. H. Evans, M.A. 5th Edition. Edited by P. T. Main, M.A. 4s.
- Astronomy**, An Introduction to Plane. By P. T. Main, M.A. Fcap. 8vo. cloth. 4s.
- Astronomy**, Practical and Spherical. By R. Main, M.A. 8vo. 14s.
- Astronomy**, Elementary Chapters on, from the 'Astronomie Physique' of Biot. By H. Goodwin, D.D. 8vo. 3s. 6d.
- Pure Mathematics and Natural Philosophy**, A Compendium of Facts and Formulæ in. By G. R. Sinalley. 2nd Edition, revised by J. McDowell, M.A. Fcap. 8vo. 3s. 6d.
- Elementary Course of Mathematics**. By H. Goodwin, D.D. 6th Edition. 8vo. 16s.
- Problems and Examples**, adapted to the 'Elementary Course of Mathematics.' 3rd Edition. 8vo. 5s.
- Solutions of Goodwin's Collection of Problems and Examples**. By W. W. Hutt, M.A. 3rd Edition, revised and enlarged. 8vo. 9s.
- Pure Mathematics**, Elementary Examples in. By J. Taylor. 8vo. 7s. 6d.
- Mechanics of Construction**. With numerous Examples. By S. Fenwick, F.R.A.S. 8vo. 12s.
- Pure and Applied Calculation**, Notes on the Principles of. By Rev. J. Challis, M.A. Demy 8vo. 15s.
- Physics**, The Mathematical Principle of. By Rev. J. Challis, M.A. Demy 8vo. 5s.

TECHNOLOGICAL HANDBOOKS.

Edited by H. TRUEMAN WOOD, Secretary of the
Society of Arts.

1. **Dyeing and Tissue Printing**. By W. Crookes, F.R.S. 5s.
2. **Glass Manufacture**. By Henry Chance, M.A.; H. J. Powell, B.A.; and H. G. Harris. 3s. 6d.
3. **Cotton Manufacture**. By Richard Marsden, Esq., of Manchester. [In the press.]
4. **Telegraphs and Telephones**. By W. H. Preece, F.R.S. [Preparing.]

HISTORY, TOPOGRAPHY, &c.

- Rome and the Campagna.** By R. Burn, M.A. With 85 Engravings and 26 Maps and Plans. With Appendix. 4to. 3l. 3s.
- Old Rome.** A Handbook for Travellers. By R. Burn, M.A. With Maps and Plans. Demy 8vo. 10s. 6d.
- Modern Europe.** By Dr. T. H. Dyer. 2nd Edition, revised and continued. 5 vols. Demy 8vo. 2l. 12s. 6d.
- The History of the Kings of Rome.** By Dr. T. H. Dyer. 8vo. 16s.
- The History of Pompeii: its Buildings and Antiquities.** By T. H. Dyer. 3rd Edition, brought down to 1874. Post 8vo. 7s. 6d.
- Ancient Athens: its History, Topography, and Remains.** By T. H. Dyer. Super-royal 8vo. Cloth. 1l. 5s.
- The Decline of the Roman Republic.** By G. Long. 5 vols. 8vo. 14s. each.
- A History of England during the Early and Middle Ages.** By C. H. Pearson, M.A. 2nd Edition revised and enlarged. 8vo. Vol. I. 16s. Vol. II. 14s.
- Historical Maps of England.** By C. H. Pearson. Folio. 2nd Edition revised. 3l. 6d.
- History of England, 1800-15.** By Harriet Martineau, with new and copious Index. 1 vol. 3s. 6d.
- History of the Thirty Years' Peace, 1815-46.** By Harriet Martineau. 4 vols. 3s. 6d. each.
- A Practical Synopsis of English History.** By A. Bowes. 4th Edition. 8vo. 2s.
- Student's Text-Book of English and General History.** By D. Beale. Crown 8vo. 2s. 6d.
- Lives of the Queens of England.** By A. Strickland. Library Edition, 8 vols. 7s. 6d. each. Cheaper Edition, 6 vols. 5s. each. Abridged Edition, 1 vol. 6s. 6d.
- Eginhard's Life of Karl the Great (Charlemagne).** Translated with Notes, by W. Glaister, M.A., B.C.L. Crown 8vo. 4s. 6d.
- Outlines of Indian History.** By A. W. Hughes. Small post 8vo. 3s. 6d.
- The Elements of General History.** By Prof. Tytler. New Edition, brought down to 1874. Small post 8vo. 3s. 6d.

ATLASES.

- An Atlas of Classical Geography.** 24 Maps. By W. Hughes and G. Long, M.A. New Edition. Imperial 8vo. 12s. 6d.
- A Grammar-School Atlas of Classical Geography.** Ten Maps selected from the above. New Edition. Imperial 8vo. 5s.
- First Classical Maps.** By the Rev. J. Tate, M.A. 3rd Edition. Imperial 8vo. 7s. 6d.
- Standard Library Atlas of Classical Geography.** Imp. 8vo. 7s. 6d.

PHILOLOGY.

WEBSTER'S DICTIONARY OF THE ENGLISH LANGUAGE. With Dr. Mahn's Etymology. 1 vol., 1628 pages, 3090 Illustrations. 21s. With Appendices and 70 additional pages of Illustrations, 1919 pages, 31s. 6d.

'THE BEST PRACTICAL ENGLISH DICTIONARY EXTANT.'—*Quarterly Review*, 1873.

Prospectuses, with specimen pages, post free on application.

New Dictionary of the English Language. Combining Explanation with Etymology, and copiously illustrated by Quotations from the best Authorities. By Dr. Richardson. New Edition, with a Supplement. 2 vols. 4to. 4l. 14s. 6d.; half russia, 5l. 15s. 6d.; russia, 6l. 12s. Supplement separately. 4to. 12s.

An 8vo. Edit. without the Quotations, 15s.; half russia, 20s.; russia, 24s.

Supplementary English Glossary. Containing 12,000 Words and Meanings occurring in English Literature, not found in any other Dictionary. By T. L. O. Davies. Demy 8vo. 16s.

Folk-Etymology. A Dictionary of Words perverted in Form or Meaning by False Derivation or Mistaken Analogy. By Rev. A. S. Palmer. Demy 8vo. 21s.

Brief History of the English Language. By Prof. James Hadley, LL.D., Yale College. Fcap. 8vo. 1s.

The Elements of the English Language. By E. Adams, Ph.D. 15th Edition. Post 8vo. 4s. 6d.

Philological Essays. By T. H. Key, M.A., F.R.S. 8vo. 10s. 6d.

Language, its Origin and Development. By T. H. Key, M.A., F.R.S. 8vo. 14s.

Synonyms and Antonyms of the English Language. By Archdeacon Smith. 2nd Edition; Post 8vo. 5s.

Synonyms Discriminated. By Archdeacon Smith. Demy 8vo. 16s.

Bible English. By T. L. O. Davies. 5s.

The Queen's English. A Manual of Idiom and Usage. By the late Dean Alford 6th Edition. Fcap. 8vo. 5s.

Etymological Glossary of nearly 2500 English Words derived from the Greek. By the Rev. E. J. Boyce. Fcap. 8vo. 3s. 6d.

A Syriac Grammar. By G. Phillips, D.D. 3rd Edition, enlarged. 8vo. 7s. 6d.

A Grammar of the Arabic Language. By Rev. W. J. Beaumont, M.A. 12mo. 7s.

DIVINITY, MORAL PHILOSOPHY, &c.

Novum Testamentum Græcum, Textus Stephanici, 1550. By F. H. Scrivener, A.M., LL.D., D.C.L. New Edition. 16mo. 4s. 6d. Also on Writing Paper, with Wide Margin. Half-bound. 12s.

By the same Author.

Codex Bezae Cantabrigiæ. 4to. 26s.

A Full Collation of the Codex Sinaiticus with the Received Text of the New Testament, with Critical Introduction. 2nd Edition, revised. Fcap. 8vo. 5s.

A Plain Introduction to the Criticism of the New Testament. With Forty Facsimiles from Ancient Manuscripts. 3rd Edition. 8vo.

[In the press.]

Six Lectures on the Text of the New Testament. For English Readers. Crown 8vo. 6s.

- The New Testament for English Readers.** By the late H. Alford, D.D. Vol. I. Part I. 3rd Edit. 12s. Vol. I. Part II. 2nd Edit. 10s. 6d. Vol. II. Part I. 2nd Edit. 16s. Vol. II. Part II. 2nd Edit. 16s.
- The Greek Testament.** By the late H. Alford, D.D. Vol. I. 6th Edit. 1l. 8s. Vol. II. 6th Edit. 1l. 4s. Vol. III. 5th Edit. 18s. Vol. IV. Part I. 4th Edit. 18s. Vol. IV. Part II. 4th Edit. 14s. Vol. IV. 1l. 12s.
- Companion to the Greek Testament.** By A. C. Barrett, M.A. 4th Edition, revised. Fcap. 8vo. 5s.
- The Book of Psalms.** A New Translation, with Introductions, &c. By the Very Rev. J. J. Stewart Perowne, D.D. 8vo. Vol. I. 5th Edition, 18s. Vol. II. 5th Edit. 16s.
- Abridged for Schools. 3rd Edition. Crown 8vo. 10s. 6d.
- History of the Articles of Religion.** By C. H. Hardwick. 3rd Edition. Post 8vo. 5s.
- History of the Creeds.** By J. R. Lumby, D.D. 2nd Edition. Crown 8vo. 7s. 6d.
- Pearson on the Creed.** Carefully printed from an early edition. With Analysis and Index by E. Walford, M.A. Post 8vo. 5s.
- An Historical and Explanatory Treatise on the Book of Common Prayer.** By Rev. W. G. Humphry, B.D. 6th Edition, enlarged. Small post 8vo. 4s. 6d.
- The New Table of Lessons Explained.** By Rev. W. G. Humphry, B.D. Fcap. 1s. 6d.
- A Commentary on the Gospels for the Sundays and other Holy Days of the Christian Year.** By Rev. W. Denton, A.M. New Edition. 3 vols. 8vo. 54s. Sold separately.
- Commentary on the Epistles for the Sundays and other Holy Days of the Christian Year.** By Rev. W. Denton, A.M. 2 vols. 36s. Sold separately.
- Commentary on the Acts.** By Rev. W. Denton, A.M. Vol. I. 8vo. 18s. Vol. II. 14s.
- Notes on the Catechism.** By Rev. Canon Barry, D.D. 6th Edit. Fcap. 2s.
- Catechetical Hints and Helps.** By Rev. E. J. Boyce, M.A. 4th Edition, revised. Fcap. 2s. 6d.
- Examination Papers on Religious Instruction.** By Rev. E. J. Boyce. Sewed. 1s. 6d.
- Church Teaching for the Church's Children.** An Exposition of the Catechism. By the Rev. F. W. Harper. Sq. fcap. 2s.
- The Winton Church Catechist.** Questions and Answers on the Teaching of the Church Catechism. By the late Rev. J. S. B. Monsell, LL.D. 3rd Edition. Cloth, 3s.; or in Four Parts, sewed.
- The Church Teacher's Manual of Christian Instruction.** By Rev. M. F. Sadler. 24th Thousand. 2s. 6d.
- Short Explanation of the Epistles and Gospels of the Christian Year, with Questions.** Royal 32mo. 2s. 6d.; calf, 4s. 6d.
- Butler's Analogy of Religion; with Introduction and Index** by Rev. Dr. Steere. New Edition. Fcap. 3s. 6d.
- **Three Sermons on Human Nature, and Dissertation on Virtue.** By W. Whewell, D.D. 4th Edition. Fcap. 8vo. 2s. 6d.

- Lectures on the History of Moral Philosophy in England.** By W. Whewell, D.D. Crown 8vo. 8s.
- Kent's Commentary on International Law.** By J. T. Abdy, LL.D. New and Cheap Edition. Crown 8vo. 10s. 6d.
- A Manual of the Roman Civil Law.** By G. Leapingwell, LL.D. 8vo. 12s.

FOREIGN CLASSICS.

A series for use in Schools, with English Notes, grammatical and explanatory, and renderings of difficult idiomatic expressions.
Fcap. 8vo.

- Schiller's Wallenstein.** By Dr. A. Buchheim. 3rd Edit. 6s. 6d.
Or the Lager and Piccolomini, 3s. 6d. Wallenstein's Tod, 3s. 6d.
- **Maid of Orleans.** By Dr. W. Wagner. 3s. 6d.
- **Maria Stuart.** By V. Kastner. 3s.
- Goethe's Hermann and Dorothea.** By E. Bell, M.A., and E. Wölfel. 2s. 6d.
- German Ballads, from Uhland, Goethe, and Schiller.** By C. L. Bielefeld. 3rd Edition. 3s. 6d.
- Charles XII., par Voltaire.** By L. Direy. 4th Edition. 3s. 6d.
- Aventures de Télémaque, par Fénelon.** By C. J. Delille. 2nd Edition. 4s. 6d.
- Select Fables of La Fontaine.** By F. E. A. Gasc. 14th Edition. 3s.
- Picciola, by X. B. Saintine.** By Dr. Dubuc. 11th Thousand. 3s. 6d.

FRENCH CLASS-BOOKS.

- Twenty Lessons in French.** With Vocabulary, giving the Pronunciation. By W. Brebner. Post 8vo. 4s.
- French Grammar for Public Schools.** By Rev. A. C. Clapin, M.A. Fcap. 8vo. 9th Edition, revised. 2s. 6d.
- French Primer.** By Rev. A. C. Clapin, M.A. Fcap. 8vo. 4th Edit. 1s.
- Primer of French Philology.** By Rev. A. C. Clapin. Fcap. 8vo. 1s.
- Le Nouveau Trésor; or, French Student's Companion.** By M. E. S. 16th Edition. Fcap. 8vo. 3s. 6d.

F. E. A. GASC'S FRENCH COURSE.

- First French Book.** Fcap. 8vo. 76th Thousand. 1s. 6d.
- Second French Book.** 42nd Thousand. Fcap. 8vo. 2s. 6d.
- Key to First and Second French Books.** Fcap. 8vo. 3s. 6d.
- French Fables for Beginners, in Prose, with Index.** 15th Thousand. 12mo. 2s.
- Select Fables of La Fontaine.** New Edition. Fcap. 8vo. 3s.
- Histoires Amusantes et Instructives.** With Notes. 14th Thousand. Fcap. 8vo. 2s. 6d.

- Practical Guide to Modern French Conversation.** 12th Thousand. Fcap. 8vo. 2s. 6d.
- French Poetry for the Young.** With Notes. 4th Edition. Fcap. 8vo. 2s.
- Materials for French Prose Composition; or, Selections from the best English Prose Writers.** 16th Thousand. Fcap. 8vo. 4s. 6d. Key, 6s.
- Prosateurs Contemporains.** With Notes. 8vo. 6th Edition, revised. 5s.
- Le Petit Compagnon; a French Talk-Book for Little Children.** 11th Thousand. 16mo. 2s. 6d.
- An Improved Modern Pocket Dictionary of the French and English Languages.** 30th Thousand, with Additions. 16mo. Cloth. 4s. Also in 2 vols., in neat leatherette, 5s.
- Modern French-English and English-French Dictionary.** 2nd Edition, revised. In 1 vol. 12s. 6d. (formerly 2 vols. 25s.)

GOMBERT'S FRENCH DRAMA.

Being a Selection of the best Tragedies and Comedies of Molière, Racine, Corneille, and Voltaire. With Arguments and Notes by A. Gombert. New Edition, revised by F. E. A. Gasc. Fcap. 8vo. 1s. each; sewed, 6d.

CONTENTS.

MOLIERE:—Le Misanthrope. L'Avare. Le Bourgeois Gentilhomme. Le Tartuffe. Le Malade Imaginaire. Les Femmes Savantes. Les Fourberies de Scapin. Les Précieuses Ridicules. L'Ecole des Femmes. L'Ecole des Maris. Le Médecin malgré Lui.

RACINE:—Phédre. Esther. Athalie. Iphigénie. Les Plaideurs. La Thébaïde; ou, Les Frères Ennemis. Andromaque. Britannicus.

P. CORNEILLE:—Le Cid. Horace. Cinna. Polyucte.

VOLTAIRE:—Zaïre.

GERMAN CLASS-BOOKS.

- Materials for German Prose Composition.** By Dr Buchheim. 7th Edition Fcap. 4s. 6d. Key, 3s.
- Wortfolge, or Rules and Exercises on the Order of Words in German Sentences.** By Dr. F. Stock. 1s. 6d.
- A German Grammar for Public Schools.** By the Rev. A. C. Clapin and F. Holl Müller. 2nd Edition. Fcap. 2s. 6d.
- Kotzebue's Der Gefangene.** With Notes by Dr. W. Stromberg. 1s.

ENGLISH CLASS-BOOKS.

- A Brief History of the English Language.** By Prof. Jas. Hadley, LL.D., of Yale College. Fcap. 8vo. 1s.
- The Elements of the English Language.** By E. Adams, Ph.D. 18th Edition. Post 8vo. 4s. 6d.
- The Rudiments of English Grammar and Analysis.** By E. Adams, Ph.D. 8th Edition. Fcap. 8vo. 2s.

By C. P. MASON, Fellow of Univ. Coll. London.

- First Notions of Grammar for Young Learners.** Fcap. 8vo.
10th Thousand. Cloth. 8d.
- First Steps in English Grammar for Junior Classes.** Demy
18mo. 32nd Thousand. 1s.
- Outlines of English Grammar for the use of Junior Classes.**
31st Thousand. Crown 8vo. 2s.
- English Grammar, including the Principles of Grammatical
Analysis.** 25th Edition. 86th Thousand. Crown 8vo. 3s. 6d.
- A Shorter English Grammar, with copious Exercises.** 8th Thou-
sand. Crown 8vo. 3s. 6d.
- English Grammar Practice, being the Exercises separately.** 1s.

- Practical Hints on Teaching.** By Rev. J. Menet, M.A. 5th Edit.
Crown 8vo. cloth, 2s. 6d.; paper, 2s.
- How to Earn the Merit Grant.** By H. Major. [*In the press.*]
- Test Lessons in Dictation.** 2nd Edition. Paper cover, 1s. 6d.
- Questions for Examinations in English Literature.** By Rev.
W. W. Skeat, Prof. of Anglo-Saxon at Cambridge University. 2s. 6d.
- Drawing Copies.** By P. H. Delamotte. Oblong 8vo. 12s. Sold
also in parts at 1s. each.
- Poetry for the School-room.** New Edition. Fcap. 8vo. 1s. 6d.
- Geographical Text-Book; a Practical Geography.** By M. E. S.
12mo. 2s.
The Blank Maps done up separately, 4to. 2s. coloured.
- Loudon's (Mrs.) Entertaining Naturalist.** New Edition. Revised
by W. S. Dallas, F.L.S. 5s.
- **Handbook of Botany.** New Edition, greatly enlarged by
D. Wooster. Fcap. 2s. 6d.
- The Botanist's Pocket-Book.** With a copious Index. By W. R.
Hayward. 3rd Edit. revised. Crown 8vo. Cloth limp. 4s. 6d.
- Experimental Chemistry, founded on the Work of Dr. Stöckhardt.**
By C. W. Heaton. Pest 8vo. 5s.
- Double Entry Elucidated.** By B. W. Foster. 12th Edit. 4to.
3s. 6d.
- A New Manual of Book-keeping.** By P. Crellin, Accountant.
Crown 8vo. 3s. 6d.

Picture School-Books. In Simple Language, with numerous
Illustrations. Royal 16mo.

School Primer. 6d.—School Reader. By J. Tilleard. 1s.—Poetry Book
for Schools. 1s.—The Life of Joseph. 1s.—The Scripture Parables. By the
Rev. J. E. Clarke. 1s.—The Scripture Miracles. By the Rev. J. E. Clarke.
1s.—The New Testament History. By the Rev. J. G. Wood, M.A. 1s.—The
Old Testament History. By the Rev. J. G. Wood, M.A. 1s.—The Story of
Bunyan's Pilgrim's Progress. 1s.—The Life of Christopher Columbus. By
Sarah Crompton. 1s.—The Life of Martin Luther. By Sarah Crompton. 1s.

BOOKS FOR YOUNG READERS.

A Series of Reading Books designed to facilitate the acquisition of the power of Reading by very young Children. In 8 vols. limp cloth, 8d. each.

- | | |
|--|--|
| The Cat and the Hen. Sam and his Dog Redleg.
Bob and Tom Lec. A Wreck. | } Suitable
for
Infants. |
| The New-born Lamb. The Rosewood Box. Poor
Fan. Sheep Dog. | |
| The Story of Three Monkeys. | } Suitable
for
Standards
I. & II. |
| Story of a Cat. Told by Herself. | |
| The Blind Boy. The Mute Girl. A New Tale of
Babes in a Wood. | |
| The Dey and the Knight. The New Bank Note.
The Royal Visit. A King's Walk on a Winter's Day. | |
| Queen Bee and Busy Bee. | |
| Gull's Cragg. | |
| A First Book of Geography. By the Rev. C. A. Johns.
Illustrated. Double size, 1s. | |

BELL'S READING-BOOKS.

FOR SCHOOLS AND PAROCHIAL LIBRARIES.

The popularity which the 'Books for Young Readers' have attained is a sufficient proof that teachers and pupils alike approve of the use of interesting stories, with a simple plot in place of the dry combination of letters and syllables, making no impression on the mind, of which elementary reading-books generally consist.

The Publishers have therefore thought it advisable to extend the application of this principle to books adapted for more advanced readers.

Now Ready. Post 8vo. Strongly bound.

- | | |
|--|--|
| Grimm's German Tales. (Selected.) 1s. | } Suitable
for
Standards
II. & III. |
| Andersen's Danish Tales. (Selected.) 1s. | |
| Great Englishmen. Short Lives for Young Children. 1s. | |
| Lamb's Tales from Shakespeare. (Selected.) 1s. | |
| Edgeworth's Tales. A Selection. 1s. | |
| Friends in Fur and Feathers. By Gwynfryn. 1s. | } Standards
IV. & V. |
| Parables from Nature. (Selected.) By Mrs. Gatty. 1s. | |
| Masterman Ready. By Capt. Marryat. (Abgd.) 1s. 6d. | |
| Settlers in Canada. By Capt. Marryat. (Abdg.) 1s. 6d. | |
| Gulliver's Travels. (Abridged.) 1s. | |
| Robinson Crusoe. 1s. 6d. | |
| Marie; or, Glimpses of Life in France. By A. R. Ellis.
1s. | |
| Poetry for Boys. By D. Munro. 1s. | } Standard
VI. |
| Southey's Life of Nelson. (Abridged.) 1s. | |
| Life of the Duke of Wellington, with Maps and Plans. 1s. | |
| The Romance of the Coast. By J. Runciman.
[In the press.] | |

LONDON:

Printed by STRANGEWAYS & SONS, Tower Street, Upper St. Martin's Lane.





Physics

Mech

B

266462

Author ..Besant, William Henry.....

TitleElementary hydrostatics.....

**University of Toronto
Library**

**DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET**

Acme Library Card Pocket
Under Pat. "Ret. Ind. File"
Made by LIBRARY BUREAU

